



King's Research Portal

DOI:

[10.1093/ijnp/pyw091](https://doi.org/10.1093/ijnp/pyw091)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Fountoulakis, K. N., Young, A., Yatham, L., Grunze, H., Vieta, E., Blier, P., Moeller, H. J., & Kasper, S. (2016). The International College of Neuropsychopharmacology (CINP) Treatment Guidelines for Bipolar Disorder in Adults (CINP-BD-2017), Part 1: Background and Methods of the Development of Guidelines. *International Journal of Neuropsychopharmacology*. <https://doi.org/10.1093/ijnp/pyw091>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



REVIEW

The International College of Neuropsychopharmacology (CINP) Treatment Guidelines for Bipolar Disorder in Adults (CINP-BD-2017), Part 1: Background and Methods of the Development of Guidelines

Konstantinos N. Fountoulakis, MD; Allan Young, MD; Lakshmi Yatham, MD; Heinz Grunze, MD; Eduard Vieta, MD; Pierre Blier, MD; Hans Jurgen Moeller, MD; Siegfried Kasper, MD

3rd Department of Psychiatry, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece (Dr Fountoulakis); Centre for Affective Disorders, Institute of Psychiatry, Psychology and Neuroscience, King's College, London, UK (Dr Young); Department of Psychiatry, University of British Columbia, Mood Disorders Centre of Excellence, Djavad Mowafaghian Centre for Brain Health, Vancouver, Canada (Dr Yatham); Paracelsus Medical University, Salzburg, Austria (Dr Grunze); Hospital Clinic, Institute of Neuroscience, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Catalonia, Spain (Dr Vieta); The Royal Institute of Mental Health Research, Department of Psychiatry, University of Ottawa, Ottawa, Canada (Dr Blier); Psychiatric Department, Ludwig Maximilians University, Munich, Germany (Dr Moeller); Department of Psychiatry and Psychotherapy, Medical University Vienna, MUV, AKH, Vienna, Austria (Dr Kasper).

Correspondence: Konstantinos N. Fountoulakis, MD, 6, Odysseos str (1st Parodos Ampelou str.), 55535 Pylaia Thessaloniki, Greece (kfount@med.auth.gr).

Abstract

Background: This paper includes a short description of the important clinical aspects of Bipolar Disorder with emphasis on issues that are important for the therapeutic considerations, including mixed and psychotic features, predominant polarity, and rapid cycling as well as comorbidity.

Methods: The workgroup performed a review and critical analysis of the literature concerning grading methods and methods for the development of guidelines.

Results: The workgroup arrived at a consensus to base the development of the guideline on randomized controlled trials and related meta-analyses alone in order to follow a strict evidence-based approach. A critical analysis of the existing methods for the grading of treatment options was followed by the development of a new grading method to arrive at efficacy and recommendation levels after the analysis of 32 distinct scenarios of available data for a given treatment option.

Conclusion: The current paper reports details on the design, method, and process for the development of CINP guidelines for the treatment of Bipolar Disorder. The rationale and the method with which all data and opinions are combined in order to produce an evidence-based operationalized but also user-friendly guideline and a specific algorithm are described in detail in this paper.

Keywords: Bipolar Disorder; anticonvulsants; antidepressants; antipsychotics; evidence-based guidelines; lithium; mania; bipolar depression; mood stabilizers; treatment

Received: June 29, 2016; Revised: September 17, 2016; Accepted: October 20, 2016

© The Author 2016. Published by Oxford University Press on behalf of CINP.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Introduction

General Background, Disclosure, and Aim

Treatment guidelines are becoming an ever more important part of medical reality, especially since the translation of research findings to everyday clinical practice is becoming increasingly difficult with the accumulation of complex and often conflicting research findings that are thereafter also included in metaanalysis. Guidelines aim to assist clinicians but also policymakers to arrive at decisions concerning the treatment and care of patients. They set the standard of care and training for health professionals and they also identify priority areas for further research, since they are based primarily on the available evidence, but also in areas where evidence is not available, on expert opinion (Fountoulakis, 2015i).

In the field of Bipolar Disorder (BD), accumulated knowledge is often complex, confusing, and in many instances contrasts with the beliefs and practices that appear to have been set in stone in psychiatric culture and training for the last few decades.

To fulfil this need for expert translation of research findings into clinical practice for the benefit of patients, the International College of Neuropsychopharmacology (CINP) launched an effort to critically appraise the literature and provide guidance to clinicians in the form of a precise treatment algorithm. It is hoped that this algorithm for the treatment of BD will help the clinician to follow the state-of-the-art evidence, thus enabling their clinical practice to be based on an informed decision-making process. This guideline has been commissioned by the CINP, and the workgroup consisted of experts with extensive research and clinical experience in the field of BDs. There was no funding from any source for the development of the guidelines and the activities of the workgroup.

All the members of the workgroup were psychiatrists who are in active clinical practice and were selected according to their expertise and with the aim to cover a multitude of some different cultures. All of them were involved in research and other academic activities, and therefore it is possible that through such activities some contributors have received income related to medicines discussed in this guideline. All conflicts of interest are mentioned at the end of this paper, which is the introductory paper to the CINP BD guidelines. It should also be noted that some drugs recommended in the guideline may not be available in all countries, and labeling and dosing might vary.

The aim of the current endeavor was to develop a guideline and precise algorithm for treatment of BD in adults for use in primary and secondary care. Children, adolescents, and the elderly are not the focus of this guidance. The guideline and algorithm have been developed after a complete review of the literature and with the use of stringent criteria. Both the guideline and the precise algorithm try to balance research vs clinical wisdom but give primacy to the available evidence.

To comply with the journal's word limit for manuscripts and for easy readability, the CINP guidelines have been organized and presented as a series of 4 distinct papers. This paper is the first of this series and will cover the general background of the guideline and algorithm, that is, the historical perspective and general clinical and treatment issues followed for the development of the guideline and the algorithm. The second paper summarizes, classifies, and grades the treatment data on BD while the third paper includes the guideline and the treatment algorithm themselves. The fourth and final paper addresses the unmet needs and areas that should be the focus of attention and specific research in the future.

Historical Perspective

Depression and bipolarity were mentioned in Eber's papyrus in ancient Egypt around 3000 BC (Okasha and Okasha, 2000) and in the Hippocratic texts. Plato (424–348 BC) and Aristotles (384–322 BC) further elaborated on the concept and Aristotle was the first to describe accurately the affections of desire, anger, fear, courage, envy, joy, hatred, and pity. Later, Galen (131–201 AD), Themison of Laodicea (1st century BC) and Aretaeus of Cappadocia (2nd century AD) as well as Arab scholars and especially Avicenna (980–1037) further elaborated on the concept of mood disorders (Fountoulakis, 2015b).

Jean-Philippe Esquirol (1772–1840) was the first to clearly point out that melancholia was a disorder of the mood with “partial insanity” (monomania) and used the word “lypemia.” Finally, Jean-Pierre Falret (1794–1870) and Jules Gabriel Francois Baillarger (1809–1890) established the connection between depression and mania and gave it the name of “folie circulaire” or “folie à double forme,” but it was Emil Kraepelin (1856–1926) who established manic-depressive illness as a distinct nosological entity and separated it from schizophrenia on the basis of heredity, longitudinal follow-up, and a supposed favorable outcome (Kraepelin, 1921). His pupil Wilhelm Weygandt (1870–1939) published the first textbook on mixed clinical states (Weygandt, 1899). Following a similar line of thinking, and in spite of some major objections to the Kraepelinian approach, Karl Jaspers (1883–1969) described aspects of mixed depressive states that he named “querulant mania,” “nagging depression,” or “wailing melancholia” (Jaspers, 1913), while Eugene Bleuler coined the term “affective illness” and by this he broadened the concept of manic-depression.

In 1957 Karl Leonhard (1902–1988) proposed that the term “bipolar disorder” should replace manic-depression, and he also made a distinction between monopolar (unipolar depression) and bipolar illness (Leonhard, 1957a, 1957b; Leonhard, 1979).

In 1870, Silas Weir Mitchell (1829–1914) was the first to recommend lithium as an anticonvulsant, hypnotic, and as medication for “general nervousness” (Mitchell, 1870, 1877). In 1871, William Alexander Hammond (1828–1900) was probably the first to prescribe a modern and effective psychotropic agent, and this was lithium (Mitchell and Hadzi-Pavlovic, 2000). Carl Lange (1834–1900) and Frederik Lange (1842–1907) had used lithium in the treatment of depression since 1886 (Lenox and Watson, 1994). However, in spite of encouraging results, by the turn of the 20th century, the “brain gout” theory of mood disorders disappeared as a medical entity and the use of lithium in psychiatry was abandoned.

In 1949 John Cade (1912–1980) reported positive results from the treatment of 10 acutely manic patients (Cade, 1949, 2000); however, 2 years later he reported the first death caused by lithium toxicity in a patient whose bipolar illness otherwise responded extremely well to treatment. Later, Mogens Schou (1918–2005) undertook a randomized controlled trial of lithium in mania (Schou et al., 1954; Bech, 2006), and eventually the efficacy of lithium during the maintenance phase was established (Gershon and Yuwiler, 1960; Baastrup, 1964; Baastrup and Schou, 1967; Angst et al., 1969, 1970; Baastrup et al., 1970; Schou et al., 1970; Johnstone et al., 1988; Schioldann, 1999; Mitchell and Hadzi-Pavlovic, 2000; Bech, 2006; Schioldann, 2006; Schioldann, 2011).

Valproate was introduced in 1966 as an anticonvulsant (Lambert et al., 1966) and later carbamazepine (Okuma et al., 1979) followed. Neuroleptics were introduced by Jean Delay

(1907–1987) and Pierre Deniker (1917–1999) in 1955, and probably many of their patients were suffering from acute mania or schizoaffective disorder (Delay and Deniker, 1955). In 1958 Roland Kuhn (1912–2005) reported on the efficacy of the first antidepressant, imipramine (Kuhn, 1958).

There were several reports in the 1970s suggesting that in bipolar depression the use of antidepressants might induce mania, mixed episodes, and rapid cycling (Wehr and Goodwin, 1987; Wehr et al., 1988). In 1994 the first detailed operational treatment guidelines were published by the American Psychiatric Association and after 2000, systematic industry-sponsored studies of second generation antipsychotics and haloperidol were performed. Also, during this period the first meta-analytic studies emerged, and the evidence-based medicine principles gained ground in treatment recommendations.

Clinical Description

While the basic conception of BD suggested that it is characterized by the alternation of manic and depressive episodes with a return to the premorbid level of functioning between the episodes and to favorable outcome compared with schizophrenia (Kraepelin, 1921), today we know that this is not always the case (Tohen et al., 1990; Grande et al., 2016). Not only BD is a much more complex disorder than this, but also the outcome varies. The most prominent clinical facets are shown in Table 1 (Fountoulakis, 2015a, 2015c, 2015d, 2015e, 2015j, 2015n).

The fact that often the correct diagnosis is made only after 8 to 10 years have passed because the first episode is psychotic-like or depressive and the correct diagnosis can be made only after a manic or a mixed episode emerges (Angst, 2007) is especially problematic. It has been estimated that more than one-half of hospitalized patients originally manifesting a depressive episode will turn out to be bipolars in the next 20 years (Angst et al., 2005a). It is of utmost importance for both clinicians and researchers to create a biographical chart with the patient's course over time and including any important event in the developmental history of the patient and emphasizing the main events and hallmarks of his/her life and his/her full psychiatric and medical history. Such a chart clarifies both the diagnosis and the course of the disease and also the response to therapeutic interventions, since any delay in the proper diagnosis also delays proper treatment (Altamura et al., 2010; Drancourt et al., 2013).

Table 1. List of the Multiple Clinical Aspects of Manic-Depressive Illness

1. Manic episodes
2. Depressive episodes
3. Mixed episodes
4. Subthreshold manic symptoms
5. Subthreshold depressive symptoms
6. 'Mixed' states and 'roughening'
7. Mood lability/cyclothymia/'personality-like' behavior
8. Predominant polarity
9. Frequency of episodes/rapid cycling
10. Psychotic features
11. Neurocognitive disorder
12. Functional deficit and disability
13. Drug/alcohol abuse
14. Comorbid anxiety and other mental disorders
15. Self-destructive behavior and suicidality

In terms of individual symptoms, fatigue and psychomotor retardation dominate the clinical picture in 75% of patients during acute bipolar depression. Irritability is present in almost 75% of patients (Winokur et al., 1969), delusions are present in 12 to 66% (Winokur et al., 1969; Carlson and Strober, 1978; Rosenthal et al., 1980; Black and Nasrallah, 1989), and hallucinations in 8 to 50% (Winokur et al., 1969; Carlson and Strober, 1978; Rosenthal et al., 1980; Black and Nasrallah, 1989; Baethge et al., 2005). Psychotic features seem to constitute a stable trait that tends to repeat itself across episodes (Helms and Smith, 1983; Nelson et al., 1984; Aronson et al., 1988a, 1988b). Depending of the study sample composition, changes in appetite for food are seen in almost all patients (Winokur et al., 1969), with one-fourth manifesting overeating and one-fourth losing significant weight (Casper et al., 1985). Almost all bipolar depressed patients experience some kind of sleep problem (Winokur et al., 1969; Casper et al., 1985). A subgroup of bipolar depressed patients (up to 25%) often exhibit excessive sleep and have difficulty getting up in the morning (Winokur et al., 1969). Decreased sexual desire is seen in more than 75% of patients (Winokur et al., 1969; Casper et al., 1985) and concerns both sexes. Approximately two-thirds of bipolar depressed patients present with multiple physical pains and complaints (e.g., headache, epigastric pain, precordial distress, etc.) in the absence of any physical illness, especially in primary care (Winokur et al., 1969).

Euphoria is observed in 30 to 97% of acutely manic patients (Clayton and Pitts, 1965; Winokur et al., 1969; Beigel and Murphy, 1971; Carlson and Goodwin, 1973; Taylor and Abrams, 1973; Winokur and Tsuang, 1975; Abrams and Taylor, 1976; Leff et al., 1976; Loudon et al., 1977; Taylor and Abrams, 1977; Cassidy et al., 1998a), while unrestrained and expansive mood is seen in 44 to 66% (Taylor and Abrams, 1973, 1977; Loudon et al., 1977). Patients are dissatisfied and intolerant and the vast majority manifest mood lability and instability (42 to 95%) (Winokur et al., 1969; Carlson and Goodwin, 1973; Abrams and Taylor, 1976; Loudon et al., 1977; Taylor and Abrams, 1977; Cassidy et al., 1998a). Irritability is also very frequent (51–100%) (Winokur et al., 1969; Carlson and Goodwin, 1973; Taylor and Abrams, 1973, 1977; Winokur and Tsuang, 1975; Abrams and Taylor, 1976; Loudon et al., 1977; Cassidy et al., 1998a; Serretti and Olgiati, 2005). However, even significant depressive symptoms are experienced by as many as 29 to 100% of acutely manic patients (Winokur et al., 1969; Beigel and Murphy, 1971; Kotin and Goodwin, 1972; Carlson and Goodwin, 1973; Murphy and Beigel, 1974; Loudon et al., 1977; Prien et al., 1988; Cassidy et al., 1998a; Bauer et al., 2005).

Accelerated psychomotor activity is observed in the vast majority of patients (56–100%) (Winokur et al., 1969; Carlson and Goodwin, 1973; Taylor and Abrams, 1973; Abrams and Taylor, 1976; Leff et al., 1976; Loudon et al., 1977; Carlson and Strober, 1978; Cassidy et al., 1998a, 1988b; Cassidy et al., 1998a; Serretti and Olgiati, 2005) and pressured speech in almost all patients (Clayton and Pitts, 1965; Winokur et al., 1969; Carlson and Goodwin, 1973; Taylor and Abrams, 1973; Abrams and Taylor, 1976; Leff et al., 1976; Loudon et al., 1977; Carlson and Strober, 1978; Cassidy et al., 1998b; Serretti and Olgiati, 2005); hypersexuality is present in 25 to 80% of patients with 23 to 33% of them having significant sexual exposure (Allison and Wilson, 1960; Clayton and Pitts, 1965; Winokur et al., 1969; Carlson and Goodwin, 1973; Taylor and Abrams, 1973, 1977; Abrams and Taylor, 1976; Leff et al., 1976; Loudon et al., 1977; Carlson and Strober, 1978). Decreased need for sleep (hyposomnia) is present in 63 to 100% of patients (Clayton and Pitts, 1965; Winokur et al., 1969; Leff et al., 1976; Loudon et al., 1977; Carlson and Strober, 1978;

Cassidy et al., 1998b; Serretti and Olgiati, 2005) and psychotic features in 33 to 96% of patients (Winokur et al., 1969; Carlson and Strober, 1978; Rosenthal et al., 1980; Black and Nasrallah, 1989).

Overall, psychotic features are so common that acute mania should be considered primarily a psychotic state (Koukopoulos, 2006). Delusions are present in 24 to 96% of manic patients, and it is interesting that persecutory ideas are equally frequent with delusions of grandiose (Bowman and Raymond, 1932; Rennie, 1942; Astrup et al., 1959; Clayton and Pitts, 1965; Winokur et al., 1969; Beigel and Murphy, 1971; Carlson and Goodwin, 1973; Taylor and Abrams, 1973, 1977; Murphy and Beigel, 1974; Abrams and Taylor, 1976; Leff et al., 1976; Loudon et al., 1977; Carlson and Strober, 1978; Rosenthal et al., 1980; Winokur, 1984; Black and Nasrallah, 1989; Serretti et al., 2002; Keck et al., 2003; Goodwin and Jamison, 2007). Hallucinations are less frequent and present in 13 to 66% of cases; they can either be congruent or noncongruent, with auditory, visual, and olfactory ones being almost equally frequent (Lange, 1922; Bowman and Raymond, 1932; Astrup et al., 1959; Winokur et al., 1969; Taylor and Abrams, 1973, 1977; Abrams and Taylor, 1976; Carlson and Strober, 1978; Rosenthal et al., 1980; Winokur, 1984; Black and Nasrallah, 1989; Serretti et al., 2002; Keck et al., 2003; Goodwin and Jamison, 2007).

Psychotic symptoms in BD are predictive of a more detrimental course, including a higher rate of rehospitalizations (Caetano et al., 2006; Ozyildirim et al., 2010).

Almost one-third of acutely manic patients are “confused” and 46 to 75% are violent (Carlson and Goodwin, 1973; Taylor and Abrams, 1973, 1977; Abrams and Taylor, 1976; Cassidy et al., 1998b). The term confused refers to manic disorganization and not to organic drop in the level of consciousness. As many as 14 to 56% of patients manifest severe regression, catatonia, posturing, and negativism, often making differential diagnosis from schizophrenia difficult (Lange, 1922; Carlson and Goodwin, 1973; Taylor and Abrams, 1973, 1977; Carlson and Strober, 1978; Abrams and Taylor, 1981; Braunig et al., 1998; Kruger et al., 2003), and 10 to 20% have fecal incontinence (Taylor and Abrams, 1973, 1977; Abrams and Taylor, 1976). A summary of the frequencies of appearance of various symptoms during the two different acute phases of the illness is shown in Table 2.

Formally, those episodes with manic symptoms but less pronounced in terms of severity and with a shorter duration are labeled hypomanic. Hypomania is much more common than mania (Angst, 1998), but its recognition is mostly achieved mainly by interviewing significant others and not the patient. Hypomanic episodes cause mild or no impairment at all, and on the contrary, in some cases, they may even contribute to success in business, leadership roles, and the arts. Psychotic symptoms are less frequent (around 20%) in comparison to full-blown manic episodes, but they do occur (Mazzarini et al., 2010).

Mixed episodes are defined as the coexistence of both depressive and manic symptoms; however, the term was abandoned with DSM-5, which includes mixed features as a specifier only. The DSM-5 demands the presence of a full-blown episode of either pole together with at least 3 symptoms of the opposite pole being present in order to allow the label of “mixed features” specifier.

It is reported that in 69.6% of cases the course resembles that of a recurrent episodic illness, while in 25% of cases there is a chronic course without clear remissions between episodes. In only 5.4% is there a single episode of mania. Suicidal ideation is present in 78.6% of patients at some time in their life. Only around 5% of BD patients have chronic mania (Akiskal, 2000).

Table 2. Summary of the Frequencies of Appearance of Various Symptoms during the Two Different Acute Phases of BD

Symptom	Episodes	
	Manic	Depressive
Euphoria	30–97%	
Expansive mood	44–66	
Depressive symptoms	29–100%	100%
Mood lability	42–95%	
Irritability	51–100%	75%
Psychomotor retardation		75%
Psychomotor acceleration	56–100%	
Pressured speech	100%	
Psychotic features	33–96%	
Delusions	24–96%	12–66%
Hallucinations	13–66%	8–50%
Weight loss		25%
Weight gain		25%
Hyposomnia	63–100%	
Oversleeping		25%
Loss of libido		25%
Hypersexuality	25–80%	
Significant sexual exposure	23–33%	
Confused	33%	
Violent	46–75%	
Regression, catatonia etc.	14–56%	
Fecal incontinence	10–20%	
Physical complains		66%

Karl Leonhard was the first to report the presence of a predominant polarity with 17.9% of patients having a manic- and 25.6% having a depressive-predominant polarity (Leonhard, 1963). The concept was further formulated by Jules Angst (1978) and Carlo Perris (Perris and d’Elia, 1966a, 1966b) and has recently been utilized for long-term prognosis and to assist clinicians in long-term therapeutic design (Quitkin et al., 1986; Judd et al., 2003; Colom et al., 2006). The most reliable definition of predominant polarity demands that at least two-thirds of episodes belong to one of the poles (Colom et al., 2006; Rosa et al., 2008; Garcia-Lopez et al., 2009; Mazzarini et al., 2009; Tohen et al., 2009; Vieta et al., 2009; Nivoli et al., 2011; Baldessarini et al., 2012; Pacchiarotti et al., 2013a; Carvalho et al., 2014a, 2014b).

Somewhere between 15% and 50% of BD patients are reported to manifest some type of seasonal variation of symptomatology (Hunt et al., 1992; Faedda et al., 1993; Goikolea et al., 2007; Shand et al., 2011). Two opposing seasonal variations have been described: fall-winter depression with or without spring-summer mania or hypomania; and spring-summer depression with or without fall-winter mania or hypomania (Faedda et al., 1993). Most studies support the first subtype (Walter, 1977; Parker and Walter, 1982; Mulder et al., 1990; Peck, 1990; Partonen and Lonnqvist, 1996; Clarke et al., 1999; Lee et al., 2007; Murray et al., 2011).

The concept of rapid cycling appeared for the first time in the 70s in a landmark paper by Dunner and Fieve (1974). In general the classic rapid-cycling includes cycles with duration of weeks to months. Ultra-rapid cycling is reported when mood cycling has frequency of weeks to days, and ultradian cycling when there is significant mood variation within a 24-hours period (Kramlinger and Post, 1996). Other terms include ultra-ultra rapid and ultradian rapid and refer to weekly or daily cycling, which is not uncommon in BD patients (Kramlinger and Post, 1996). Most studies suggest a 5 to 33.3% up-to-1-year prevalence

(Kukopulos et al., 1980; Nurnberger et al., 1988; Coryell et al., 1992; Schneck et al., 2004, 2008; Azorin et al., 2008; Cruz et al., 2008; Garcia-Amador et al., 2009; Lee et al., 2010) and 25.8 to 43% lifetime prevalence (Dittmann et al., 2002; Coryell et al., 2003; Yildiz and Sachs, 2004; Hajek et al., 2008; Lee et al., 2010).

In terms of neurocognitive function, the literature suggests that the neurocognitive deficit in BD patients concerns almost all domains and phases of the illness with only a few exceptions. Its magnitude is at the severe range during the acute episodes and at the medium range during euthymia, while the origin of the deficit remains unclear. In terms of neurocognitive function, BD patients do quantitatively better than patients with schizophrenia, but the qualitative pattern of the deficit is similar in the 2 disorders. There are no clear differences between BD subtypes. The deficit is present early in the course of the disorder. At least in some patients it might emerge before the onset of the first mood episode, and in the majority of patients it progresses probably in relationship with the manifestation of psychotic symptoms. The verbal memory and executive function deficit probably constitute endophenotypes, while the role of medication as a causative factor is limited (Tsitsipa and Fountoulakis, 2015; Cullen et al., 2016).

Finally, in contrast to the original conceptualization of BD by Emil Kraepelin a century ago, unfortunately it seems that only a minority of BD patients achieve complete functional recovery (Goldberg et al., 1995a, 1995b; Keck et al., 1998; Strakowski et al., 1998; Daban et al., 2006; Martinez-Aran et al., 2007; Mur et al., 2007).

Classification

ICD and DSM include BD as a diagnostic entity but with significant differences between them (Fountoulakis, 2015h). It is important to note that almost all the research literature follows the DSM classification, while almost all countries worldwide have the obligation to use the ICD in their official documents, including hospital records, etc. The ICD-10-CM helps to bridge these 2 different classification systems for administration purposes. In ICD-10 (WHO, 1992, 1994), BD is included in the chapter on mood (affective) disorders (F30-F39). While in previous editions of the DSM, both unipolar and bipolar disorders were grouped under the chapter on mood disorders, on the contrary in DSM-5 (American Psychiatric Association, 2013), BDs were separated from unipolar depression. The “bipolar” chapter includes BD and cyclothymic disorder, while the “depression” chapter includes disruptive mood dysregulation disorder, major depressive disorder, persistent depressive disorder (dysthymia), and premenstrual dysphoric disorder. Both chapters include “unspecified,” “other,” and “due to” categories.

Another important difference between the 2 classification systems is that ICD requires the presence of at least 2 episodes of pathological disturbance of mood while DSM does not. DSM recognizes the presence of 2 subtypes of BD, that is, of BD-I (BD with manic episodes) and BD-II (BD with hypomanic but not manic episodes). BD-II is not part of the ICD-10 diagnostic list, which accepts hypomania as a diagnostic entity (F30.0), but it is considered simply a low-severity mania.

In ICD-10 a mixed affective episode (F38.0) is defined as an affective episode of at least 2 weeks duration that is characterized by either a mixture or a rapid alternation (usually within a few hours) of hypomanic, manic, and depressive symptoms. In DSM-5 a radical change was the abolishment of the concept of mixed episodes. In previous versions of the DSM, mixed episodes were defined as the coexistence of full-blown manic and

depressive episodes simultaneously. Although such a coexistence is rather rare, almost one-third of patients recruited in pharmaceutical trials of acute mania were diagnosed as mixed. Thus there exists ample data, although neither properly analyzed nor published. Instead of the diagnosis of mixed episodes, DSM-5 introduced the mixed features specifier concept. According to this, a mood episode (either manic or depressed) has mixed features if at least 3 criteria of the opposite pole (from a specific list) coexist. It is important to note that according to DSM-5, mixed features can also be attributed to a unipolar major depressive episode without changing the diagnosis to BD.

Another important change in the DSM-5 is the introduction of the anxious distress specifier, which demands the presence of at least 2 criteria from a list of 5 (tension, restlessness, concentration difficulties, worry, fear of losing control).

The ICD-10 classification accepts the presence of “somatic syndrome,” which seems analogous but it is not identical to “melancholic features” of DSM-5 (Fountoulakis et al., 1999). The atypical features, rapid cycling, and anxious distress are described in DSM-5 but not in ICD-10. Also, ICD-10 does not differentiate psychotic symptoms into mood congruent vs mood incongruent. The other specifiers, catatonia, peripartum onset, and seasonal pattern, are not included in the ICD-10 either. It is also important to note that ICD-10 recognizes catatonia only in the frame of schizophrenia, while DSM-5 uses this specifier also in affective disorders.

There is an issue concerning the diagnosis of cases with sub-threshold manic symptoms or long-lasting hyperthymia. While the traditional bipolar vs unipolar distinction is widely used and adopted by classification systems, it is doubtful whether it can capture the essence of the huge heterogeneity observed in mood disorders and their dynamic nature with frequent switches and changes in the clinical profile. The greatest disadvantage of both classification systems is that they perform better (and focus) when interepisodic remission is present; instead, the everyday real-life patient is more likely to suffer from a chronic disorder with residual and mixed symptoms. The term spectrum was first used in psychiatry in 1968 for the schizophrenia spectrum (Kety et al., 1968).

The proposed mood spectrum models unify categorical classification, which is essential, with a dimensional view, which is true to nature; both are needed and both are empirically testable. Today the term bipolar spectrum is mainly used in 2 complementary senses: (1) a spectrum of severity, which embraces psychotic and nonpsychotic major and minor BDs (including bipolar dysthymia, recurrent brief and minor depressions), cyclothymic disorders, hypomania and, at its broadest, even borderline disorders and cyclothymic temperament; (2) a proportional mood spectrum, which considers the 2 components, mania and depression, on the level of major and minor mood disorders. This proportional model is an extension of Kleist's concept of BD as a combination of the 2 monopolar disorders of depression and mania (Kleist, 1937). Thus these 2 approaches to spectrum reflect 2 distinct continua: from normal to pathological and from unipolar to bipolar.

An important part of the bipolar spectrum is cyclothymic disorder, which is considered to be an attenuated form of BD. Their behavior is characterized by the alternation of extremes (Akiskal et al., 1977). Depending on the threshold of traits used in determining the presence of hyperthymia, cyclothymic patients may constitute 10 to 20% of those with major depressive disorder. Also, cyclothymia is often a prodromal of BD (Akiskal et al., 1979). Another important part of the bipolar spectrum are those patients who experience an antidepressant-induced switch.

Thus, many patients with so-called unipolar depression are actually pseudounipolar.

Some authors suggest that a significant part of the literature consists mostly of expert opinion overemphasizing various links between bipolar and unipolar mood disorders and personality disorders (Paris et al., 2007; Patten and Paris, 2008). Recently, the first solid international epidemiological data in support of the bipolar spectrum have been published (Merikangas et al., 2007, 2011; Angst et al., 2010). According to these authors there is a direct association between increasingly restrictive definitions of BD and indicators of clinical severity, including symptom severity, role impairment, comorbidity, suicidality, and treatment. For example, the proportion of mood episodes rated as clinically severe increased from 42.5% for subthreshold BD to 68.8% for BD-II to 74.5% for BD-I. However, since clinical diagnosis and severity share confounding factors and definitions overlap, it is also important to note that these studies also showed that the proportion of cases reporting severe role impairment ranged from 46.3% for subthreshold BD to 57.1% for BD-I (Merikangas et al., 2011).

On the basis of both epidemiological data and clinical wisdom, a limited number of models reflecting the structure of the bipolar spectrum have been proposed. The first effort was a dimensional concept (from normal to pathological) proposed by Kretschmer in 1921 for schizophrenia (schizothymic-schizoid-schizophrenic) and for affective disorders (cyclothymic temperament-cycloid 'psychopathy'-manic-depressive disorder). Bleuler suggested a similar concept in 1922. In 1977 Akiskal proposed a cyclothymic-bipolar spectrum (Akiskal et al., 1977). A simple model system was introduced in 1978 by Jules Angst (1978; Angst et al., 1978), who used the the following codes: M for severe mania, D for severe depression (unipolar depression), m for less severe mania (hypomania), and d for less severe depression. In 1981 Gerald Klerman suggested a mania spectrum (Klerman, 1981, 1987) and in the late 1990s Akiskal proposed 6 subtypes, some of which are further subdivided according to their unique clinical features. A summary of his proposed subtype schema is as follows (Akiskal and Pinto, 1999; Akiskal and Benazzi, 2005; Ng et al., 2007; Fountoulakis, 2008).

Epidemiology

In the last few decades there has been an increasing interest in psychiatric epidemiology. For BD, a point that plays a major role in the estimation of the prevalence rates is the definition of hypomania and of mixed, irritable, or dysphoric forms of manic episodes. This is further complicated by the presence of inaccurate recall and the low sensitivity of the interview instruments concerning subthreshold symptomatology and nonclassical clinical pictures (Kessler et al., 1997a).

A number of important studies exist and provide important but inconclusive information. The Amish study (Egeland and Hostetter, 1983; Egeland et al., 1983; Hostetter et al., 1983) reported similar prevalence rates between unipolar depression and bipolar illness and also similar rates between genders. It is impressive that 79% of patients with BD-I were previously diagnosed as suffering from schizophrenia. The Epidemiological Catchment Area study (ECA) (Eaton et al., 1981; Regier et al., 1984, 1988, 1993; Bourdon et al., 1992) reported a lifetime prevalence of 0.8% for BD-I (0.3–1.2%) and an annual prevalence of 0.6% (0.2–1%) with similar prevalence for males and females. The annual incidence was 0.4% (0.1–0.6%) of cases, which corresponds to approximately 3.2 (0.8–4.8) per 100 000 residents. The median age at onset was 18 years. A reanalysis of the ECA data

with the addition of subthreshold bipolarity produced a total lifetime prevalence of 6.4% with 0.5% being a lifetime prevalence of BD-II (Judd and Akiskal, 2003). The National Comorbidity Survey (NCS) (Kessler et al., 1993, 1994a, 1994b, 1995, 1996, 1997b; Blazer et al., 1994; Wittchen et al., 1994; Warner et al., 1995; Kendler et al., 1996; Magee et al., 1996) reported a lifetime prevalence of 1.7% for BD-I and an annual prevalence of 1.3% with similar prevalence for males and females. The median age at onset was 21 years. The NCS-R (Kessler et al., 2004, 2005a, 2005b, 2012a, 2012b; Kessler and Merikangas, 2004; Merikangas et al., 2007; Angst et al., 2010; Nierenberg et al., 2010) reported a lifetime prevalence of 1.0% for BD-I and an annual prevalence of 0.6% with again similar prevalence for males and females. The median age at onset was 19 years. For BD-II the lifetime prevalence was 1.1% and the annual prevalence was 0.8% with similar prevalence for males and females. The median age at onset was 20 years. There was a small difference between males and females in the BD-II rates, with female rates being slightly higher. The Cross National Collaboration Group study included data from 7 countries (US, Canada, Puerto Rico, Germany, Taiwan, South Korea and New Zealand) (Weissman et al., 1996) and reported variable rates for different countries, but overall the rates seemed moderately consistent cross-nationally. The Zurich study (Angst et al., 1984, 2005b; Wicki and Angst, 1991) reported an annual prevalence of BD-I of 0.7% and a lifetime prevalence for the bipolar spectrum of 5.5%. The Nottingham study (Brewin et al., 1997) reported the 2-year incidence rate reported for BD was 0.005%, which corresponds to an annual incidence of 2.5/100 000. The Netherlands study (Bijl et al., 2002; Regeer et al., 2002; ten Have et al., 2002) suggested a lifetime prevalence of BD equal to 2.0%. The annual incidence was equal to 2.7/100 000. There was no significant difference between males and females. The Australian National Survey reported the year prevalence of euphoric BD (combined BD-I and BD-II) was 0.5% (Mitchell et al., 2004). The Butajira study from Ethiopia reported a lifetime prevalence of BD-I disorder of 0.5%, with the rate being 0.6% for males and 0.3% for females. The mean age of cases was 29.5 years, with no significant sex difference. The mean age of first recognition of illness was 22 years. There was no significant sex difference in the age at onset of manic or depressive phases (Negash et al., 2005). A more recent cross-sectional, face-to-face, household survey in 11 countries in the Americas, Europe, and Asia reported that the lifetime prevalence was 0.6% for BD-I and 0.4% for BD-II, while the year prevalence was 0.4% and 0.3%, respectively (Merikangas et al., 2011). A few studies report on the epidemiology of bipolar spectrum and suggest that in the adult population the lifetime prevalence of the bipolar spectrum is between 3 and 8.3% (Weissman and Myers, 1978; Angst et al., 1984, 2005b; Oliver and Simmons, 1985; Wicki and Angst, 1991; Heun and Maier, 1993; Angst, 1998; Szadoczky et al., 1998; Hirschfeld et al., 2003a, 2003b; Judd and Akiskal, 2003; Moreno and Andrade, 2005; Faravelli et al., 2006; Kessler et al., 2006).

Overall and according to the WHO, BD affected an estimated 29.5 million persons worldwide in 2004 (WHO, 2008). The available data suggest that the life prevalence of BD-I is around 1%, with probably a similar rate concerning BD-II. The full bipolar spectrum probably has lifetime prevalence around 5%. There are no striking differences between genders. However these figures should be considered as only indicative, since important discrepancies exist among studies and countries, as mentioned above. The rather small difference between annual and lifetime rates suggests that BD is both an episodic but also a chronic mental disorder with high recurrence rates.

The various studies from around the world suggest that the age at onset is late adolescence or early adulthood, around the age of 18 to 20 years, but also they suggest that approximately one-fourth of BD patients have the onset before the age of 13 (Perlis et al., 2004; Post et al., 2008; Stringaris et al., 2010; Merikangas et al., 2012), and among other things this suggests caution in the use of stimulants for the treatment of children with ADHD and worse overall outcome (Agnew-Blais and Danese, 2016).

Staging

After the introduction of operationalized diagnostic criteria for all contemporary classification systems, the need to define and rate seriousness, progression, changes in physiology, and damage made and the extent and the specific characteristics of the disease emerged. Staging is the term that defines this procedure (Fountoulakis, 2015k). The field in medicine where staging is most successful and enjoys great importance is that of clinical oncology. Since 1993 there were many attempts to arrive at a staging model for psychiatry (Fava and Kellner, 1993; Yung and McGorry, 1996, 2007; McGorry et al., 2006, 2007, 2010; McGorry, 2007, 2010b; Vieta et al., 2011; Cosci and Fava, 2013). The concept of staging if and when applied has a number of implications. Almost by definition it suggests that early stages are easier to treat, while later stages are rather refractory to treatment. Thus these later stages might need the application of treatment options with more adverse events, higher risk, and less overall benefit (Post et al., 2010) or some kind of palliative care should be considered.

The earliest research contribution to the effort of staging BD was the description of the stages of mania in the early 1970s when Carlson and Goodwin not only described discrete stages in the development and course of acute mania, but also they described a rollback phenomenon that is the clinical condition improves by manifesting the same stages but at a reverse order (Carlson and Goodwin, 1973). Up to n, 5 major staging models have been proposed for BD (Berk et al., 2007a, 2007b; Kapczinski et al., 2009; Post, 2010; Post et al., 2012; Cosci and Fava, 2013; Frank et al., 2014). Although there is some support for the proposed staging models, the research base is thin, the heterogeneity of the data is significant, and the studies include small sample sizes. A number of vicious logical cycles could be in place. Most of the data are cross-sectional (Kapczinski et al., 2014), and the need for a transdiagnostic and longitudinal research approach is prominent (Lin et al., 2013).

The data so far support the presence of an asymptomatic at-risk phase and a nonspecific prodromal phase. This prodromal phase seems to be common for a number of mental disorders, and prediction is extremely difficult on the basis of current knowledge. The literature is also supportive of the presence of an early stage of the full-blown illness, during which the episodes are well defined and there are no or very few inter-episode residual symptoms, good response to treatment, and little disability. It also supports the presence of a late stage that is associated with a more chronic and refractory disease, probably with depressive predominant polarity, psychotic features, and significant disability. It is disappointing that there is little research on the treatment effect at late stages (Berk et al., 2012), with only a few exceptions (Torrent et al., 2013). The use of biomarkers might, in the near future, facilitate the validation of staging systems and their therapeutic utility (Vieta, 2015).

Therapeutic Issues

The treatment of BD is complex (Fountoulakis, 2008) and for several decades the treatment of BD was theoretically based on

the concept of mood stabilizers. This term was originally used during the 1950s to refer to a combination of amphetamine and a barbiturate to treat patients with neurotic instability but not patients with BD. The term mood normalizer was proposed by Mogens Schou for lithium (Schou, 1963), but eventually the stabilizer concept prevailed, probably because the focus of research with lithium was on the long-term prophylaxis.

During the last decade, however, there was a plethora of data, mainly because of the introduction of atypical antipsychotics as possible treatment options. However, this gave the chance also for older substances to be tested under rigorously defined research conditions. These studies revealed that the treatment could be more complex than previously believed and several issues exist. The clinician should be aware of many specific indications, contraindications, details, and traps (Fountoulakis et al., 2005, 2007a, 2007b, 2008, 2015m; Gonda et al., 2009).

The concept of mood stabilizers is disputed, since the data do not support an equally wide efficacy for different compounds like lithium, valproate, or carbamazepine to warrant such a label. On the contrary, there are negative data concerning specific areas, while our knowledge is quite limited concerning other areas. One particular problem, which only recently has been acknowledged, is that probably some facets of the disorder are refractory to treatment. Another important problem is that not only is the evidence limited concerning the treatment of specific facets and issues of BD (Fountoulakis, 2010; Fountoulakis et al., 2012, 2013), but also continued scientific training and reading is inadequate. Thus, research findings are not making it to everyday clinical practice. Focused educational intervention might be necessary to change this attitude. Part of this problem is reflected in the common practice among clinicians to use medication on the basis of a class effect. This means that they consider that a whole class of medications possesses a specific action. This class effect is often considered in combination with a syndromal approach, which means that irrespective of the nosological entity, a specific kind of symptoms respond to a specific class of medication.

For example, according to this combined approach all antipsychotics are equally effective against psychotic symptoms irrespective of disorder diagnosis, and the same holds for all antidepressants against depressive symptoms. This is the most commonly used approach in everyday clinical practice and has a huge impact on public mental health. Its significant advantage is that it provides the clinician with fast and simple rules to determine treatment. On the other hand, its greatest problem is that this approach has been proven false, especially in the case of BD where it is specifically combined with a very broad mood stabilizers concept (Fountoulakis et al., 2011). The extent to which this truly influences the everyday clinical practice worldwide is unknown but is probably significant. The extent to which this concept influences the outcome of BD is similarly unknown, although theoretically a more evidence-based approach should improve the overall outcome of BD patients.

It is important to note that with the introduction of the second-generation antipsychotics, antipsychotics became a cornerstone for the treatment of BP also according to treatment guidelines. On the contrary, a number of studies showed that the usefulness of antidepressants that were traditionally seen in Europe as a meaningful treatment option for bipolar depression is questionable (Pacchiarotti et al., 2013b). Additionally, the maintenance/long-term treatment became more complex, since it has proven that agents previously considered to be mood stabilizers were essentially more effective for one pole than the other (Popovic et al., 2012). This is definitely a fast-moving field,

and it is certainly difficult for a clinician to follow new findings and incorporate them into his or her everyday clinical practice.

On the other hand, the data on the usefulness of psychosocial interventions are limited, and their value against specific symptoms and problems remains unknown (Fountoulakis et al., 2009; Reinares et al., 2014).

One very special issue is agitation and its treatment. There is a significant number of published papers on the pharmacological (Citrome, 2004; Battaglia, 2005; Nordstrom and Allen, 2007; Nordstrom et al., 2012) but also on the nonpharmacological treatment of agitation (Marder, 2006; Amann et al., 2013), while recently a consensus paper on how to treat agitation in BD patients has been published (Garriga et al., 2016).

Special Issues

Since BD is characterized by phases that respond to a completely different way to treatment, it is of outmost importance to define phases of treatment and comorbidity (Fountoulakis, 2015f; Vrublevska and Fountoulakis, 2015).

It is relatively easy to define acute either manic/hypomanic or depressive episodes. However the terms continuation and maintenance are often interchangeably used in the terminology of RCTs and thus create significant confusion (Frank et al., 1991; Ghaemi et al., 2004). Continuation treatment lasts up to 12 months, and the duration depends on an estimate of when the episode would have remitted spontaneously. On the other hand, maintenance treatment starts after remission and thus after continuation and covers several years. Although a strict definition demands at least 2 months of sustained recovery for the patient to be considered in remission (American Psychiatric Association, 2000), the reality is that only a minority of patients in RCTs achieve complete remission. This makes the use of terms (relapse vs recurrence and continuation vs maintenance) problematic. In the nomenclature of RCTs, the terms relapse and maintenance are preferred. The FDA policy is to accept data, based on patients in remission for <2 months, thus adding to the continuation vs maintenance confusion of definitions (Calabrese et al., 2006).

The term relapse is also problematic in BD. A narrow definition suggests that relapses are of the same polarity with the index episode, and they tend to occur within the first months of improvement. However with a polymorphic disease like BD, it might be inappropriate not to include in relapses the early emergence of an episode of the opposite pole. It is important to note that licensing authorities accept the latter approach.

The acute episode after which BD patients are enrolled into maintenance trials is called the index episode. To date, most maintenance trials follow an enriched design, that is, only patients who have remitted under the investigation agent during the acute phase are enrolled into the double blind maintenance phase. This design has interesting consequences, since it biases the sample both towards a specific predominant polarity and also towards a favorable response to the specific agent (Cipriani et al., 2014). These 2 comments constitute important limitations in the generalizability of the results and make very difficult the translation of research findings into the everyday clinical practice in the case of patients who, rather than continued on the same medication, are switched to another one during the maintenance phase (Grande et al., 2014).

Economic Considerations

It is very difficult to calculate the true economic cost of a polymorphic disorder like BD. The cost includes direct spending due

to hospitalizations and medication, cost of supporting infrastructure of the various National Health Systems, somatic comorbidity, indirect and out-of-pocket costs, as well as the absenteeism from work and premature death (Fountoulakis, 2015g).

For the UK the total cost has been estimated to be £2.055 billion in 1999/2000 prices (Das Gupta and Guest, 2002). It is interesting that 86% of this cost was the result of productivity loss and unemployment, while only 10% was cost related with NHS services. Medication costs in primary care were approximately £8.5 million, corresponding to 0.4% of total cost and 4.3% of NHS cost. A more recent study showed that the NHS cost has been doubled, with medication costs rising disproportionately and reaching £25.2 million, that is 7.4% of NHS cost (Young et al., 2011). In the US, the cost of medication was rather very low during the 1990s and reached 2% of the total cost after 2000, but the exact figure is unknown (Wyatt and Henter, 1995; Begley et al., 2001; McCrone et al., 2008; Dilsaver, 2011). In Germany the total annual cost was calculated to be 5.8 million euros, with 98% being due to productivity loss (Runge and Grunze, 2004). Similar estimations come from other areas of the world, with the calculations basing on different prevalence rates and health systems and societal structures (Hakkaart-van Roijen et al., 2004; Fisher et al., 2007; Ekman et al., 2013).

It is clear that the cost of medication treatment constitutes a very small percentage of the total cost of BD (Hidalgo-Mazzei et al., 2015). Medication treatment is, however, the intervention with the greatest impact on the course of the illness and the intervention that makes possible other actions to exist by resolving acute episodes in a reliable way. Furthermore, it decreases the long-term impairment and improves insight and collaboration by the side of the patient. Nevertheless, it also seems clear that medication cost is disproportionately rising, at least in some places of the world and for periods of time, and this constitutes an additional factor of concern. One should be very careful, because a small reduction in medication costs as a consequence of giving priority to cheaper agents and disregarding clinical data could easily result in a significant and disproportional increase in the total cost of the disease.

The CINP workgroup decided not to take medication cost or availability of medication into consideration. It chose to rely exclusively on clinical data, leaving the cost and availability issues to local and national groups who would like to implement the CINP guidelines in a specific country or region and would be obliged to take into consideration also the local socio-political and economic environment.

Methodology

The workgroup decided after consensus to follow the following methodology for the development of the treatment algorithm with the steps listed below:

- a) Defining the sources of data and choosing which to use
- b) Development of a grading method
- c) Search of the literature
- d) Grading of the data
- e) Defining the clinical parameters to take into consideration
- f) Development of a precise treatment algorithm
- g) Development of the clinical guideline

Defining the Sources of Data and Choosing Whom to Use

Randomized Controlled Trials (RCTs)

This type of study constitutes the main source of evidence. Without them it is impossible to say whether an agent or

method possesses efficacy or not, since it is impossible to control for confounding variables any other way. Randomization of patients to parallel treatment arms, including placebo, allows one to attribute confidently observed differences in efficacy between these arms to the effects of the treatments (McAlister et al., 1999a, 1999b; Pocock and Elbourne, 2000).

However, with BD there is an important problem. Because of ethical, practical, and most often economic limitations, it is not always possible to apply the RCT method across all facets and issues of BD. For many of them, data are available only on the basis of posthoc analyses or secondary outcomes. Another major limitation is that this kind of study is very expensive, and thus most of them are industry sponsored with the objective to obtaining the label for the specific product. Although such trials follow the regulatory agencies' design, they have limitations on generalizability. Also it is well known that only a small minority of highly selected patients is eligible to enter these studies, and thus the generalizability of results is problematic. The study duration is often relatively short and this is true also for maintenance trials, in part because the existence of a placebo arm carries a high attrition rate.

An important pitfall concerns the actual results of the RCT, which often are different from those published. It is not unusual that when a trial is negative on the basis of its primary outcome, a publication is done on the basis of positive secondary outcomes. This is essentially misleading, but fortunately it is a phenomenon that has been less frequent during the recent years.

Meta-Analysis

Meta-analysis is a technique that combines data (not simply pooled) from several trials and returns a specific quantitative answer to a specific question that usually is which treatment is superior in comparison with others or placebo. Sometimes but not always it also provides an absolute estimate of the treatment effect size.

There are a number of significant limitations for the meta-analytic methods (Huf et al., 2011a, 2011b). There is a need for the studies included in the meta-analysis to be similar in design and with sufficient information being available. Meta-analytic studies often violate this rule and include a diverse group of trials in the analysis (e.g., studies of monotherapy and combination treatment, fixed and flexible dosage studies, etc.) with unknown consequences (Fountoulakis et al., 2014).

Common problems of meta-analyses include small sample sizes, inadequate power, study heterogeneity, lack of extractable data, lack of interchangeable measurement instruments and definitions of outcomes, and other differences in the design of studies whose data are utilized. Negative trials are often not published and this poses an important limitation to the meta-analytic approach. Today the trials sponsored by official foundations can be traced in trial repositories. However, their detailed results are unlikely to be retrieved and even if they are retrieved, they have not undergone the essential peer review process (which adds credibility) like those published, and their quality could be questionable.

The question whether it is appropriate to use data from the largest possible number of disparate studies vs the need for including data only from essentially identical studies is a matter of debate and has also been discussed specifically concerning acute mania trials where these different approaches gave conflicting results (Yildiz et al., 2010; Cipriani et al., 2011). Practically, all meta-analytical studies utilize compromises to deal with the above problems and limitations. These compromises might

have profound effects on the validity and generalizability of their results (Noble, 2006; Mismetti et al., 2007; Huf et al., 2011b).

Some authors consider meta-analysis to be on the top of the evidence-based pyramid of data sources. This approach suggests that its results are superior to the results of the RCTs, and subsequently it is meant that a positive meta-analysis is superior to a number of negative RCTs even in the case of the absence of any positive RCTs. However, the authors of the current paper consider that in most cases meta-analysis has a lower evidence level than RCTs and therefore graded it below them, primarily because of a significant number of limitations and drawbacks that often make the results of meta-analysis equivocal.

Open Trials

Open trials do not utilize the double blind design and they are not placebo controlled. Therefore they are easier to conduct, their number and size are greater, and the quality of patients enrolled is closer to that seen in the real world. Their great limitation is that their open nature induces significant bias, and thus they are by no means considered to be even close to being the gold standard or a reliable source of evidence data. Their role should be considered complementary. It is not unusual that treatment modalities with many positive open trials fail in RCTs, with topiramate in BD being a striking example (Suppes, 2002).

Review and Opinion Papers

Review and opinion papers mainly constitute educational tools, which attempt to translate the research findings into ready-to-use tools for the everyday clinical practice. They are extremely useful for the average clinician; however, they usually echo the opinion of the author, and thus they might contain significant bias. Their overall reliability and validity is questionable and only a few add significantly to our understanding by critically analyzing the existing data. Their ever-increasing number in the literature might constitute a problem, since they often obscure research findings by reproducing widely established biases and misconceptions. This is an important problem especially in the field of BD treatment.

Sources to Include

The authors decided by consensus to include only RCTs and meta-analyses in the development of the current treatment algorithm, since they have the highest validity for judgment. The authors reserved the privilege to judge and use the second and third source on an individual basis and according to their research and clinical experience for the latter steps of the algorithm where a Delphi method to arrive at decisions was utilized.

Development of a Grading Method

The authors decided to develop a grading method for the evaluation of available data concerning the treatment of BD. Such methods have existed since the early 1980s (Fletcher and Spitzer, 1980), but the de novo development of such a method was judged to be absolutely necessary, because the existing grading methods were not sufficiently appropriate for use in this particular set of data. In the frame of this process, the most widely accepted grading methods were studied, and their advantages and disadvantages were identified and taken into consideration in relationship to the specific needs of the current study. All grading methods include a method to assess the quality of data and a method to arrive at recommendations on the basis of the extent to which we can be confident that the desirable effects of an intervention outweigh the undesirable effects. The values

and preferences factor as well, but the cost was not taken into consideration by the workgroup.

Starting in 1992, 5 steps were developed to summarize the process of individual-level decision making and they were published in 2005 (Dawes et al., 2005). They include:

- The formulation of a precise and answerable question and avoiding uncertainty and vague statements (Richardson et al., 1995; Schlosser et al., 2007).
- The performance of a systematic search and retrieval of the evidence available (Rosenberg et al., 1998).
- The critical review and classification of the retrieved evidence with the recognition of the presence of systematic errors, various types of bias, confounders, reliability and validity issues, etc. The clinical significance and the generalizability of the results should also be taken into account (Parkes et al., 2001; Horsley et al., 2011).
- Application of results in practice.
- Evaluation of performance (Jamtvedt et al., 2003, 2006a, 2006b; Ivers et al., 2012).

It is important to assess the quality of the evidence that comes from the sources described above. The quality assessment is based on the strength of their freedom from the various biases that beset medical research. In this frame, triple-blind, placebo-controlled trials with allocation concealment and complete follow-up involving a homogeneous patient population and medical condition should be considered to constitute the highest grade, while case reports should be considered to constitute the lowest grade. Expert opinion should not be considered to be a source of evidence, although it could be a valuable tool for the development of guidelines (Tonelli, 1999).

Until recently there were a number of grading systems for assessing the quality of evidence that were developed by different organizations. One of them is the U.S. Preventive Services Task Force (U.S. Preventive Services Task Force, 1989; Sherman et al., 2011) and another system is the Oxford (UK) Center for Evidence Based Medicine Levels of Evidence, which also is useful for the grading of diagnostic tests, prognostic markers, or harm (Oxford (UK) Center for Evidence Based Medicine Levels of Evidence Working Group) and constituted the basis for the use of the BCLC staging system for diagnosing and monitoring hepatocellular carcinoma

in Canada (Paul et al., 2012). Another method to grade data is the Patient Outcomes Research Team (PORT) method (Lehman and Steinwachs, 1998), which has been used by the World Federation of Societies of Biological Psychiatry for the development of the WFSBP guidelines (Grunze et al., 2002, 2003, 2004). In 1992 the Agency for Health Care Policy and Research and the National Institute of Mental Health established a PORT for Schizophrenia at the University of Maryland School of Medicine and the Johns Hopkins University School of Public Health. The PORT investigators adopted the criteria on levels of evidence used for development of the Agency for Health Care Policy and Research Depression Guidelines.

The most detailed and precise modern method seems to be the GRADE method (short for Grading of Recommendations Assessment, Development and Evaluation) for the development of guidelines (Guyatt et al., 2008b; Jaeschke et al., 2008), which clearly separates quality of evidence from level of recommendation and suggests it is necessary to include a clear question that should include all 4 components of clinical management (patients, an intervention, a comparison, and the outcomes of interest) (Oxman and Guyatt, 1988) and to grade the outcomes into those who are critical for the decision making and those who are not (Schunemann et al., 2006). In this frame, the assessment of the quality of evidence is important, since it reflects the confidence whether the effect is adequate to support recommendations. The determinants of quality are study limitations, inconsistency of results, indirectness of evidence, imprecision, and reporting bias (Guyatt et al., 2011a, 2011b, 2011c, 2011d, 2011e, 2013). There is some option to upgrade the quality when the effect size is very high (Guyatt et al., 2011f). The GRADE method provides guidance to grade the data from a variety of sources (Guyatt et al., 2008a), but it is not sensitive for datasets that focus solely on RCTs like the dataset of the current workgroup. According to the GRADE grading system, all the data included in the current effort to develop guidelines are of high quality. From the limitations recognized by the GRADE (lack of allocation concealment, lack of blinding, large losses to follow-up, failure to adhere to an intention to treat analysis, and stopping early for benefit or failure to report outcomes), only large losses to follow-up and stopping early for benefit or failure to report outcomes could be applicable to the current study. A comparison of all the grading methods is shown in Table 3.

Table 3. Comparative Presentation of Different Grading Methods

USPSTF	OCEBM	GRADE	PORT
Level I: Evidence obtained from at least one properly designed randomized controlled trial.	Systematic review of randomized trials or n-of-1 trials	High quality	Level A: Good research-based evidence, with some expert opinion, to support the recommendation
Level II-1: Evidence obtained from well-designed controlled trials without randomization.	Randomized trial or observational study with dramatic effect	Medium quality	Level B: Fair research-based evidence, with substantial expert opinion, to support the recommendation
Level II-2: Evidence obtained from well-designed cohort or case-control analytic studies, preferably from more than one center or research group.	Nonrandomized controlled cohort/follow-up study	Low quality	
Level II-3: Evidence obtained from multiple time series designs with or without the intervention. Dramatic results in uncontrolled trials might also be regarded as this type of evidence.	Case-series, case-control studies, or historically controlled studies	Very low quality	Level C: Recommendation based primarily on expert opinion, with minimal research-based evidence, but significant clinical experience
Level III: Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.	Mechanism-based reasoning		

The recommendation methods constitute a step forward and are determined by the balance of risk vs benefit of the intervention and the level of evidence on which this information is based. A comparison of the recommendation methods of the U.S. Preventive Services Task Force uses (Sherman et al., 2011) that utilizes a 5-levels system and the GRADE system that has only 2 categories concerning recommendations and characterizes them as strong (conditional) and weak (discretionary) (Guyatt et al., 2008b, 2008c) and also considers cost (Brunetti et al., 2013) is shown in Table 4.

As defined previously, only RCTs were taken into consideration, a fact that puts all the data at the highest grading according to all

systems. However, the workgroup was concerned about a number of issues, including inconsistency of results between RCTs, conflicting results between RCTs and meta-analyses, issues explored only on the basis of secondary outcomes, etc. After recognizing all these sources of problematic quality, 32 individual scenarios were identified and are listed in Table 5. Afterwards they were ranked after consensus and grouped into levels. Two solutions were proposed. The ranking, the 4- and 5-levels solution, and the final grading system are shown in Table 6. The description of the grading and the recommendation systems are shown in Table 7.

At this point it is important to note that the absence of evidence is not identical with the presence of negative data.

Table 4. Comparative Presentation of Recommendation Methods

USPSTF	GRADE
Level A: Good scientific evidence suggests that the benefits of the clinical service substantially outweigh the potential risks.	Strong
Level B: At least fair scientific evidence suggests that the benefits of the clinical service outweighs the potential risks.	
Level C: At least fair scientific evidence suggests that there are benefits provided by the clinical service, but the balance between benefits and risks are too close for making general recommendations.	Weak
Level D: At least fair scientific evidence suggests that the risks of the clinical service outweighs potential benefits.	
Level I: Scientific evidence is lacking, of poor quality, or conflicting, such that the risk versus benefit balance cannot be assessed.	

Abbreviations: GRADE, Grading of Recommendations Assessment, Development and Evaluation for the Development of Guidelines; OCEBM, Oxford (UK) Center for Evidence Based Medicine; PORT, Patient Outcomes Research Team; USPSTF, U.S. Preventive Services Task Force.

Table 5. The 32 Different Scenarios That Were Identified, Listed, and Graded

Primary Outcome Scenarios
1. At least 1 positive 2-active arm RCTs vs placebo exist, plus positive 1 active arm RCTs. No negative RCTs
2. At least 2 positive RCTs vs placebo exist. No negative RCTs
3. One positive RCT vs placebo exists. No negative RCTs
4. Some positive plus some negative RCTs vs placebo. Positive all meta-analyses
5. Some positive plus some negative RCTs vs placebo. Mixed results from meta-analyses
6. Some positive plus some negative RCTs vs placebo. Negative all meta-analyses
7. More positive but some negative RCTs vs placebo. Positive all meta-analyses
8. More positive but some negative RCTs vs placebo. Mixed results from meta-analyses
9. More positive but some negative RCTs vs placebo. Negative all meta-analyses
10. More negative but some positive RCTs vs placebo. Positive all meta-analyses
11. More negative but some positive RCTs vs placebo. Mixed results from meta-analyses
12. More negative but some positive RCTs vs placebo. Negative all meta-analyses
13. Only 1 negative trial exists vs placebo
14. Only negative trials exist vs placebo. Meta analyses all negative
15. Only negative trials exist vs placebo. Meta analyses all positive
16. Only negative trials exist vs placebo. Meta analyses mixed
Posthoc scenarios
17. Only 1 positive from posthoc analyses vs placebo
18. At least 2 positive from posthoc analyses vs placebo
19. Only 1 negative from posthoc analyses vs placebo
20. At least 2 negative from posthoc analyses vs placebo. Positive all meta-analyses
21. At least 2 negative from posthoc analyses vs placebo. Negative all meta-analyses
22. At least 2 negative from posthoc analyses vs placebo. mixed meta-analyses
23. More negative than positive from posthoc analyses vs placebo. Positive all meta-analyses
24. More negative than positive from posthoc analyses vs placebo. Negative all meta-analyses
25. More negative than positive from posthoc analyses vs placebo. mixed meta-analyses
26. More positive than negative from posthoc analyses vs placebo. Positive all meta-analyses
27. More positive than negative from posthoc analyses vs placebo. Negative all meta-analyses
28. More positive than negative from posthoc analyses vs placebo. mixed meta-analyses
Other scenarios
29. Only 1 failed trial, no other data
30. At least 2 failed trials, no other data
31. Only prematurely terminated trials
32. Although trials exist, the data are not available in a way to arrive at reliable conclusions

Abbreviations: GRADE, Grading of Recommendations Assessment, Development and Evaluation) for the Development of Guidelines; USPSTF, U.S. Preventive Services Task Force.

Table 6. The Ranking, the 4- and 5-levels Solution, and the Final Grading System for the 32 Different Scenarios

Scenario	Rank	Solutions		
		5-Grade	4-Grade	Grade system
At least 1 positive 2-active arm RCTs vs placebo exist, plus positive 1 active arm RCTs. No negative RCTs	1	A	A	1
At least 2 positive RCTs vs, placebo exist. No negative RCTs	1	A	A	1
One positive RCT vs placebo exists. No negative RCTs	2	A	B	2
More positive but some negative RCTs vs placebo. Positive all meta-analyses	2	A	B	2
Some positive plus some negative RCTs vs placebo. Positive all meta-analyses	3	B	B	2
More negative but some positive RCTs vs placebo. Positive all meta-analyses	4	B	B	2
Only negative trials exist vs placebo. Meta analyses all positive	4	B	B	2
At least 2 positive from posthoc analyses vs placebo	5	B	C	3
Only 1 positive from posthoc analyses vs placebo.	5	B	C	3
Some positive plus some negative RCTs vs placebo. Mixed results from meta-analyses	6	C	C	3
More positive but some negative RCTs vs placebo. Mixed results from meta-analyses	6	C	C	3
More positive than negative from posthoc analyses vs placebo. Positive all meta-analyses	7	D	C	3
More negative than positive from posthoc analyses vs placebo. Positive all meta-analyses	7	D	C	3
At least 2 negative from posthoc analyses vs placebo. Positive all meta-analyses	7	D	C	3
More positive than negative from posthoc analyses vs placebo. mixed meta-analyses	8	E	C	3
More negative but some positive RCTs vs placebo. Mixed results from meta-analyses	9	E	D	4
Only negative trials exist vs placebo. Meta analyses mixed	9	E	D	4
At least 2 negative from posthoc analyses vs placebo. Mixed meta-analyses	10	E	D	4
More negative than positive from posthoc analyses vs placebo. Mixed meta-analyses	10	E	D	4
Some positive plus some negative RCTs vs placebo. Negative all meta-analyses	neg	neg	neg	5
More positive but some negative RCTs vs placebo. Negative all meta-analyses	neg	neg	neg	5
More negative but some positive RCTs vs placebo. Negative all meta-analyses	neg	neg	neg	5
Only 1 negative trial exists vs placebo	neg	neg	neg	5
Only negative trials exist vs placebo. Meta analyses all negative	neg	neg	neg	5
Only 1 negative from posthoc analyses vs placebo	neg	neg	neg	5
At least 2 negative from posthoc analyses vs placebo. Negative all meta-analyses	neg	neg	neg	5
More negative than positive from posthoc analyses vs placebo. Negative all meta-analyses	neg	neg	neg	5
More positive than negative from posthoc analyses vs placebo. Negative all meta-analyses	neg	neg	neg	5
Only prematurely terminated trials	neg	neg	neg	5
Although trials exist, the data are not available in a way to arrive at reliable conclusions	neg	neg	neg	5
Only 1 failed trial, no other data	unknown	unknown	unknown	
At least 2 failed trials, no other data	unknown	unknown	unknown	

All treatment agents were graded also in terms of safety and tolerability. All combination options were graded at best with 2, since they put the patient at a higher risk for manifesting adverse events.

Search of the Literature

The workgroup decided that the PRISMA method (Hopewell et al., 2008; Liberati et al., 2009; Moher et al., 2009a, 2009b) should be followed in the search of the literature, which will include 3 kinds of papers:

- i. RCTs (placebo controlled as well as clinical trials with an active comparator with the compounds used as monotherapy or add-on therapy).
- ii. Posthoc analyses of RCTs
- iii. Meta-analyses and review papers
- iv. Treatment guidelines papers

The search strategies will include:

1. To locate RCTs, the combination of the words 'bipolar,' 'manic,' 'mania,' 'manic depression,' and 'manic depressive' and 'randomized' will be used.
2. Webpages containing lists of clinical trials will be scanned. These sites include <http://clinicaltrials.gov> and <http://www.clinicalstudyresults.org> as well as the official sites of all the pharmaceutical companies with products used for the treatment of BD.

3. Relevant review articles will be scanned and their reference lists will be utilized.
4. The MEDLINE will be searched with the combination of keywords 'guidelines' or 'algorithms' with 'mania,' 'manic,' 'bipolar,' 'manic-depressive,' or 'manic depression.'
5. The treatment guidelines will also be scanned and their reference lists will be utilized.
6. Only papers in English language will be included.

Additionally, an unstructured search of the literature will be performed concerning the adverse events and other safety issues of treatment options

The workgroup considered the fact that it is difficult to locate unpublished studies, especially old ones, and even more difficult to retrieve their results. Thus it was decided that the focus should be put mainly on published studies which are definitely peer reviewed, are of higher quality, and provide more details than meeting abstracts or report sheets. However, whenever an unpublished trial should be located, it is mentioned in the specific part of the manuscript. The authors decided not to seek for additional information concerning unpublished trials from manufacturers, because this might increase the retrieval bias.

Grading of the Data

The grading of the data will follow their retrieval and will be done according to the method developed and described in the

current paper. The grading will be included in the second paper concerning the CINP guidelines for BD

Defining the Clinical Parameters to Take into Consideration

In the real-world setting, the therapist encounters patients with specific clinical features that often determine the choice of treatment on the basis of clinical experience and wisdom rather than evidence. These features include the so-called core manic and core depressive features, psychotic features, anxiety, the co-occurrence of manic and depressive symptoms in a variety of combinations that often do not correspond to concepts accepted by modern classification systems, agitation, and rapid cycling. It is interesting to address the complete constellation of symptoms instead of a specific group. The problem is that the data often focus on the second rather than the first option. It is important also to consider the predominant polarity and subtype of BD (BD-I vs BD-II), the personal history of the patient, and more specifically previous response or refractoriness to treatment and adverse events (including switch).

The data will be scanned concerning the treatment of all the above conditions and modifiers and relevant conclusions will be made concerning whether they can be used as clinical cues for the selection of appropriate treatment.

Development of a Precise Algorithm

The development of a precise algorithm for experimental reasons will be the first task. This algorithm will be based exclusively on the evidence and will be the next step after the data and the interventions are graded in terms of recommendation. This algorithm will be based on the data in a narrow and strict sense and

might provide with very precise but limited treatment options for the everyday clinical practice. There will be no trade between the evidence-based approach and clinical utility; the first will be absolutely dominant. This algorithm will reflect the exact state of the art concerning hard data but will lack any clinical wisdom, and it is expected that its application in everyday clinical practice will be problematic. Therefore it should be considered as experimental, and clinicians who will wish to apply it in their clinical practice should do so by taking into consideration these advantages and disadvantages. The algorithm will be included in the second paper concerning the CINP guidelines for BD, and it will be accompanied by a detailed table with the grading recommendation of all available interventions during all the phases of BD and in relevance with the presence of specific clinical features.

At a later time point a software application will be developed by the CINP to assist with the use of the algorithm.

Development of the Clinical Guideline

The development of the guideline will follow after the data and the interventions have been graded and the precise algorithm has been developed. The guideline will be included in the third paper concerning the CINP guidelines for BD. The workgroup decided after consensus on the following rules for the development of the guidelines:

- i. Overall the guideline should be based on existing research hard evidence, but also it should make sense for the everyday clinical practice and should be user friendly. Although their nature will be based on the evidence-based approach, this should not go too far concerning the interpretation of the research findings and the potential clinical implications.

Table 7. Summary of the Method for the Grading of the Data and Recommendation as Decided by the Workgroup on the Basis of Both Efficacy and Safety Tolerability

Grading on Basis of Efficacy

Level 1	Good research-based evidence, supported by at least 2 placebo controlled studies of sufficient magnitude and good quality. In case of the presence of negative RCTs, positive RCTs should outnumber negative ones
Level 2	Fair research-based evidence, from one randomised, double-blind placebo controlled trial. Also in case one or more trials exist, however, they fail to fulfil all the criteria above (e.g., very small sample size or no placebo control) as well as in case of positive meta-analysis alone.
Level 3	Some evidence from comparative studies without placebo arm or from posthoc analyses.
Level 4	Inconclusive data or poor quality of RCTs
Level 5	Negative data

Grading on the basis of safety and tolerability

Level 1	Very good tolerability, few side effects which are not enduring, they do not cause significant distress and are not life-threatening and they do not compromise the overall somatic health of the patient
Level 2	Moderate tolerability, many side effects which could be enduring, and cause significant distress but they are not life-threatening although they could compromise the overall somatic health of the patient. Agents with very good overall tolerability but with rare life-threatening adverse events, could be classified here only if the lethality risk can be essentially considered to be negligible with the application of procedures and protocols (e.g., laboratory testing, titration schedules, etc.)
Level 3	Poor tolerability, many side effects which are enduring, cause significant distress, compromise the overall somatic health of the patient or are life-threatening. Agents with moderate overall tolerability and rare life-threatening adverse events should be classified here even in cases the lethality risk can be essentially considered to be negligible with the application of procedures and protocols (e.g., laboratory testing, titration schedules, etc.)

Recommendations for treatment (combination of efficacy and safety/tolerability)

Level 1	Level 1 or 2 for efficacy and 1 for safety/tolerability
Level 2	Level 1 or 2 for efficacy and 2 for safety/tolerability
Level 3	Level 3 for efficacy and 1 or 2 for safety/tolerability
Level 4	Level 4 for efficacy or 3 for safety/tolerability
Level 5	Level 5 for efficacy (not recommended)

- ii. Agents and treatment modalities with proven efficacy across all 3 phases of the illness (acute mania, acute bipolar depression, and maintenance phase concerning the prevention of both manic and depressive episodes) should be given priority.
- iii. No economic and availability issues will be taken into consideration. National bodies that might wish to utilize the CINP guidelines could add such analyses tailored to the specific country or region.

Discussion

The current paper sets the frame for the development of the CINP treatment guidelines for BD. It contains all the background information, including important clinical features, staging methods, and important treatment issues and details. It also elaborates on the methodology to be used and describes the development of a grading system that will be suitable for use with the kind of data under consideration.

The overall aim of the workgroup was to push guidelines one step further by evaluating the available data in depth and also by identifying clinical issues that need specific interventions that could be supported by the data. A significant contribution is expected to be the precise experimental algorithm that will constitute an option for further study.

Acknowledgment

The authors thank Professor Guy Goodwin for his valuable input in the authoring of this manuscript.

Statement of Interest

K.N.F. has received grants and served as consultant, advisor, or CME speaker for the following entities: AstraZeneca, Bristol-Myers Squibb, Eli Lilly, Ferrer, Gedeon Richter, Janssen, Lundbeck, Otsuka, Pfizer, the Pfizer Foundation, Sanofi-Aventis, Servier, Shire, and others.

E.V. has received grants and served as consultant, advisor, or CME speaker for the following entities: Allergan, AstraZeneca, Bristol-Myers Squibb, Dainippon Sumitomo Pharma, Ferrer, Forest Research Institute, Gedeon Richter, Glaxo-Smith-Kline, Janssen, Lilly, Lundbeck, Otsuka, Pfizer, Roche, Sanofi-Aventis, Servier, Shire, Sunovion, Takeda, the Brain and Behaviour Foundation, the Spanish Ministry of Science and Innovation (CIBERSAM), the Seventh European Framework Programme (ENBREC), and the Stanley Medical Research Institute. A.H.Y. is employed by King's College London; is Honorary Consultant SLam (NHS UK); has paid lectures by and participated in advisory boards for all major pharmaceutical companies with drugs used in affective and related disorders; and has no share holdings in pharmaceutical companies. He was lead Investigator for Embolden Study (AZ), BCI Neuroplasticity study, and Aripiprazole Mania Study; investigator initiated studies from AZ, Eli Lilly, Lundbeck, and Wyeth; and has received grant funding (past and present) from: NIHR-BRC (UK); NIMH (USA); CIHR (Canada); NARSAD (USA); Stanley Medical Research Institute (USA); MRC (UK); Wellcome Trust (UK); Royal College of Physicians (Edinburgh); BMA (UK); UBC-VGH Foundation (Canada); WEDC (Canada); CCS Depression Research Fund (Canada); MSFHR (Canada); and NIHR (UK).

H.G. within the last 3 years received grant/research support from: NIHR UK, MRC UK, NTW, and NHS Foundation Trust; receipt of honoraria or consultation fees from: Gedeon-Richter, Lundbeck, and Hofmann-LaRoche; and participated in

a company-sponsored speaker's bureau at BMS, Ferrer, Janssen-Cilag, Otsuka, Lundbeck, and Pfizer.

L.Y. has been on speaker/advisory boards for, or has received research grants from Alkermes, Allergan, AstraZeneca, Bristol Myers Squibb, CANMAT, CIHR, Eli Lilly, Forest, GlaxoSmithKline, Intas, Janssen, the Michael Smith Foundation for Health Research, Pfizer, Servier, Sumitomo Dainippon, Sunovion, and the Stanley Foundation.

S.K. within the last 3 years received grants/research support, consulting fees, and honoraria from Angelini, AOP Orphan Pharmaceuticals AG, AstraZeneca, Eli Lilly, Janssen, KRKA-Pharma, Lundbeck, Neuraxpharm, Pfizer, Pierre Fabre, Schwabe, and Servier.

H.J.M. received honoraria for lectures or advisory activities or received grants by the following pharmaceutical companies: Lundbeck, Servier, Schwabe, and Bayer. He was president or in the executive board of the following organizations: CINP, ECNP, WFSBP, EPA, and chairman of the WPA-section on Pharmacopsychiatry.

P.B. has received research grants, honoraria for participation in advisory boards, and/or gave presentations from Allergan, AstraZeneca, Bristol Myers Squibb, Canadian Institute for Health Research, Eli Lilly, Lundbeck, Janssen, Ontario Brain Institute, Meda-Valeant, Merck, Otsuka, Pierre Fabre Medicaments, Pfizer, Shire, Sunovion, and Takeda.

References

- Abrams R, Taylor MA (1976) Mania and schizo-affective disorder, main type: a comparison. *Am J Psychiatry* 133:445–447.
- Abrams R, Taylor MA (1981) Importance of schizophrenic symptoms in the diagnosis of mania. *Am J Psychiatry* 138:658–661.
- Agnew-Blais J, Danese A (2016) Childhood maltreatment and unfavourable clinical outcomes in bipolar disorder: a systematic review and meta-analysis. *Lancet Psychiatry* 3:342–349.
- Akiskal H (2000) Mood disorders. In: *Comprehensive Textbook of Psychiatry* (Sadock B, Sadock V, eds), pp1338–1377. Philadelphia: Lippincott Williams & Wilkins.
- Akiskal HS, Pinto O (1999) The evolving bipolar spectrum. Prototypes I, II, III, and IV. *Psychiatr Clin North Am* 22:517–534, vii.
- Akiskal HS, Benazzi F (2005) Atypical depression: a variant of bipolar II or a bridge between unipolar and bipolar II? *J Affect Disord* 84:209–217.
- Akiskal HS, Djenderedjian AM, Rosenthal RH, Khani MK (1977) Cyclothymic disorder: validating criteria for inclusion in the bipolar affective group. *Am J Psychiatry* 134:1227–1233.
- Akiskal HS, Rosenthal RH, Rosenthal TL, Kashgarian M, Khani MK, Puzantian VR (1979) Differentiation of primary affective illness from situational, symptomatic, and secondary depressions. *Arch Gen Psychiatry* 36:635–643.
- Allison JB, Wilson WP (1960) Sexual behavior of manic patients: a preliminary report. *South Med J* 53:870–874.
- Altamura AC, Dell'Osso B, Berlin HA, Buoli M, Bassetti R, Mundo E (2010) Duration of untreated illness and suicide in bipolar disorder: a naturalistic study. *Eur Arch Psychiatry Clin Neurosci* 260:385–391.
- Amann BL, Wesuls R, Landin Romero R, Grunze H (2013) [De-escalation and atypical antipsychotics in the treatment of acute mania]. *Fortschr Neurol Psychiatr* 81 Suppl 1:S9–16.
- American Psychiatric Association (2000) *Diagnostic and Statistical Manual of Mental Disorders 4th Edition, Text Revision, DSM-IV-TR*. Washington, DC: American Psychiatric Publishing.
- American Psychiatric Association (2013) *Diagnostic and Statistical Manual of Mental Disorders 5th ed., DSM-5*. Washington, DC: American Psychiatric Association.

- Angst J (1978) The course of affective disorders. II. Typology of bipolar manic-depressive illness. *Arch Psychiatr Nervenkr* 226:65–73.
- Angst J (1998) The emerging epidemiology of hypomania and bipolar II disorder. *J Affect Disord* 50:143–151.
- Angst J (2007) The bipolar spectrum. *Br J Psychiatry* 190:189–191.
- Angst J, Grof P, Schou M (1969) Lithium. *Lancet* 1:1097.
- Angst J, Dobler-Mikola A, Binder J (1984) The Zurich study--a prospective epidemiological study of depressive, neurotic and psychosomatic syndromes. I. Problem, methodology. *Eur Arch Psychiatry Neurol Sci* 234:13–20.
- Angst J, Felder W, Frey R, Stassen HH (1978) The course of affective disorders. I. Change of diagnosis of monopolar, unipolar, and bipolar illness. *Arch Psychiatr Nervenkr* 226:57–64.
- Angst J, Sellaro R, Stassen HH, Gamma A (2005a) Diagnostic conversion from depression to bipolar disorders: results of a long-term prospective study of hospital admissions. *J Affect Disord* 84:149–157.
- Angst J, Weis P, Grof P, Baastrup PC, Schou M (1970) Lithium prophylaxis in recurrent affective disorders. *Br J Psychiatry* 116:604–614.
- Angst J, Gamma A, Neuenschwander M, Ajdacic-Gross V, Eich D, Rossler W, Merikangas KR (2005b) Prevalence of mental disorders in the Zurich Cohort Study: a twenty year prospective study. *Epidemiol Psychiatr Soc* 14:68–76.
- Angst J, Cui L, Swendsen J, Rothen S, Cravchik A, Kessler RC, Merikangas KR (2010) Major depressive disorder with sub-threshold bipolarity in the National Comorbidity Survey Replication. *Am J Psychiatry* 167:1194–1201.
- Aronson TA, Shukla S, Hoff A, Cook B (1988a) Proposed delusional depression subtypes: preliminary evidence from a retrospective study of phenomenology and treatment course. *J Affect Disord* 14:69–74.
- Aronson TA, Shukla S, Gujavarty K, Hoff A, DiBuono M, Khan E (1988b) Relapse in delusional depression: a retrospective study of the course of treatment. *Compr Psychiatry* 29:12–21.
- Astrup C, Fossum A, Holmboe R (1959) A follow-up of 270 patients with acute affective psychoses. *Acta Psychiatr Scand Suppl* 34:1–65.
- Azorin JM, Kaladjian A, Adida M, Hantouche EG, Hameg A, Lancrenon S, Akiskal HS (2008) Factors associated with rapid cycling in bipolar I manic patients: findings from a French national study. *CNS Spectr* 13:780–787.
- Baastrup PC (1964) The use of lithium in manic-depressive psychosis. *Compr Psychiatry* 5:396–408.
- Baastrup PC, Schou M (1967) Lithium as a prophylactic agents. Its effect against recurrent depressions and manic-depressive psychosis. *Arch Gen Psychiatry* 16:162–172.
- Baastrup PC, Poulsen JC, Schou M, Thomsen K, Amdisen A (1970) Prophylactic lithium: double blind discontinuation in manic-depressive and recurrent-depressive disorders. *Lancet* 2:326–330.
- Baethge C, Baldessarini RJ, Freudenthal K, Streeruwitz A, Bauer M, Bschor T (2005) Hallucinations in bipolar disorder: characteristics and comparison to unipolar depression and schizophrenia. *Bipolar Disord* 7:136–145.
- Baldessarini RJ, Undurraga J, Vazquez GH, Tondo L, Salvatore P, Ha K, Khalsa HM, Lepri B, Ha TH, Chang JS, Tohen M, Vieta E (2012) Predominant recurrence polarity among 928 adult international bipolar I disorder patients. *Acta psychiatrica Scandinavica* 125:293–302.
- Battaglia J (2005) Pharmacological management of acute agitation. *Drugs* 65:1207–1222.
- Bauer MS, Simon GE, Ludman E, Unutzer J (2005) 'Bipolarity' in bipolar disorder: distribution of manic and depressive symptoms in a treated population. *Br J Psychiatry* 187:87–88.
- Bech P (2006) The full story of lithium. A tribute to Mogens Schou (1918–2005). *Psychother Psychosom* 75:265–269.
- Begley CE, Annegers JF, Swann AC, Lewis C, Coan S, Schnapp WB, Bryant-Comstock L (2001) The lifetime cost of bipolar disorder in the US: an estimate for new cases in 1998. *Pharmacoeconomics* 19:483–495.
- Beigel A, Murphy DL (1971) Assessing clinical characteristics of the manic state. *Am J Psychiatry* 128:688–694.
- Berk M, Hallam KT, McGorry PD (2007a) The potential utility of a staging model as a course specifier: a bipolar disorder perspective. *J Affect Disord* 100:279–281.
- Berk M, Berk L, Udina M, Moylan S, Stafford L, Hallam K, Goldstone S, McGorry PD (2012) Palliative models of care for later stages of mental disorder: maximizing recovery, maintaining hope, and building morale. *Aust N Z J Psychiatry* 46:92–99.
- Berk M, Conus P, Lucas N, Hallam K, Malhi GS, Dodd S, Yatham LN, Yung A, McGorry P (2007b) Setting the stage: from prodrome to treatment resistance in bipolar disorder. *Bipolar Disord* 9:671–678.
- Bijl RV, De Graaf R, Ravelli A, Smit F, Vollebergh WA, Netherlands Mental Health S, Incidence S (2002) Gender and age-specific first incidence of DSM-III-R psychiatric disorders in the general population. Results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Soc Psychiatry Psychiatr Epidemiol* 37:372–379.
- Black DW, Nasrallah A (1989) Hallucinations and delusions in 1,715 patients with unipolar and bipolar affective disorders. *Psychopathology* 22:28–34.
- Blazer DG, Kessler RC, McGonagle KA, Swartz MS (1994) The prevalence and distribution of major depression in a national community sample: the National Comorbidity Survey. *Am J Psychiatry* 151:979–986.
- Bourdon KH, Rae DS, Locke BZ, Narrow WE, Regier DA (1992) Estimating the prevalence of mental disorders in U.S. adults from the Epidemiologic Catchment Area Survey. *Public Health Rep* 107:663–668.
- Bowman K, Raymond A (1932) A statistical study of delusions in the manic-depressive psychoses. *Am J Psychiatry* 88:111–121.
- Braunig P, Kruger S, Shugar G (1998) Prevalence and clinical significance of catatonic symptoms in mania. *Compr Psychiatry* 39:35–46.
- Brewin J, Cantwell R, Dalkin T, Fox R, Medley I, Glazebrook C, Kwiecinski R, Harrison G (1997) Incidence of schizophrenia in Nottingham. A comparison of two cohorts, 1978–80 and 1992–94. *Br J Psychiatry* 171:140–144.
- Brunetti M, Shemilt I, Pregno S, Vale L, Oxman AD, Lord J, Sisk J, Ruiz F, Hill S, Guyatt GH, Jaeschke R, Helfand M, Harbour R, Davoli M, Amato L, Liberati A, Schunemann HJ (2013) GRADE guidelines: 10. Considering resource use and rating the quality of economic evidence. *J Clin Epidemiol* 66:140–150.
- Cade J (1949) Lithium salts in the treatment of psychotic excitement. *Med J Aust* 36:349–352.
- Cade JF (2000) Lithium salts in the treatment of psychotic excitement. 1949. *Bull World Health Organ* 78:518–520.
- Caetano SC, Olvera RL, Hunter K, Hatch JP, Najt P, Bowden C, Pliszka S, Soares JC (2006) Association of psychosis with suicidality in pediatric bipolar I, II and bipolar NOS patients. *J Affect Disord* 91:33–37.
- Calabrese JR, Goldberg JF, Ketter TA, Suppes T, Frye M, White R, DeVeaugh-Geiss A, Thompson TR (2006) Recurrence in bipolar I disorder: a posthoc analysis excluding relapses in two

- double-blind maintenance studies. *Biol Psychiatry* 59:1061–1064.
- Carlson GA, Goodwin FK (1973) The stages of mania. A longitudinal analysis of the manic episode. *Arch Gen Psychiatry* 28:221–228.
- Carlson GA, Strober M (1978) Affective disorder in adolescence: issues in misdiagnosis. *J Clin Psychiatry* 39:59–66.
- Carvalho AF, Quevedo J, McIntyre RS, Soeiro-de-Souza MG, Fountoulakis KN, Berk M, Hyphantis TN, Vieta E (2014a) Treatment implications of predominant polarity and the polarity index: a comprehensive review. *Int J Neuropsychopharmacol*.
- Carvalho AF, McIntyre RS, Dimelis D, Gonda X, Berk M, Nunes-Neto PR, Cha DS, Hyphantis TN, Angst J, Fountoulakis KN (2014b) Predominant polarity as a course specifier for bipolar disorder: a systematic review. *J Affect Disord* 163:56–64.
- Casper RC, Redmond DE Jr, Katz MM, Schaffer CB, Davis JM, Koslow SH (1985) Somatic symptoms in primary affective disorder. Presence and relationship to the classification of depression. *Arch Gen Psychiatry* 42:1098–1104.
- Cassidy F, Murry E, Forest K, Carroll BJ (1998a) Signs and symptoms of mania in pure and mixed episodes. *J Affect Disord* 50:187–201.
- Cassidy F, Forest K, Murry E, Carroll BJ (1998b) A factor analysis of the signs and symptoms of mania. *Arch Gen Psychiatry* 55:27–32.
- Cipriani A, Barbui C, Rendell J, Geddes JR (2014) Clinical and regulatory implications of active run-in phases in long-term studies for bipolar disorder. *Acta Psychiatr Scand* 129:328–342.
- Cipriani A, Barbui C, Salanti G, Rendell J, Brown R, Stockton S, Purgato M, Spinelli LM, Goodwin GM, Geddes JR (2011) Comparative efficacy and acceptability of antimanic drugs in acute mania: a multiple-treatments meta-analysis. *Lancet* 378:1306–1315.
- Citrome L (2004) New treatments for agitation. *Psychiatr Q* 75:197–213.
- Clarke M, Moran P, Keogh F, Morris M, Kinsella A, Larkin C, Walsh D, O'Callaghan E (1999) Seasonal influences on admissions for affective disorder and schizophrenia in Ireland: a comparison of first and readmissions. *Eur Psychiatry* 14:251–255.
- Clayton PJ, Pitts FN, Jr. (1965) Affect disorder. IV. Mania. *Compr Psychiatry* 6:313–322.
- Colom F, Vieta E, Daban C, Pacchiarotti I, Sanchez-Moreno J (2006) Clinical and therapeutic implications of predominant polarity in bipolar disorder. *J Affect Disord* 93:13–17.
- Coryell W, Endicott J, Keller M (1992) Rapidly cycling affective disorder. Demographics, diagnosis, family history, and course. *Arch Gen Psychiatry* 49:126–131.
- Coryell W, Solomon D, Turvey C, Keller M, Leon AC, Endicott J, Schettler P, Judd L, Mueller T (2003) The long-term course of rapid-cycling bipolar disorder. *Arch Gen Psychiatry* 60:914–920.
- Cosci F, Fava GA (2013) Staging of mental disorders: systematic review. *Psychother Psychosom* 82:20–34.
- Cruz N, Vieta E, Comes M, Haro JM, Reed C, Bertsch J (2008) Rapid-cycling bipolar I disorder: course and treatment outcome of a large sample across Europe. *J Psychiatr Res* 42:1068–1075.
- Cullen B, Ward J, Graham NA, Deary IJ, Pell JP, Smith DJ, Evans JJ (2016) Prevalence and correlates of cognitive impairment in euthymic adults with bipolar disorder: A systematic review. *J Affect Disord* 205:165–181.
- Daban C, Martinez-Aran A, Torrent C, Tabares-Seisdedos R, Balanza-Martinez V, Salazar-Fraile J, Selva-Vera G, Vieta E (2006) Specificity of cognitive deficits in bipolar disorder versus schizophrenia. A systematic review. *Psychother Psychosom* 75:72–84.
- Das Gupta R, Guest JF (2002) Annual cost of bipolar disorder to UK society. *Br J Psychiatry* 180:227–233.
- Dawes M, Summerskill W, Glasziou P, Cartabellotta A, Martin J, Hopayian K, Porzsolt F, Burls A, Osborne J, Second International Conference of Evidence-Based Health Care T, Developers (2005) Sicily statement on evidence-based practice. *BMC Med Educ* 5:1.
- Delay J, Deniker P (1955) Neuroleptic effects of chlorpromazine in therapeutics of neuropsychiatry. *J Clin Exp Psychopathol* 16:104–112.
- Dilsaver SC (2011) An estimate of the minimum economic burden of bipolar I and II disorders in the United States: 2009. *J Affect Disord* 129:79–83.
- Dittmann S, Biedermann NC, Grunze H, Hummel B, Scharer LO, Kleindienst N, Forsthoff A, Matzner N, Walser S, Walden J (2002) The Stanley Foundation Bipolar Network: results of the naturalistic follow-up study after 2.5 years of follow-up in the German centres. *Neuropsychobiology* 46 Suppl 1:2–9.
- Drancourt N, Etain B, Lajnef M, Henry C, Raust A, Cochet B, Mathieu F, Gard S, Mbailara K, Zanouy L, Kahn JP, Cohen RF, Wajsbrot-Elgrabli O, Leboyer M, Scott J, Bellivier F (2013) Duration of untreated bipolar disorder: missed opportunities on the long road to optimal treatment. *Acta Psychiatr Scand* 127:136–144.
- Dunner DL, Fieve RR (1974) Clinical factors in lithium carbonate prophylaxis failure. *Arch Gen Psychiatry* 30:229–233.
- Eaton WW, Regier DA, Locke BZ, Taube CA (1981) The Epidemiologic Catchment Area Program of the National Institute of Mental Health. *Public Health Rep* 96:319–325.
- Egeland JA, Hostetter AM (1983) Amish Study, I: Affective disorders among the Amish, 1976–1980. *Am J Psychiatry* 140:56–61.
- Egeland JA, Hostetter AM, Eshleman SK, 3rd (1983) Amish Study, III: the impact of cultural factors on diagnosis of bipolar illness. *Am J Psychiatry* 140:67–71.
- Ekman M, Granstrom O, Omerov S, Jacob J, Landen M (2013) The societal cost of bipolar disorder in Sweden. *Soc Psychiatry Psychiatr Epidemiol* 48:1601–1610.
- Faedda GL, Tondo L, Teicher MH, Baldessarini RJ, Gelbard HA, Floris GF (1993) Seasonal mood disorders. Patterns of seasonal recurrence in mania and depression. *Arch Gen Psychiatry* 50:17–23.
- Faravelli C, Rosi S, Alessandra Scarpato M, Lampronti L, Amedei SG, Rana N (2006) Threshold and subthreshold bipolar disorders in the Sesto Fiorentino Study. *J Affect Disord* 94:111–119.
- Fava GA, Kellner R (1993) Staging: a neglected dimension in psychiatric classification. *Acta Psychiatr Scand* 87:225–230.
- Fisher LJ, Goldney RD, Dal Grande E, Taylor AW, Hawthorne G (2007) Bipolar disorders in Australia. A population-based study of excess costs. *Soc Psychiatry Psychiatr Epidemiol* 42:105–109.
- Fletcher SW, Spitzer WO (1980) Approach of the Canadian Task Force to the Periodic Health Examination. *Ann Intern Med* 92:253–254.
- Fountoulakis K (2015a) Long-term course. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp81–107. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015b) Historical perspective. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp1–25. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015c) Neurocognitive functioning in bipolar disorder: a comprehensive review of recent data. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression*

- (Fountoulakis K, ed), pp109–162. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015d) Personality disorders (narcissistic, anti-social, borderline). In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp197–204. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015e) Alcohol and substance use in bd patients. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp205–223. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015f) Special issues. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp659–684. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015g) Disability and overall burden related with bipolar disorder. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp361–388. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015h) Classification and epidemiology. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp341–360. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015i) Treatment guidelines. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp643–658. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015j) Comorbidity. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp225–340. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015k) Staging of bipolar disorder. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp437–459. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015l) Biological therapies. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp 461–625. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015m) Psychosocial treatment and interventions. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp627–642. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis K (2015n) Clinical description. In: *Bipolar Disorder: An Evidence-Based Guide to Manic Depression* (Fountoulakis K, ed), pp27–80. Berlin Heidelberg: Springer-Verlag.
- Fountoulakis KN (2008) The contemporary face of bipolar illness: complex diagnostic and therapeutic challenges. *CNS Spectr* 13:763–774, 777–769.
- Fountoulakis KN (2010) An update of evidence-based treatment of bipolar depression: where do we stand? *Curr Opin Psychiatry* 23:19–24.
- Fountoulakis KN, Samara MT, Siamouli M (2014) Burning issues in the meta-analysis of pharmaceutical trials for depression. *J Psychopharmacol* 28:106–117.
- Fountoulakis KN, Grunze H, Panagiotidis P, Kaprinis G (2008) Treatment of bipolar depression: an update. *J Affect Disord* 109:21–34.
- Fountoulakis KN, Gonda X, Siamouli M, Rihmer Z (2009) Psychotherapeutic intervention and suicide risk reduction in bipolar disorder: a review of the evidence. *J Affect Disord* 113:21–29.
- Fountoulakis KN, Gonda X, Vieta E, Rihmer Z (2011) Class effect of pharmacotherapy in bipolar disorder: fact or misbelief? *Ann Gen Psychiatry* 10:8.
- Fountoulakis KN, Kontis D, Gonda X, Yatham LN (2013) A systematic review of the evidence on the treatment of rapid cycling bipolar disorder. *Bipolar Disord* 15:115–137.
- Fountoulakis KN, Iacovides A, Nimatoudis I, Kaprinis G, Ierodiakonou C (1999) Comparison of the diagnosis of melancholic and atypical features according to DSM-IV and somatic syndrome according to ICD-10 in patients suffering from major depression. *Eur Psychiatry* 14:426–433.
- Fountoulakis KN, Kontis D, Gonda X, Siamouli M, Yatham LN (2012) Treatment of mixed bipolar states. *Int J Neuropsychopharmacol* 15:1015–1026.
- Fountoulakis KN, Vieta E, Sanchez-Moreno J, Kaprinis SG, Goikolea JM, Kaprinis GS (2005) Treatment guidelines for bipolar disorder: a critical review. *J Affect Disord* 86:1–10.
- Fountoulakis KN, Vieta E, Siamouli M, Valenti M, Magiria S, Oral T, Fresno D, Giannakopoulos P, Kaprinis GS (2007a) Treatment of bipolar disorder: a complex treatment for a multi-faceted disorder. *Ann Gen Psychiatry* 6:27.
- Fountoulakis KN, Bech P, Panagiotidis P, Siamouli M, Kantartzis S, Papadopoulou A, Papadopoulou M, Kaprinis S, Kourila E, Iacovides A, St Kaprinis G (2007b) Comparison of depressive indices: reliability, validity, relationship to anxiety and personality and the role of age and life events. *J Affect Disord* 97:187–195.
- Frank E, Nimgaonkar VL, Phillips ML, Kupfer DJ (2014) All the world's a (clinical) stage: rethinking bipolar disorder from a longitudinal perspective. *Mol Psychiatry*.
- Frank E, Prien RF, Jarrett RB, Keller MB, Kupfer DJ, Lavori PW, Rush AJ, Weissman MM (1991) Conceptualization and rationale for consensus definitions of terms in major depressive disorder. Remission, recovery, relapse, and recurrence. *Arch Gen Psychiatry* 48:851–855.
- Garcia-Amador M, Colom F, Valenti M, Horga G, Vieta E (2009) Suicide risk in rapid cycling bipolar patients. *J Affect Disord* 117:74–78.
- Garcia-Lopez A, De Dios-Perrino C, Ezquiaga E (2009) Polarity of the first episode and predominant polarity in a cohort of bipolar outpatients. *Eur Neuropsychopharmacol* 19:S571.
- Garriga M et al. (2016) Assessment and management of agitation in psychiatry: Expert consensus. *World J Biol Psychiatry* 17:86–128.
- Gershon S, Yuwiler A (1960) Lithium ion: a specific psychopharmacological approach to the treatment of mania. *J Neuropsychiatr* 1:229–241.
- Ghaemi SN, Pardo TB, Hsu DJ (2004) Strategies for preventing the recurrence of bipolar disorder. *J Clin Psychiatry* 65 Suppl 10:16–23.
- Goikolea JM, Colom F, Martinez-Aran A, Sanchez-Moreno J, Giordano A, Bulbena A, Vieta E (2007) Clinical and prognostic implications of seasonal pattern in bipolar disorder: a 10-year follow-up of 302 patients. *Psychol Med* 37:1595–1599.
- Goldberg JF, Harrow M, Grossman LS (1995a) Recurrent affective syndromes in bipolar and unipolar mood disorders at follow-up. *Br J Psychiatry* 166:382–385.
- Goldberg JF, Harrow M, Grossman LS (1995b) Course and outcome in bipolar affective disorder: a longitudinal follow-up study. *Am J Psychiatry* 152:379–384.
- Gonda X, Fountoulakis KN, Rihmer Z, Lazary J, Laszik A, Akiskal KK, Akiskal HS, Bagdy G (2009) Towards a genetically validated new affective temperament scale: a delineation of the temperament phenotype of 5-HTTLPR using the TEMPS-A. *J Affect Disord* 112:19–29.
- Goodwin F, Jamison K (2007) *Manic-depressive illness*, 2nd Edition. New York: Oxford University Press.
- Grande I, Berk M, Birmaher B, Vieta E (2016) *Bipolar disorder*. *Lancet* 387:1561–1572.
- Grande I, Bernardo M, Bobes J, Saiz-Ruiz J, Alamo C, Vieta E (2014) Antipsychotic switching in bipolar disorders: a systematic review. *Int J Neuropsychopharmacol* 17:497–507.

- Grunze H, Kasper S, Goodwin G, Bowden C, Moller HJ (2004) The World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for the biological treatment of bipolar disorders, part III: maintenance treatment. *World J Biol Psychiatry* 5:120–135.
- Grunze H, Kasper S, Goodwin G, Bowden C, Baldwin D, Licht R, Vieta E, Moller HJ (2002) World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for biological treatment of bipolar disorders. Part I: Treatment of bipolar depression. *World J Biol Psychiatry* 3:115–124.
- Grunze H, Kasper S, Goodwin G, Bowden C, Baldwin D, Licht RW, Vieta E, Moller HJ (2003) The World Federation of Societies of Biological Psychiatry (WFSBP) Guidelines for the Biological Treatment of Bipolar Disorders, Part II: treatment of mania. *World J Biol Psychiatry* 4:5–13.
- Guyatt G, Oxman AD, Sultan S, Brozek J, Glasziou P, Alonso-Coeillo P, Atkins D, Kunz R, Montori V, Jaeschke R, Rind D, Dahm P, Akl EA, Meerpohl J, Vist G, Berliner E, Norris S, Falck-Ytter Y, Schunemann HJ (2013) GRADE guidelines: 11. Making an overall rating of confidence in effect estimates for a single outcome and for all outcomes. *J Clin Epidemiol* 66:151–157.
- Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schunemann HJ, Group GW (2008a) What is “quality of evidence” and why is it important to clinicians? *BMJ* 336:995–998.
- Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coeillo P, Schunemann HJ, Group GW (2008b) GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 336:924–926.
- Guyatt GH, Oxman AD, Kunz R, Falck-Ytter Y, Vist GE, Liberati A, Schunemann HJ, Group GW (2008c) Going from evidence to recommendations. *BMJ* 336:1049–1051.
- Guyatt GH, Oxman AD, Vist G, Kunz R, Brozek J, Alonso-Coeillo P, Montori V, Akl EA, Djulbegovic B, Falck-Ytter Y, Norris SL, Williams JW, Jr., Atkins D, Meerpohl J, Schunemann HJ (2011a) GRADE guidelines: 4. Rating the quality of evidence--study limitations (risk of bias). *J Clin Epidemiol* 64:407–415.
- Guyatt GH, Oxman AD, Montori V, Vist G, Kunz R, Brozek J, Alonso-Coeillo P, Djulbegovic B, Atkins D, Falck-Ytter Y, Williams JW, Jr., Meerpohl J, Norris SL, Akl EA, Schunemann HJ (2011b) GRADE guidelines: 5. Rating the quality of evidence--publication bias. *J Clin Epidemiol* 64:1277–1282.
- Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, Alonso-Coeillo P, Falck-Ytter Y, Jaeschke R, Vist G, Akl EA, Post PN, Norris S, Meerpohl J, Shukla VK, Nasser M, Schunemann HJ, Group GW (2011c) GRADE guidelines: 8. Rating the quality of evidence--indirectness. *J Clin Epidemiol* 64:1303–1310.
- Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, Alonso-Coeillo P, Glasziou P, Jaeschke R, Akl EA, Norris S, Vist G, Dahm P, Shukla VK, Higgins J, Falck-Ytter Y, Schunemann HJ, Group GW (2011d) GRADE guidelines: 7. Rating the quality of evidence--inconsistency. *J Clin Epidemiol* 64:1294–1302.
- Guyatt GH, Oxman AD, Kunz R, Brozek J, Alonso-Coeillo P, Rind D, Devereaux PJ, Montori VM, Freyschuss B, Vist G, Jaeschke R, Williams JW Jr, Murad MH, Sinclair D, Falck-Ytter Y, Meerpohl J, Whittington C, Thorlund K, Andrews J, Schunemann HJ (2011e) GRADE guidelines 6. Rating the quality of evidence--imprecision. *J Clin Epidemiol* 64:1283–1293.
- Guyatt GH et al. (2011f) GRADE guidelines: 9. Rating up the quality of evidence. *J Clin Epidemiol* 64:1311–1316.
- Hajek T, Hahn M, Slaney C, Garnham J, Green J, Ruzickova M, Zvolsky P, Alda M (2008) Rapid cycling bipolar disorders in primary and tertiary care treated patients. *Bipolar Disord* 10:495–502.
- Hakkaart-van Roijen L, Hoeijenbos MB, Regeer EJ, ten Have M, Nolen WA, Veraart CP, Rutten FF (2004) The societal costs and quality of life of patients suffering from bipolar disorder in the Netherlands. *Acta Psychiatr Scand* 110:383–392.
- Helms PM, Smith RE (1983) Recurrent psychotic depression. Evidence of diagnostic stability. *J Affect Disord* 5:51–54.
- Heun R, Maier W (1993) The distinction of bipolar II disorder from bipolar I and recurrent unipolar depression: results of a controlled family study. *Acta Psychiatr Scand* 87:279–284.
- Hidalgo-Mazzei D, Undurraga J, Reinares M, Bonnin Cdel M, Saez C, Mur M, Nieto E, Vieta E (2015) The real world cost and health resource utilization associated to manic episodes: The MANACOR study. *Rev Psiquiatr Salud Ment* 8:55–64.
- Hirschfeld RM, Calabrese JR, Weissman MM, Reed M, Davies MA, Frye MA, Keck PE Jr, Lewis L, McElroy SL, McNulty JP, Wagner KD (2003a) Screening for bipolar disorder in the community. *J Clin Psychiatry* 64:53–59.
- Hirschfeld RM, Holzer C, Calabrese JR, Weissman M, Reed M, Davies M, Frye MA, Keck P, McElroy S, Lewis L, Tierce J, Wagner KD, Hazard E (2003b) Validity of the mood disorder questionnaire: a general population study. *Am J Psychiatry* 160:178–180.
- Hopewell S, Clarke M, Moher D, Wager E, Middleton P, Altman DG, Schulz KF, Group C (2008) CONSORT for reporting randomized controlled trials in journal and conference abstracts: explanation and elaboration. *PLoS Med* 5:e20.
- Horsley T, Hyde C, Santesso N, Parkes J, Milne R, Stewart R (2011) Teaching critical appraisal skills in healthcare settings. *Cochrane Database Syst Rev* (Online):CD001270.
- Hostetter AM, Egeland JA, Endicott J (1983) Amish Study, II: consensus diagnoses and reliability results. *Am J Psychiatry* 140:62–66.
- Huf W, Kalcher K, Kasper S (2011a) Widespread methodological problems limit validity of meta-analytic results. *Psychother Psychosom* 80:246; author reply 247–248.
- Huf W, Kalcher K, Pail G, Friedrich ME, Filzmoser P, Kasper S (2011b) Meta-analysis: fact or fiction? How to interpret meta-analyses. *World J Biol Psychiatry* 12:188–200.
- Hunt N, Sayer H, Silverstone T (1992) Season and manic relapse. *Acta Psychiatr Scand* 85:123–126.
- Ivers N, Jamtvedt G, Flottorp S, Young JM, Odgaard-Jensen J, French SD, O'Brien MA, Johansen M, Grimshaw J, Oxman AD (2012) Audit and feedback: effects on professional practice and healthcare outcomes. *Cochrane Database Syst Rev* (Online) 6:CD000259.
- Jaeschke R, Guyatt GH, Dellinger P, Schunemann H, Levy MM, Kunz R, Norris S, Bion J, Group GW (2008) Use of GRADE grid to reach decisions on clinical practice guidelines when consensus is elusive. *BMJ* 337:a744.
- Jamtvedt G, Young JM, Kristoffersen DT, Thomson O'Brien MA, Oxman AD (2003) Audit and feedback: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev* (Online):CD000259.
- Jamtvedt G, Young JM, Kristoffersen DT, O'Brien MA, Oxman AD (2006a) Does telling people what they have been doing change what they do? A systematic review of the effects of audit and feedback. *Qual Saf Health Care* 15:433–436.
- Jamtvedt G, Young JM, Kristoffersen DT, O'Brien MA, Oxman AD (2006b) Audit and feedback: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev* (Online):CD000259.
- Jaspers K (1913) *General Psychopathology*. Chicago: The Johns Hopkins University Press (1997).

- Johnstone EC, Crow TJ, Frith CD, Owens DG (1988) The Northwick Park "functional" psychosis study: diagnosis and treatment response. *Lancet* 2:119–125.
- Judd LL, Akiskal HS (2003) The prevalence and disability of bipolar spectrum disorders in the US population: re-analysis of the ECA database taking into account subthreshold cases. *J Affect Disord* 73:123–131.
- Judd LL, Akiskal HS, Schettler PJ, Coryell W, Endicott J, Maser JD, Solomon DA, Leon AC, Keller MB (2003) A prospective investigation of the natural history of the long-term weekly symptomatic status of bipolar II disorder. *Arch Gen Psychiatry* 60:261–269.
- Kapczinski F, Dias VV, Kauer-Sant'Anna M, Frey BN, Grassi-Oliveira R, Colom F, Berk M (2009) Clinical implications of a staging model for bipolar disorders. *Expert Rev Neurother* 9:957–966.
- Kapczinski F et al. (2014) Staging systems in bipolar disorder: an International Society for Bipolar Disorders Task Force Report. *Acta Psychiatr Scand*.
- Keck PE Jr, McElroy SL, Strakowski SM, West SA, Sax KW, Hawkins JM, Bourne ML, Haggard P (1998) 12-month outcome of patients with bipolar disorder following hospitalization for a manic or mixed episode. *Am J Psychiatry* 155:646–652.
- Keck PE Jr, McElroy SL, Havens JR, Altshuler LL, Nolen WA, Frye MA, Suppes T, Denicoff KD, Kupka R, Leverich GS, Rush AJ, Post RM (2003) Psychosis in bipolar disorder: phenomenology and impact on morbidity and course of illness. *Compr Psychiatry* 44:263–269.
- Kendler KS, Gallagher TJ, Abelson JM, Kessler RC (1996) Lifetime prevalence, demographic risk factors, and diagnostic validity of nonaffective psychosis as assessed in a US community sample. The National Comorbidity Survey. *Arch Gen Psychiatry* 53:1022–1031.
- Kessler RC, Merikangas KR (2004) The National Comorbidity Survey Replication (NCS-R): background and aims. *Int J Methods Psychiatr Res* 13:60–68.
- Kessler RC, McGonagle KA, Swartz M, Blazer DG, Nelson CB (1993) Sex and depression in the National Comorbidity Survey. I: lifetime prevalence, chronicity and recurrence. *J Affect Disord* 29:85–96.
- Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995) Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 52:1048–1060.
- Kessler RC, Rubinow DR, Holmes C, Abelson JM, Zhao S (1997a) The epidemiology of DSM-III-R bipolar I disorder in a general population survey. *Psychol Med* 27:1079–1089.
- Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE (2005a) Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62:617–627.
- Kessler RC, McGonagle KA, Nelson CB, Hughes M, Swartz M, Blazer DG (1994a) Sex and depression in the National Comorbidity Survey. II: Cohort effects. *J Affect Disord* 30:15–26.
- Kessler RC, Nelson CB, McGonagle KA, Liu J, Swartz M, Blazer DG (1996) Comorbidity of DSM-III-R major depressive disorder in the general population: results from the US National Comorbidity Survey. *Br J Psychiatry Suppl*:17–30.
- Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE (2005b) Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62:593–602.
- Kessler RC, Akiskal HS, Angst J, Guyer M, Hirschfeld RM, Merikangas KR, Stang PE (2006) Validity of the assessment of bipolar spectrum disorders in the WHO CIDI 3.0. *J Affect Disord* 96:259–269.
- Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS (1994b) Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Arch Gen Psychiatry* 51:8–19.
- Kessler RC, Anthony JC, Blazer DG, Bromet E, Eaton WW, Kendler K, Swartz M, Wittchen HU, Zhao S (1997b) The US National Comorbidity Survey: overview and future directions. *Epidemiol Psychiatr Soc* 6:4–16.
- Kessler RC, Avenevoli S, Costello J, Green JG, Gruber MJ, McLaughlin KA, Petukhova M, Sampson NA, Zaslavsky AM, Merikangas KR (2012a) Severity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication Adolescent Supplement. *Arch Gen Psychiatry* 69:381–389.
- Kessler RC, Berglund P, Chiu WT, Demler O, Heeringa S, Hiripi E, Jin R, Pennell BE, Walters EE, Zaslavsky A, Zheng H (2004) The US National Comorbidity Survey Replication (NCS-R): design and field procedures. *Int J Methods Psychiatr Res* 13:69–92.
- Kessler RC, Avenevoli S, Costello EJ, Georgiades K, Green JG, Gruber MJ, He JP, Koretz D, McLaughlin KA, Petukhova M, Sampson NA, Zaslavsky AM, Merikangas KR (2012b) Prevalence, persistence, and sociodemographic correlates of DSM-IV disorders in the National Comorbidity Survey Replication Adolescent Supplement. *Arch Gen Psychiatry* 69:372–380.
- Kety S, Rosenthal D, Wender P, Schulschinger F (1968) The types and prevalence of mental illness in the biological and adoptive families of adopted schizophrenics. *J Psychiatr Res*:345–362.
- Kleist K (1937) Zustandsbilder und Krankheitsarten im Lichte der Gehirnpathologie. *Psychiatrisch-neurologische Wochenschrift*:420–422.
- Klerman GL (1981) The spectrum of mania. *Compr Psychiatry* 22:11–20.
- Klerman GL (1987) The classification of bipolar disorders. *Psychiatric Annals* 17:13–17.
- Kotin J, Goodwin FK (1972) Depression during mania: clinical observations and theoretical implications. *Am J Psychiatry* 129:679–686.
- Koukopoulos A (2006) The primacy of mania. In: *Bipolar Psychopharmacology: Caring for the patient* (Akiskal H, Tohen M, eds). New York: John Wiley & Sons.
- Kraepelin E (1921) *Manic-Depressive Insanity and Paranoia*. Edinburgh: Livingstone.
- Kramlinger KG, Post RM (1996) Ultra-rapid and ultradian cycling in bipolar affective illness. *Br J Psychiatry* 168:314–323.
- Kruger S, Cooke RG, Spegg CC, Braunig P (2003) Relevance of the catatonic syndrome to the mixed manic episode. *J Affect Disord* 74:279–285.
- Kuhn R (1958) The treatment of depressive states with G 22355 (imipramine hydrochloride). *Am J Psychiatry* 115:459–464.
- Kukopulos A, Reginaldi D, Laddomada P, Floris G, Serra G, Tondo L (1980) Course of the manic-depressive cycle and changes caused by treatment. *Pharmakopsychiatr Neuropsychopharmakol* 13:156–167.
- Lambert PA, Carraz G, Borselli S, Carbel S (1966) [Neuropsychotropic action of a new anti-epileptic agent: depamide]. *Ann Med Psychol (Paris)* 124:707–710.
- Lange J (1922) *Katatonsche erscheinungen im rahmen manischer erkrankungen*. Berlin: Julius Springer.
- Lee HC, Tsai SY, Lin HC (2007) Seasonal variations in bipolar disorder admissions and the association with climate: a population-based study. *J Affect Disord* 97:61–69.
- Lee S, Tsang A, Kessler RC, Jin R, Sampson N, Andrade L, Karam EG, Mora ME, Merikangas K, Nakane Y, Popovici DG, Posada-

- Villa J, Sagar R, Wells JE, Zarkov Z, Petukhova M (2010) Rapid-cycling bipolar disorder: cross-national community study. *Br J Psychiatry* 196:217–225.
- Leff JP, Fischer M, Bertelsen A (1976) A cross-national epidemiological study of mania. *Br J Psychiatry* 129:428–442.
- Lehman AF, Steinwachs DM (1998) Translating research into practice: the Schizophrenia Patient Outcomes Research Team (PORT) treatment recommendations. *Schizophr Bull* 24:1–10.
- Lenox RH, Watson DG (1994) Lithium and the brain: a psychopharmacological strategy to a molecular basis for manic depressive illness. *Clin Chem* 40:309–314.
- Leonhard K (1957a) [Pathogenesis of manic-depressive disease]. *Nervenarzt* 28:271–272.
- Leonhard K (1957b) [Cycloid psychoses, often erroneously considered as schizophrenia]. *Psychiatr Neurol Med Psychol (Leipz)* 9:359–365.
- Leonhard K (1963) Die prapsychotische Temperamente bei den monopolen und bipolaren phasischen Psychosen. *Psychiat et Neurol (Basel)* 146:105–115.
- Leonhard K (1979) *The classification of endogenous psychoses*, 5th ed. New York: Irvington.
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D (2009) The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 339:b2700.
- Lin A, Reniers RL, Wood SJ (2013) Clinical staging in severe mental disorder: evidence from neurocognition and neuroimaging. *Br J Psychiatry Suppl* 54:s11–17.
- Loudon JB, Blackburn IM, Ashworth CM (1977) A study of the symptomatology and course of manic illness using a new scale. *Psychol Med* 7:723–729.
- Magee WJ, Eaton WW, Wittchen HU, McGonagle KA, Kessler RC (1996) Agoraphobia, simple phobia, and social phobia in the National Comorbidity Survey. *Arch Gen Psychiatry* 53:159–168.
- Marder SR (2006) A review of agitation in mental illness: treatment guidelines and current therapies. *J Clin Psychiatry* 67 Suppl 10:13–21.
- Martinez-Aran A, Vieta E, Torrent C, Sanchez-Moreno J, Goikolea JM, Salameo M, Malhi GS, Gonzalez-Pinto A, Daban C, Alvarez-Grandi S, Fountoulakis K, Kaprinis G, Tabares-Seisdedos R, Ayuso-Mateos JL (2007) Functional outcome in bipolar disorder: the role of clinical and cognitive factors. *Bipolar Disord* 9:103–113.
- Mazzarini L, Colom F, Pacchiarotti I, Nivoli AM, Murru A, Bonnin CM, Cruz N, Sanchez-Moreno J, Kotzalidis GD, Girardi P, Tatarelli R, Vieta E (2010) Psychotic versus non-psychotic bipolar II disorder. *J Affect Disord* 126:55–60.
- Mazzarini L, Pacchiarotti I, Colom F, Sani G, Kotzalidis GD, Rosa AR, Sanna L, De Rossi P, Girardi N, Bonnin CM, Sanchez-Moreno J, Vazquez GH, Gasto C, Tatarelli R, Vieta E (2009) Predominant polarity and temperament in bipolar and unipolar affective disorders. *J Affect Disord* 119:28–33.
- McAlister FA, Graham I, Karr GW, Laupacis A (1999a) Evidence-based medicine and the practicing clinician. *J Gen Intern Med* 14:236–242.
- McAlister FA, Laupacis A, Wells GA, Sackett DL (1999b) *Users' Guides to the Medical Literature: XIX. Applying clinical trial results B. Guidelines for determining whether a drug is exerting (more than) a class effect.* *JAMA* 282:1371–1377.
- McCrone P, Dhanasiri S, Patel A, Knapp M, Lawton-Simth S (2008) *Paying the Price*. London: The King's Fund.
- McGorry PD (2007) Issues for DSM-V: clinical staging: a heuristic pathway to valid nosology and safer, more effective treatment in psychiatry. *Am J Psychiatry* 164:859–860.
- McGorry PD (2010a) Risk syndromes, clinical staging and DSM V: new diagnostic infrastructure for early intervention in psychiatry. *Schizophr Res* 120:49–53.
- McGorry PD (2010b) Staging in neuropsychiatry: a heuristic model for understanding, prevention and treatment. *Neurotox Res* 18:244–255.
- McGorry PD, Nelson B, Goldstone S, Yung AR (2010) Clinical staging: a heuristic and practical strategy for new research and better health and social outcomes for psychotic and related mood disorders. *Can J Psychiatry* 55:486–497.
- McGorry PD, Hickie IB, Yung AR, Pantelis C, Jackson HJ (2006) Clinical staging of psychiatric disorders: a heuristic framework for choosing earlier, safer and more effective interventions. *Aust N Z J Psychiatry* 40:616–622.
- McGorry PD, Purcell R, Hickie IB, Yung AR, Pantelis C, Jackson HJ (2007) Clinical staging: a heuristic model for psychiatry and youth mental health. *Med J Aust* 187:S40–42.
- Merikangas KR, Cui L, Kattan G, Carlson GA, Youngstrom EA, Angst J (2012) Mania with and without depression in a community sample of US adolescents. *Arch Gen Psychiatry* 69:943–951.
- Merikangas KR, Akiskal HS, Angst J, Greenberg PE, Hirschfeld RM, Petukhova M, Kessler RC (2007) Lifetime and 12-month prevalence of bipolar spectrum disorder in the National Comorbidity Survey replication. *Arch Gen Psychiatry* 64:543–552.
- Merikangas KR, Jin R, He JP, Kessler RC, Lee S, Sampson NA, Viana MC, Andrade LH, Hu C, Karam EG, Ladea M, Medina-Mora ME, Ono Y, Posada-Villa J, Sagar R, Wells JE, Zarkov Z (2011) Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative. *Arch Gen Psychiatry* 68:241–251.
- Mismetti P, Cucherat M, Laporte S (2007) [Meta-analyses or meta-analyses: same level of proof?]. *Presse Med* 36:524–530.
- Mitchell PB, Hadzi-Pavlovic D (2000) Lithium treatment for bipolar disorder. *Bull World Health Organ* 78:515–517.
- Mitchell PB, Slade T, Andrews G (2004) Twelve-month prevalence and disability of DSM-IV bipolar disorder in an Australian general population survey. *Psychol Med* 34:777–785.
- Mitchell S (1870) On the use of bromide of lithium. *Am J Med Sci* 60:443–445.
- Mitchell S (1877) Clinical lecture on nervousness in the male. *Med News* 35:177–184.
- Moher D, Liberati A, Tetzlaff J, Altman DG, Group P (2009a) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 6:e1000097.
- Moher D, Liberati A, Tetzlaff J, Altman DG, Group P (2009b) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 339:b2535.
- Moreno DH, Andrade LH (2005) The lifetime prevalence, health services utilization and risk of suicide of bipolar spectrum subjects, including subthreshold categories in the Sao Paulo ECA study. *J Affect Disord* 87:231–241.
- Mulder RT, Cosgriff JP, Smith AM, Joyce PR (1990) Seasonality of mania in New Zealand. *Aust N Z J Psychiatry* 24:187–190.
- Mur M, Portella MJ, Martinez-Aran A, Pifarre J, Vieta E (2007) Persistent neuropsychological deficit in euthymic bipolar patients: executive function as a core deficit. *J Clin Psychiatry* 68:1078–1086.
- Murphy DL, Beigel A (1974) Depression, elation, and lithium carbonate responses in manic patient subgroups. *Arch Gen Psychiatry* 31:643–648.

- Murray G, Lam RW, Beaulieu S, Sharma V, Cervantes P, Parikh SV, Yatham LN (2011) Do symptoms of bipolar disorder exhibit seasonal variation? A multisite prospective investigation. *Bipolar Disord* 13:687–695.
- Negash A, Alem A, Kebede D, Deyessa N, Shibre T, Kullgren G (2005) Prevalence and clinical characteristics of bipolar I disorder in Butajira, Ethiopia: a community-based study. *J Affect Disord* 87:193–201.
- Nelson WH, Khan A, Orr WW Jr. (1984) Delusional depression. Phenomenology, Neuroendocrine function, and tricyclic antidepressant response. *J Affect Disord* 6:297–306.
- Ng B, Camacho A, Lara DR, Brunstein MG, Pinto OC, Akiskal HS (2007) A case series on the hypothesized connection between dementia and bipolar spectrum disorders: Bipolar type VI? *J Affect Disord* 107:307–315.
- Nierenberg AA, Akiskal HS, Angst J, Hirschfeld RM, Merikangas KR, Petukhova M, Kessler RC (2010) Bipolar disorder with frequent mood episodes in the national comorbidity survey replication (NCS-R). *Mol Psychiatry* 15:1075–1087.
- Nivoli AMA, Pacchiarotti I, Rosa AR, Popovic D, Murru A, Valenti M, Bonnin CM, Grande I, Sanchez-Moreno J, Vieta E, Colom F (2011) Gender differences in a cohort study of 604 bipolar patients: The role of predominant polarity. *J Affect Disord* 133:443–449.
- Noble JH Jr. (2006) Meta-analysis: Methods, strengths, weaknesses, and political uses. *J Lab Clin Med* 147:7–20.
- Nordstrom K, Allen MH (2007) Managing the acutely agitated and psychotic patient. *CNS Spectr* 12:5–11.
- Nordstrom K, Zun LS, Wilson MP, Md VS, Ng AT, Bregman B, Anderson EL (2012) Medical evaluation and triage of the agitated patient: consensus statement of the American association for emergency psychiatry project Beta medical evaluation workgroup. *West J Emerg Med* 13:3–10.
- Nurnberger J Jr, Guroff JJ, Hamovit J, Berrettini W, Gershon E (1988) A family study of rapid-cycling bipolar illness. *J Affect Disord* 15:87–91.
- OCEBM Levels of Evidence Working Group The Oxford 2011 Levels of Evidence <http://www.cebm.net/index.aspx?o=5653>. In: Oxford Centre for Evidence-Based Medicine.
- Okasha A, Okasha T (2000) Notes on mental disorders in pharaonic Egypt. *Hist Psychiatry* XI:413–424.
- Okuma T, Inanaga K, Otsuki S, Sarai K, Takahashi R, Hazama H, Mori A, Watanabe M (1979) Comparison of the antimanic efficacy of carbamazepine and chlorpromazine: a double-blind controlled study. *Psychopharmacology (Berl)* 66:211–217.
- Oliver JM, Simmons ME (1985) Affective disorders and depression as measured by the Diagnostic Interview Schedule and the Beck Depression Inventory in an unselected adult population. *J Clin Psychol* 41:469–477.
- Oxman AD, Guyatt GH (1988) Guidelines for reading literature reviews. *CMAJ* 138:697–703.
- Ozyildirim I, Cakir S, Yazici O (2010) Impact of psychotic features on morbidity and course of illness in patients with bipolar disorder. *Eur Psychiatry* 25:47–51.
- Pacchiarotti I, Nivoli AMA, Mazzarini L, Kotzalidis GD, Sani G, Koukopoulos A, Scott J, Strejilevich S, Sanchez-Moreno J, Murru A, Valenti M, Girardi P, Vieta E, Colom F (2013a) The symptom structure of bipolar acute episodes: in search for the mixing link. *J Affect Disord* 149:56–66.
- Pacchiarotti I et al. (2013b) The International Society for Bipolar Disorders (ISBD) task force report on antidepressant use in bipolar disorders. *Am J Psychiatry* 170:1249–1262.
- Paris J, Gunderson J, Weinberg I (2007) The interface between borderline personality disorder and bipolar spectrum disorders. *Compr Psychiatry* 48:145–154.
- Parker G, Walter S (1982) Seasonal variation in depressive disorders and suicidal deaths in New South Wales. *Br J Psychiatry* 140:626–632.
- Parkes J, Hyde C, Deeks J, Milne R (2001) Teaching critical appraisal skills in health care settings. *Cochrane Database Syst Rev* (Online):CD001270.
- Partonen T, Lonnqvist J (1996) Seasonal variation in bipolar disorder. *Br J Psychiatry* 169:641–646.
- Patten SB, Paris J (2008) The bipolar spectrum--a bridge too far? *Can J Psychiatry* 53:762–768.
- Paul C, Gallini A, Archier E, Castela E, Devaux S, Aractingi S, Aubin F, Bachelez H, Cribier B, Joly P, Jullien D, Le Maitre M, Misery L, Richard MA, Ortonne JP (2012) Evidence-based recommendations on topical treatment and phototherapy of psoriasis: systematic review and expert opinion of a panel of dermatologists. *Journal of the European Academy of Dermatology and Venereology: JEADV* 26 Suppl 3:1–10.
- Peck DF (1990) Climatic variables and admissions for mania: a reanalysis. *J Affect Disord* 20:249–250.
- Perlis RH, Miyahara S, Marangell LB, Wisniewski SR, Ostacher M, DelBello MP, Bowden CL, Sachs GS, Nierenberg AA, Investigators S-B (2004) Long-term implications of early onset in bipolar disorder: data from the first 1000 participants in the systematic treatment enhancement program for bipolar disorder (STEP-BD). *Biol Psychiatry* 55:875–881.
- Perris C, d'Elia G (1966a) A study of bipolar (manic-depressive) and unipolar recurrent depressive psychoses. X. Mortality, suicide and life-cycles. *Acta Psychiatr Scand Suppl* 194:172–189.
- Perris C, d'Elia G (1966b) A study of bipolar (manic-depressive) and unipolar recurrent depressive psychoses. IX. therapy and prognosis. *Acta Psychiatr Scand Suppl* 194:153–171.
- Pocock SJ, Elbourne DR (2000) Randomized trials or observational tribulations? *N Engl J Med* 342:1907–1909.
- Popovic D, Reinares M, Goikolea JM, Bonnin CM, Gonzalez-Pinto A, Vieta E (2012) Polarity index of pharmacological agents used for maintenance treatment of bipolar disorder. *Eur Neuropsychopharmacol* 22:339–346.
- Post RM (2010) Mechanisms of illness progression in the recurrent affective disorders. *Neurotox Res* 18:256–271.
- Post RM, Fleming J, Kapczinski F (2012) Neurobiological correlates of illness progression in the recurrent affective disorders. *J Psychiatr Res* 46:561–573.
- Post RM, Luckenbaugh DA, Leverich GS, Altshuler LL, Frye MA, Suppes T, Keck PE, McElroy SL, Nolen WA, Kupka R, Grunze H, Walden J (2008) Incidence of childhood-onset bipolar illness in the USA and Europe. *Br J Psychiatry* 192:150–151.
- Post RM, Altshuler LL, Frye MA, Suppes T, Keck PE, Jr., McElroy SL, Leverich GS, Luckenbaugh DA, Rowe M, Pizzarello S, Kupka RW, Grunze H, Nolen WA (2010) Complexity of pharmacologic treatment required for sustained improvement in outpatients with bipolar disorder. *J Clin Psychiatry* 71:1176–1186; quiz 1252–1173.
- Prien RF, Himmelhoch JM, Kupfer DJ (1988) Treatment of mixed mania. *J Affect Disord* 15:9–15.
- Quitkin FM, Rabkin JG, Prien RF (1986) Bipolar disorder: are there manic-prone and depressive-prone forms? *J Clin Psychopharmacol* 6:167–172.
- Regeer EJ, Rosso ML, ten Have M, Vollebergh W, Nolen WA (2002) Prevalence of bipolar disorder: a further study in The Netherlands. *Bipolar Disord* 4 Suppl 1:37–38.
- Regier DA, Myers JK, Kramer M, Robins LN, Blazer DG, Hough RL, Eaton WW, Locke BZ (1984) The NIMH Epidemiologic Catchment Area program. Historical context, major objectives,

- and study population characteristics. *Arch Gen Psychiatry* 41:934–941.
- Regier DA, Farmer ME, Rae DS, Myers JK, Kramer M, Robins LN, George LK, Karno M, Locke BZ (1993) One-month prevalence of mental disorders in the United States and sociodemographic characteristics: the Epidemiologic Catchment Area study. *Acta Psychiatr Scand* 88:35–47.
- Regier DA, Boyd JH, Burke JD, Jr., Rae DS, Myers JK, Kramer M, Robins LN, George LK, Karno M, Locke BZ (1988) One-month prevalence of mental disorders in the United States. Based on five Epidemiologic Catchment Area sites. *Arch Gen Psychiatry* 45:977–986.
- Reinares M, Sanchez-Moreno J, Fountoulakis KN (2014) Psychosocial interventions in bipolar disorder: what, for whom, and when. *J Affect Disord* 156:46–55.
- Rennie T (1942) Prognosis in manic-depressive psychosis. *Am J Psychiatry* 98:801–814.
- Richardson WS, Wilson MC, Nishikawa J, Hayward RS (1995) The well-built clinical question: a key to evidence-based decisions. *ACP J club* 123:A12–13.
- Rosa AR, Andreazza AC, Kunz M, Gomes F, Santin A, Sanchez-Moreno J, Reinares M, Colom F, Vieta E, Kapczinski F (2008) Predominant polarity in bipolar disorder: diagnostic implications. *J Affect Disord* 107:45–51.
- Rosenberg WM, Deeks J, Lusher A, Snowball R, Dooley G, Sackett D (1998) Improving searching skills and evidence retrieval. *J R Coll Physicians Lond* 32:557–563.
- Rosenthal NE, Rosenthal LN, Stallone F, Dunner DL, Fieve RR (1980) Toward the validation of RDC schizoaffective disorder. *Arch Gen Psychiatry* 37:804–810.
- Runge C, Grunze H (2004) [Annual costs of bipolar disorders in Germany]. *Nervenarzt* 75:896–903.
- Schioldann J (1999) John Cade's seminal lithium paper turns fifty. *Acta Psychiatr Scand* 100:403–405.
- Schioldann J (2006) Obituary: Mogens Abelin Schou (1918–2005): half a century with lithium. *Hist Psychiatry* 17:247–252.
- Schioldann J (2011) 'On periodical depressions and their pathogenesis' by Carl Lange (1886). *Hist Psychiatry* 22:108–130.
- Schlosser RW, Koul R, Costello J (2007) Asking well-built questions for evidence-based practice in augmentative and alternative communication. *J Commun Disord* 40:225–238.
- Schneck CD, Miklowitz DJ, Miyahara S, Araga M, Wisniewski S, Gyulai L, Allen MH, Thase ME, Sachs GS (2008) The prospective course of rapid-cycling bipolar disorder: findings from the STEP-BD. *Am J Psychiatry* 165:370–377.
- Schneck CD, Miklowitz DJ, Calabrese JR, Allen MH, Thomas MR, Wisniewski SR, Miyahara S, Shelton MD, Ketter TA, Goldberg JF, Bowden CL, Sachs GS (2004) Phenomenology of rapid-cycling bipolar disorder: data from the first 500 participants in the Systematic Treatment Enhancement Program. *Am J Psychiatry* 161:1902–1908.
- Schou M (1963) Normothymotics, "mood-normalizers": are lithium and the imipramine drugs specific for affective disorders? *Br J Psychiatry* 109:803–809.
- Schou M, Juel-Nielsen N, Stromgren E, Voldby H (1954) The treatment of manic psychoses by the administration of lithium salts. *J Neurol Neurosurg Psychiatry* 17:250–260.
- Schou M, Baastup PC, Grof P, Weis P, Angst J (1970) Pharmacological and clinical problems of lithium prophylaxis. *Br J Psychiatry* 116:615–619.
- Schunemann HJ, Fretheim A, Oxman AD (2006) Improving the use of research evidence in guideline development: 10. Integrating values and consumer involvement. *Health Res Policy Syst* 4:22.
- Serretti A, Olgiati P (2005) Profiles of "manic" symptoms in bipolar I, bipolar II and major depressive disorders. *J Affect Disord* 84:159–166.
- Serretti A, Mandelli L, Lattuada E, Cusin C, Smeraldi E (2002) Clinical and demographic features of mood disorder subtypes. *Psychiatry Res* 112:195–210.
- Shand AJ, Scott NW, Anderson SM, Eagles JM (2011) The seasonality of bipolar affective disorder: comparison with a primary care sample using the Seasonal Pattern Assessment Questionnaire. *J Affect Disord* 132:289–292.
- Sherman M, Burak K, Maroun J, Metrakos P, Knox JJ, Myers RP, Guindi M, Porter G, Kachura JR, Rasuli P, Gill S, Ghali P, Chaudhury P, Siddiqui J, Valenti D, Weiss A, Wong R (2011) Multidisciplinary Canadian consensus recommendations for the management and treatment of hepatocellular carcinoma. *Curr Oncol* 18:228–240.
- Strakowski SM, Keck PE Jr, McElroy SL, West SA, Sax KW, Hawkins JM, Kmetz GF, Upadhyaya VH, Tugrul KC, Bourne ML (1998) Twelve-month outcome after a first hospitalization for affective psychosis. *Arch Gen Psychiatry* 55:49–55.
- Stringaris A, Santosh P, Leibenluft E, Goodman R (2010) Youth meeting symptom and impairment criteria for mania-like episodes lasting less than four days: an epidemiological enquiry. *J Child Psychol Psychiatry* 51:31–38.
- Suppes T (2002) Review of the use of topiramate for treatment of bipolar disorders. *J Clin Psychopharmacol* 22:599–609.
- Szadoczky E, Papp Z, Vitrai J, Rihmer Z, Furedi J (1998) The prevalence of major depressive and bipolar disorders in Hungary. Results from a national epidemiologic survey. *J Affect Disord* 50:153–162.
- Taylor MA, Abrams R (1973) The phenomenology of mania. A new look at some old patients. *Arch Gen Psychiatry* 29:520–522.
- Taylor MA, Abrams R (1977) Catatonia. Prevalence and importance in the manic phase of manic-depressive illness. *Arch Gen Psychiatry* 34:1223–1225.
- ten Have M, Vollebergh W, Bijl R, Nolen WA (2002) Bipolar disorder in the general population in The Netherlands (prevalence, consequences and care utilisation): results from The Netherlands Mental Health Survey and Incidence Study (NEMESIS). *J Affect Disord* 68:203–213.
- Tohen M, Waternaux CM, Tsuang MT (1990) Outcome in mania. A 4-year prospective follow-up of 75 patients utilizing survival analysis. *Arch Gen Psychiatry* 47:1106–1111.
- Tohen M, Frank E, Bowden CL, Colom F, Ghaemi SN, Yatham LN, Malhi GS, Calabrese JR, Nolen WA, Vieta E, Kapczinski F, Goodwin GM, Suppes T, Sachs GS, Chengappa KR, Grunze H, Mitchell PB, Kanba S, Berk M (2009) The International Society for Bipolar Disorders (ISBD) Task Force report on the nomenclature of course and outcome in bipolar disorders. *Bipolar Disord* 11:453–473.
- Tonelli MR (1999) In defense of expert opinion. *Acad Med* 74:1187–1192.
- Torrent C et al. (2013) Efficacy of functional remediation in bipolar disorder: a multicenter randomized controlled study. *Am J Psychiatry* 170:852–859.
- Tsitsipa E, Fountoulakis KN (2015) The neurocognitive functioning in bipolar disorder: a systematic review of data. *Ann Gen Psychiatry* 14:42.
- U.S. Preventive Services Task Force (1989) Guide to clinical preventive services: report of the U.S. Preventive Services Task Force.: DIANE Publishing.
- Vieta E (2015) Staging and psychosocial early intervention in bipolar disorder. *Lancet Psychiatry* 2:483–485.
- Vieta E, Reinares M, Rosa AR (2011) Staging bipolar disorder. *Neurotox Res* 19:279–285.

- Vieta E, Berk M, Wang W, Colom F, Tohen M, Baldessarini RJ (2009) Predominant previous polarity as an outcome predictor in a controlled treatment trial for depression in bipolar I disorder patients. *J Affect Disord* 119:22–27.
- Vrublevska J, Fountoulakis K (2015) Medical comorbidity in bipolar disorder. In: *The Bipolar Book. History, Neurobiology, and Treatment* (Yildiz A, Ruiz P, Nemeroff C, eds), pp497–506. Oxford: Oxford University Press.
- Walter SD (1977) Seasonality of mania: a reappraisal. *Br J Psychiatry* 131:345–350.
- Warner LA, Kessler RC, Hughes M, Anthony JC, Nelson CB (1995) Prevalence and correlates of drug use and dependence in the United States. Results from the National Comorbidity Survey. *Arch Gen Psychiatry* 52:219–229.
- Wehr TA, Goodwin FK (1987) Can antidepressants cause mania and worsen the course of affective illness? *Am J Psychiatry* 144:1403–1411.
- Wehr TA, Sack DA, Rosenthal NE, Cowdry RW (1988) Rapid cycling affective disorder: contributing factors and treatment responses in 51 patients. *Am J Psychiatry* 145:179–184.
- Weissman MM, Myers JK (1978) Affective disorders in a US urban community: the use of research diagnostic criteria in an epidemiological survey. *Arch Gen Psychiatry* 35:1304–1311.
- Weissman MM, Bland RC, Canino GJ, Faravelli C, Greenwald S, Hwu HG, Joyce PR, Karam EG, Lee CK, Lellouch J, Lepine JP, Newman SC, Rubio-Stipec M, Wells JE, Wickramaratne PJ, Wittchen H, Yeh EK (1996) Cross-national epidemiology of major depression and bipolar disorder. *JAMA* 276:293–299.
- Weygandt W (1899) *Über die mischzustände des manisch-depressiven irreseins*. Munich: JF Lechman.
- WHO (1992) *The ICD-10 classification of mental and behavioural disorders - clinical descriptions and diagnostic guidelines*. Geneva: World Health Organization.
- WHO (1994) *Diagnostic and Management Guidelines for Mental Disorders in Primary Health Care*. Seattle, WA: Hogrefe & Huber.
- WHO (2008) *The global burden of disease: 2004 update*. Geneva: World Health Organization.
- Wicki W, Angst J (1991) The Zurich Study. X. Hypomania in a 28- to 30-year-old cohort. *Eur Arch Psychiatry Clin Neurosci* 240:339–348.
- Winokur G (1984) Psychosis in bipolar and unipolar affective illness with special reference to schizo-affective disorder. *Br J Psychiatry* 145:236–242.
- Winokur G, Tsuang MT (1975) Elation versus irritability in mania. *Compr Psychiatry* 16:435–436.
- Winokur G, Clayton P, Reich T (1969) *Manic Depressive Illness*. Saint Louis: CV Mosby.
- Wittchen HU, Zhao S, Kessler RC, Eaton WW (1994) DSM-III-R generalized anxiety disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 51:355–364.
- Wyatt RJ, Henter I (1995) An economic evaluation of manic-depressive illness--1991. *Soc Psychiatry Psychiatr Epidemiol* 30:213–219.
- Yildiz A, Sachs GS (2004) Characteristics of rapid cycling bipolar-I patients in a bipolar speciality clinic. *J Affect Disord* 79:247–251.
- Yildiz A, Vieta E, Leucht S, Baldessarini RJ (2010) Efficacy of anti-manic treatments: meta-analysis of randomized, controlled trials. *Neuropsychopharmacology* 36:375–389.
- Young AH, Rigney U, Shaw S, Emmas C, Thompson JM (2011) Annual cost of managing bipolar disorder to the UK health-care system. *J Affect Disord* 133:450–456.
- Yung AR, McGorry PD (1996) The prodromal phase of first-episode psychosis: past and current conceptualizations. *Schizophr Bull* 22:353–370.
- Yung AR, McGorry PD (2007) Prediction of psychosis: setting the stage. *Br J Psychiatry Suppl* 51:s1–8.