



## King's Research Portal

DOI:

[10.1080/08870446.2016.1275629](https://doi.org/10.1080/08870446.2016.1275629)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Moon, Z., Moss-Morris, R., Hunter, M. S., & Hughes, L. D. (2017). Measuring illness representations in breast cancer survivors (BCS) prescribed tamoxifen: Modification and validation of the Revised Illness Perceptions Questionnaire (IPQ-BCS). *Psychology & health*, 32(4), 439-458.  
<https://doi.org/10.1080/08870446.2016.1275629>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



**Measuring Illness Representations in Breast Cancer Survivors (BCS): Modification and validation of the Revised Illness Perceptions Questionnaire (IPQ-BCS)**

Journal:	<i>European Health Psychology Society</i>
Manuscript ID	GPSH-2016-0296.R1
Manuscript Type:	Psychology and Health
Keywords:	illness perceptions, scale validation, confirmatory factor analysis, IPQ-R, breast cancer, tamoxifen

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5 **Measuring Illness Representations in Breast Cancer Survivors (BCS): Modification**  
6 **and validation of the Revised Illness Perceptions Questionnaire (IPQ-BCS)**  
7  
8  
9

10 Zoe Moon, Rona Moss-Morris, Myra S. Hunter, Lyndsay D. Hughes

11  
12  
13 King's College London  
14  
15

16 Zoe Moon, Institute of Psychiatry, Psychology & Neuroscience, King's College London. 0207 188  
17 9324. [Zoe.moon@kcl.ac.uk](mailto:Zoe.moon@kcl.ac.uk)  
18  
19

20  
21 Rona Moss-Morris, Institute of Psychiatry, Psychology & Neuroscience, King's College London.  
22 0207 188 0178. [Rona.moss-morris@kcl.ac.uk](mailto:Rona.moss-morris@kcl.ac.uk)  
23  
24

25  
26 Myra Hunter, Institute of Psychiatry, Psychology & Neuroscience, King's College London. 0207 188  
27 0189. [Myra.hunter@kcl.ac.uk](mailto:Myra.hunter@kcl.ac.uk)  
28  
29

30  
31 Author note  
32  
33

34 Lyndsay Hughes, Institute of Psychiatry, Psychology & Neuroscience, King's College  
35 London.  
36  
37

38  
39 This research was supported by Breast Cancer Now under grant number  
40 2013NovPhD201.  
41  
42

43  
44 Correspondence concerning this article should be addressed to Dr Lyndsay Hughes,  
45 Health Psychology Section, King's College London, 5<sup>th</sup> Floor Bermondsey Wing, Guy's  
46 Hospital. London, SE1 9RT. Email; [Lyndsay.hughes@kcl.ac.uk](mailto:Lyndsay.hughes@kcl.ac.uk). Tel: 0207 188 9779.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **Measuring Illness Representations in Breast Cancer Survivors (BCS) prescribed**  
4 **tamoxifen: Modification and validation of the Revised Illness Perceptions**  
5 **Questionnaire (IPQ-BCS)**  
6

7 **Abstract**  
8

9  
10 Objective: The Revised Illness Perceptions Questionnaire (IPQ-R), widely used to assess  
11 illness perceptions, may fail to measure unique characteristics of different illnesses. This  
12 study modified and validated the IPQ-R for breast cancer survivors, to provide detailed  
13 understanding of the specific illness perceptions held by these patients.  
14  
15

16  
17 Design: Initial modifications were made following qualitative interviews and were revised in  
18 a think-aloud study. The modified scale was tested in 753 breast cancer survivors prescribed  
19 tamoxifen. Modifications included adding a tamoxifen consequences scale and adapting the  
20 timeline scales to measure beliefs around risk of recurrence and cure. A confirmatory factor  
21 analysis was conducted on the modified questionnaire and an exploratory factor analysis on  
22 the causal beliefs scale. Test-retest reliability, internal consistency and construct validity were  
23 also examined.  
24  
25

26  
27 Results: The proposed eight factor structure showed acceptable model fit, with high loadings  
28 and good reliability for all subscales. Correlations between subscales were consistent with  
29 theory and previous research.  
30  
31

32  
33 Conclusions: The IPQ-BCS is valid and reliable, and provides unique understanding of  
34 specific perceptions held by this population, including beliefs surrounding risk of recurrence  
35 and consequences of ongoing hormonal treatment. Identifying these perceptions will aid  
36 development of interventions targeting depression, fear of recurrence and medication non-  
37 adherence.  
38  
39

40  
41 **Keywords:** illness perceptions, scale validation, confirmatory factor analysis, IPQ-R, breast  
42 cancer, tamoxifen  
43  
44

45  
46  
47  
48  
49 **Introduction**  
50

51  
52 Illness representations or perceptions, which form part of the Common Sense Model of Self-  
53 Regulation (CSM; Leventhal, Diefenbach, & Leventhal, 1992), predict a range of outcomes,  
54 including quality of life (QOL) (Petrie, Jago, & Devcich, 2007), fatigue (Jopson & Moss-  
55 Morris, 2003) and poor physical and mental health (Frostholm et al., 2007; Whittaker, Kemp,  
56  
57  
58  
59  
60

1  
2  
3 & House, 2007). The CSM proposes that patients' coping behaviours, such as adherence, are  
4 guided by their cognitive and emotional representations of their illness. Cognitive  
5 representations include common sense beliefs about the illness identity (the symptoms / label  
6 associated with the illness), the cause(s), consequences, timeline and controllability of the  
7 illness. Patients also have emotional representations, such as fear, which guide how they  
8 respond to the illness. Finally, patients have a meta-cognitive perception of the coherence of  
9 their illness representations (Moss-Morris et al., 2002). The development of the Illness  
10 Perceptions Questionnaire (IPQ; Weinman, Petrie, Moss-Morris, & Horne, 1996), the Brief  
11 Illness Perceptions Questionnaire (Brief IPQ; Broadbent, Petrie, Main, & Weinman, 2006)  
12 and the Revised Illness Perceptions Questionnaire (IPQ-R; Moss-Morris et al., 2002) allowed  
13 researchers to quantify illness representations and increased empirical research on the role of  
14 illness perceptions in areas such as coping, medication adherence and health outcomes.  
15  
16  
17  
18  
19  
20  
21  
22

23  
24 The IPQ-R has shown good internal reliability and test-retest reliability, as well as sound  
25 discriminant, known group and predictive validity (Moss-Morris et al., 2002). However, it  
26 was developed as a generic scale for use across different illness groups and therefore may not  
27 provide insight into the unique beliefs of different patient groups (French & Weinman, 2008).  
28 Whilst the authors of the IPQ-R recommend that the scale is modified for use in different  
29 contexts (Moss-Morris et al., 2002), validated modified versions are currently lacking.  
30 Researchers often rely on very minor modifications such as adding symptoms or causes  
31 which may not tap into illness specific beliefs. Think-aloud studies have shown that patients  
32 can struggle to answer questions on the IPQ. Patients enrolled in physiotherapy or a  
33 preoperative exercise programme had some difficulty completing the Brief IPQ and  
34 occasionally misinterpreted questions (van Oort, Schroder, & French, 2011). Another study  
35 showed that patients with type 2 diabetes had difficulties with the concepts of cure and  
36 symptoms and misunderstood the negative wording on some questions on the IPQ-R  
37 (McCorry, Scullion, McMurray, Houghton, & Dempster, 2013). This highlights the need to  
38 explore the face validity of IPQ-R items in different groups of patients and to test the face  
39 validity of modifications using think-aloud methods.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 One patient group for whom modifications may be particularly pertinent are breast cancer  
52 survivors (BCS). There are around three million BCS living in the US and another 200,000  
53 women are diagnosed with breast cancer every year (American Cancer Society, 2014). These  
54 patients have completed their active treatment and may no longer consider themselves to be  
55 ill, although continued therapy and monitoring is required. They may therefore struggle to  
56  
57  
58  
59  
60

1  
2  
3 answer questions on the IPQ-R about current illness or current symptom control. However,  
4 BCS experience a myriad of psychosocial issues and measuring illness perceptions is relevant  
5 to understanding these ongoing reactions to their previous cancer. For example, around a  
6 quarter of BCS experience depression or fatigue, and up to 70% show clinical levels of fear  
7 of cancer recurrence (FCR) (Cvetković & Nenadović, 2016; Servaes, Gielissen, Verhagen, &  
8 Bleijenberg, 2007; Thewes et al., 2012). Others also struggle to cope with long term  
9 hormonal therapy such as tamoxifen, which is prescribed for up to ten years as adjuvant  
10 treatment for women with oestrogen receptive positive breast cancer (about 75% of all breast  
11 cancers; Harrell et al., 2007). Whilst tamoxifen is one of the most effective systemic  
12 treatments available for breast cancer, it can cause unpleasant side effects (Garreau,  
13 Delamelena, Walts, Karamlou, & Johnson, 2006) and both non-adherence and non-  
14 persistence rates are often as high as 50% within five years of treatment (Hershman et al.,  
15 2010; Kostev, Haas, & Hadji, 2012; Owusu et al., 2008). Non-adherence to tamoxifen is  
16 associated with significantly increased risk of recurrence and mortality (Barron, Cahir, Sharp,  
17 & Bennett, 2013; Hershman et al., 2011; Makubate, Donnan, Dewar, Thompson, &  
18 McCowan, 2013). However, little is known about how illness perceptions and beliefs may  
19 affect adherence in this population.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31  
32 An IPQ modified to address beliefs about a past illness, possibility of recurrence and ongoing  
33 adjuvant treatment will allow researchers to investigate illness representations alongside BCS  
34 specific coping (including adherence) and psychological outcomes. The CSM has been  
35 suggested as a useful framework for understanding FCR (Fardell et al., 2016) and other breast  
36 cancer survivorship issues (Kaptein et al., 2015). Further, identifying illness perceptions  
37 idiosyncratic to BCS could aid development of interventions, which have the potential to  
38 improve psychological wellbeing and QOL (Simard et al., 2013).  
39  
40  
41  
42  
43  
44

45 This study aimed to modify the IPQ-R for use with BCS. We focused specifically on women  
46 taking tamoxifen in order to get a more homogenous sample and to tap into illness beliefs  
47 specific to adjuvant therapy. Following advice from French and Weinman (2008), we used a  
48 mixed methods approach to modify and validate the questionnaire.  
49  
50  
51

52 The specific objectives were:

- 53  
54 1. To conduct qualitative interviews based on the CSM to elicit key beliefs held by BCS  
55 taking tamoxifen;
- 56  
57 2. To use these interviews to develop a modified version of the IPQ-R (the IPQ-BCS);  
58  
59  
60

3. To test the face validity of this modified questionnaire using think aloud interviews and modify further if indicated;
4. To assess the factor structure, internal consistency and test-retest reliability of the modified IPQ-BCS in a large cross sectional study of BCS;
5. To assess construct validity of the new subscales using inter-correlations between subscales and relationships between subscales and psychological variables (beliefs about medications and distress).

It was hypothesised that IPQ-R subscales would show correlations similar to that found in previous research (Hagger & Orbell, 2005; Moss-Morris et al., 2002). We hypothesised that distress would be associated with higher consequences, identity, emotional representations and risk of recurrence beliefs; that tamoxifen necessity beliefs would correlate with treatment control; and that tamoxifen concerns would correlate consequences and identity beliefs.

## **Method**

The study was approved by the Northampton National Research Ethics Committee (REF 14/EM/1207).

### ***Qualitative study***

#### *Participants and procedure*

Participants were recruited through an oncology clinic in a London hospital and through online advertisements, as part of a larger study investigating women's experiences of taking tamoxifen. Patients were eligible if they were female, over the age of 18 and had been prescribed tamoxifen post primary breast cancer. Patients were told about the research by their clinician, and if interested they were introduced to a researcher and given an information sheet. Women who responded to online advertisements were screened for eligibility and given information about the study.

A follow up call was made two days later to arrange an interview. This was part of a larger qualitative study to explore women's experiences of taking tamoxifen. Patients were interviewed face to face or over the telephone. Informed consent was taken prior to each interview. Interviews were recorded and transcribed verbatim. Participants were first asked a series of general questions about their experience of tamoxifen, before being asked specific questions regarding their illness perceptions for modification of the IPQ-R (See Table 1 for

1  
2  
3 interview schedule). Thirty-two women took part in the larger qualitative study, of whom 18  
4 were asked the additional questions specifically relating to the modification of the  
5 questionnaire. Data collection for these additional questions ceased once data saturation was  
6 reached and only these questions were analysed in this study. Thus, data from 18 women  
7 were analysed. Participant demographics are shown in Table 2.  
8  
9

10  
11 [TABLE 1 NEAR HERE]  
12  
13

#### 14 *Item development*

15  
16  
17 Interviews were analysed using deductive analysis. Using the CSM as a framework, themes  
18 were generated around prevalent beliefs and perceptions. Changes to the questionnaire were  
19 made to reflect the language used by participants. A key theme was that women did not  
20 identify as currently having breast cancer. All questions were amended to avoid asking  
21 women about their breast cancer in the present tense. Original and amended items are shown  
22 in Supplementary Material.  
23  
24  
25  
26

27  
28 A second theme suggested women attributed few symptoms to breast cancer. Therefore the  
29 identity scale was amended to assess symptoms which were (a) attributed to breast cancer, (b)  
30 to tamoxifen treatment and (c) to their previous / other treatment. Analysis of the interviews  
31 elicited specific tamoxifen related symptoms. Ten new symptoms, such as hot flushes and  
32 change in sex drive were added to the original list of fourteen symptoms in the core version  
33 of the IPQ-R (See Table 3 for list of additional symptoms).  
34  
35  
36  
37

38  
39 When asked about control, consequences and causes women tended to discuss their risk of  
40 recurrence instead of their breast cancer. Therefore, to effectively assess control beliefs, the  
41 personal and treatment control subscales were amended so that the word '*illness*' was  
42 replaced with '*risk of recurrence*'. The treatment control items were asked specifically in  
43 relation to tamoxifen. In addition to the existing illness consequences scale, a new scale was  
44 added to assess the consequences of taking tamoxifen, as this was a dominant theme  
45 identified in the interviews. With regards to timeline beliefs, the interviews showed that  
46 women did not have symptoms which come and go. The cyclic timeline scale was removed  
47 and a new scale was added to assess risk of recurrence. Likewise, the timeline acute / chronic  
48 scale was amended to assess the extent to which women believe that their breast cancer is  
49 cured, as the interviews showed that these beliefs were much more pertinent than beliefs  
50 around the chronic nature of breast cancer itself. The coherence scale was modified to  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 measure understanding of tamoxifen treatment rather than breast cancer. Finally, as women  
4 discussed fear around risk of recurrence rather than fear around breast cancer, the emotional  
5 representations scale was amended to reflect this. The cause scale was modified by adding  
6 breast cancer specific causes such as hormonal influence and removing causes which were  
7 not applicable. Examples of changes to specific items are shown in Table 3.  
8  
9

### 10 11 *Think-aloud study*

12  
13  
14 After item modification, a think-aloud study was conducted to examine if items on the new  
15 IPQ-R were being understood and interpreted in the expected way. Eleven women from the  
16 interview study were invited to take part in the think aloud study and eight agreed. Think  
17 aloud studies involve patients verbalising their thought process as they answer the  
18 questionnaire (Ericsson & Simon, 1998). These methods have been used previously to  
19 examine questionnaires assessing illness perceptions (van Oort et al., 2011), theory of  
20 planned behaviour (French, Cooke, McLean, Williams, & Sutton, 2007) and QOL  
21 (Westerman et al., 2008). Participants were asked to complete the modified IPQ-BCS and to  
22 verbalise everything they were thinking as they were completing the questionnaire. If they  
23 were quiet for a long period of time, they were prompted to think aloud as they were  
24 considering the question. The think-aloud sessions were conducted over the telephone and  
25 participants consented to be audio recorded.  
26  
27  
28  
29  
30  
31  
32  
33

34  
35 [TABLE 2 NEAR HERE]  
36

37  
38 The think-aloud interviews showed that women could understand the questionnaire and that  
39 they found it relevant and applicable. However, several issues were identified which led to  
40 further modifications. The instructions to both the identity and cause scales were modified to  
41 improve their clarity. A few participants remarked that some items in the personal and  
42 treatment control scales were worded too severely and that they were unsure how to answer  
43 them. Therefore the items were amended to reflect this. Several other items were revised  
44 slightly to enhance the chance they would be applicable for all participants or to ensure they  
45 were being correctly interpreted. Some women remarked on the repetitiveness of questions,  
46 so where possible items were deleted (See Supplementary Material).  
47  
48  
49  
50  
51  
52

53 [TABLE 3 NEAR HERE]  
54  
55  
56  
57  
58  
59  
60

## *Quantitative study*

### *Participants and procedure*

Participants were recruited through oncology clinics at 25 NHS Trusts throughout England and through online advertisements. Patients were eligible if they had a diagnosis of primary breast cancer and if they had been prescribed tamoxifen. Participants had to be female and over the age of 18. Patients were approached by a member of their clinical team during a routine clinic appointment or received an invitation in the post from their clinical team. They were given information about the study along with the questionnaire and a return envelope. After providing informed consent, participants either completed the questionnaire in the clinic or took it home to return to the researcher. Participants who were recruited online responded to an advert and after being screened for eligibility, were sent information about the study along with a link to an online questionnaire. Participants gave informed consent whilst completing the online questionnaire. A separate sample was recruited to assess the test-retest reliability of the IPQ-R. This sample was recruited from four NHS Trusts. Participants were given information about the study from the clinical team and once consented, they completed the first questionnaire in clinic. Participants were either posted the second questionnaire or given a link to complete it online two weeks later, whichever was their preference. Telephone reminders were made if the second questionnaire had not been returned within one week.

### *Measures*

*Modified IPQ-R (IPQ-BCS)*. Participants completed the modified version of the IPQ-R (IPQ-BCS), which included subscales measuring identity, cure beliefs, risk of recurrence, tamoxifen consequences, breast cancer consequences, personal control over recurrence, tamoxifen control, coherence, emotional representations and causes. All questions were scored on a five point Likert type scale ranging from *Strongly Agree* to *Strongly Disagree* with the exception of the identity scale where participants ticked each column to indicate if they experienced that symptom. Each subscale included four items, with the exception of cure beliefs, tamoxifen consequences and emotional representations, which included five items. The identity subscale was calculated by totalling the number of symptoms which were attributed to tamoxifen. Symptoms which were added to the original list of symptoms are shown in Table 3.

1  
2  
3 *Hospital Anxiety and Depression Scale (HADS)*. The HADS is a 14 item scale measuring  
4 depression and anxiety (Zigmond & Snaith, 1983). The total distress scale was used in this  
5 study, as a large meta confirmatory factor analysis has shown evidence of a strong general  
6 HADS factor rather than two distinct subscales (Norton, Cosco, Doyle, Done, & Sacker,  
7 2013). Each item is scored on a scale of 0 - 3, with higher scores reflecting higher levels of  
8 distress. The HADS has shown good reliability in patients with breast cancer (Matthews et  
9 al., 2014; Stanton, Petrie, & Partridge, 2014).

10  
11  
12  
13  
14  
15 *Beliefs about Medicines Questionnaire (BMQ)*. The BMQ-Specific measures beliefs  
16 surrounding the necessity of taking medications and concerns about adverse effects (Horne,  
17 Weinman, & Hankins, 1999). The word medication was replaced with the word tamoxifen for  
18 all items. Each item is rated on a five point Likert type scale. A higher score on each subscale  
19 indicates stronger necessity or concern beliefs. The scale has been used many times in BCS  
20 with Cronbach's alpha values of 0.79 - 0.86 and 0.72 - 0.84 for the necessity and concerns  
21 scale respectively (Bender et al., 2014; Corter, Findlay, Broom, Porter, & Petrie, 2013; Jacob  
22 Arriola et al., 2014).

### 23 24 25 26 27 28 29 30 *Statistical analysis*

31  
32 Missing data was less than 5% and were replaced using mean substitution. A Confirmatory  
33 Factor Analysis (CFA) was conducted on the modified IPQ-BCS using Mplus version 7 to  
34 test the hypothesised model of eight subscales (cure beliefs, tamoxifen consequences, risk of  
35 recurrence, breast cancer consequences, personal control, treatment control, coherence and  
36 emotional representations). CFA is the gold standard method for evaluation of construct  
37 validity in psychometric tests (Hu & Bentler, 1999). The CFA was conducted using weighted  
38 least squares with means and variances corrected (WLSMV), as the data was measured on an  
39 ordinal categorical scale. Multiple indices were used to assess model fit. Chi-squared was not  
40 used as it is sensitive to sample size (Byrne, 2001). The Comparative Fit Index (CFI), Root  
41 Mean Square Error of Approximation (RMSEA) and Tucker Lewis Index (TLI) were used.  
42 CFI or TLI values of greater than 0.95 suggest acceptable model fit (Hu & Bentler, 1999).  
43 RMSEA values of 0.08 indicate reasonable fit and values of under 0.06 indicate good fit (Hu  
44 & Bentler, 1999). The reliability of each subscale was tested using Cronbach's alpha. Test-  
45 retest reliability was assessed using intra-class correlation of each subscale at baseline and  
46 two week follow up. Discriminant validity was assessed using inter-correlations between  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 IPQ-R dimensions. Construct validity was assessed by examining the correlations between  
4 IPQ-R dimensions and additional variables (beliefs about medications and distress).  
5  
6

7 It is recommended that the causal attribution scale be examined in an exploratory fashion  
8 (Dempster & McCorry, 2012); therefore Exploratory Factor Analysis (EFA) was used as it  
9 does not specify an underlying factor structure. Item frequencies were visually inspected and  
10 items were removed if the majority of participants did not see them as a cause. An EFA was  
11 then conducted using the SPSS R-menu for ordinal factor analysis based on polychoric  
12 correlations (Basto & Pereira, 2012). The number of factors to retain was assessed using  
13 parallel analysis (Horn, 1965). The factor analysis was conducted using Maximum  
14 Likelihood extraction and Geomin Q-Q rotation.  
15  
16  
17  
18  
19

## 20 21 **Results**

22  
23 Data was collected from 753 participants. Participants were all female and had been  
24 diagnosed with Stage I – III breast cancer (Table 2). Mean age was 53 (SD=10.5) and  
25 participants were on average 33 months post breast cancer diagnosis (SD=24, range 2 months  
26 – 16 years).  
27  
28  
29

### 30 31 *Confirmatory Factor Analysis*

32  
33 The sample size exceeded the requirements of at least 3 cases per item (Tabachnick & Fidell,  
34 2007). Visual inspection of the data showed the items generally correlated as expected within  
35 the eight subscales, indicating that a CFA was appropriate. The 35 item IPQ-BCS showed  
36 acceptable model fit (RMSEA = 0.08, 95% CI= 0.08 – 0.09, CFI = 0.95, TLI = 0.94). In  
37 order to reduce the length of the scale, one item (with the lowest factor loading) was removed  
38 from each of the three subscales with five items (tamoxifen consequences, cure beliefs and  
39 emotional representations). Removing these items did not change the overall model fit, and  
40 therefore this briefer questionnaire is preferred where all subscales have four items. Table 4  
41 shows the factor loadings for each of the items under each of the subscales. Factor loadings  
42 were all well above the required threshold of 0.40 (Ford, MacCallum, & Tait, 1986), ranging  
43 from 0.63 to 0.95.  
44  
45  
46  
47  
48  
49  
50  
51

52 [TABLE 4 NEAR HERE]  
53  
54  
55  
56  
57  
58  
59  
60

### ***Internal and test-retest reliability***

All scales showed excellent reliability, with Cronbach's alpha values ranging from .76 to .92 (Table 4). Test-retest reliability was tested in a separate sample of 48 women. Participants completed the questionnaire twice; on average 18 days apart (range 11 – 31). The intraclass correlation coefficients for each scale ranged from 0.77 to 0.94 indicating excellent test-retest reliability (Table 4).

### ***Exploratory Factor Analysis on cause items***

Item frequencies and correlations were explored visually and two items were removed from the EFA. Item 3 (*A germ or virus*) was removed as it did not correlate with other items and only 5% of participants agreed that it might be a risk factor for recurrence. Item 12 (*smoking*) was also removed, as only 24% of participants provided data for this question. *Hormonal influence* was the strongest item, with 81% of participants agreeing that it was a risk factor.

Parallel analysis was used on eleven causal items to assess the number of factors to retain, and indicated a three factor solution, explaining 46% of the total variance. Factor loadings are shown in Table 5. The first factor, labelled psychological attributions, included items relating to stress, worries and emotional state. The second factor, labelled health behaviours, included items such as diet and eating habits, and exercise. These two factors showed good reliability (.85 and .72 respectively). The final factor included item 11 (*ageing*) and item 13 (*hormonal influence*). However, hormonal influence had a factor loading of below .4 and the reliability of the factor was very low (.44). Therefore, these items might be best considered individually and not as part of a sub-scale. Item 2 (*runs in the family*) and item 5 (*chance or bad luck*) did not load onto any factors.

[TABLE 5 NEAR HERE]

### ***Examination of the Identity scale***

Each symptom was experienced by at least 13% of participants. Over 40% of participants had experienced pain, weight loss/gain, hot flushes, night sweats, fatigue, sleep difficulties, joint pain and loss of sex drive. Patients experienced on average 7.8 symptoms (SD = 5.9).

Symptoms were more commonly attributed to tamoxifen (mean = 5.8, SD = 4.9), than to breast cancer (mean=2.1, SD = 3.2) or previous/other treatment (mean = 2.0, SD = 3.6). As symptoms were rarely attributed to breast cancer, identity was represented by the total

1  
2  
3 number of symptoms attributed to tamoxifen. All symptoms were most commonly attributed  
4 to tamoxifen, with the exception of pain which was attributed to breast cancer by 29% of  
5 participants and to tamoxifen by 14% of participants. Hot flushes were the most common  
6 symptom attributed to tamoxifen (65%), followed by night sweats (55%), weight loss / gain  
7 (41%), joint pain (37%), fatigue (35%), leg cramps (35%) and vaginal dryness, itchiness or  
8 discomfort (35%). These results provide support for the validity of the symptoms included in  
9 the scale as well as the different sources of attribution.  
10  
11  
12  
13

### 14 *Construct validity*

15  
16  
17 Inter-correlations between the IPQ-BCS subscales are shown in Table 6. The direction and  
18 size of the correlations are consistent with previous research (Hagger & Orbell, 2005; Moss-  
19 Morris et al., 2002), and with what would be expected due to the underlying theory.  
20  
21 Tamoxifen consequences and breast cancer consequences were positively correlated. Both  
22 consequences scales correlated positively with emotional representations and risk of  
23 recurrence and negatively with cure beliefs, treatment control. Cure beliefs had a moderate  
24 negative correlation with risk of recurrence. Personal control and treatment control were  
25 strongly correlated. Both control scales correlated positively with coherence and cure beliefs  
26 and negatively with risk of recurrence. Emotional representations was negatively correlated  
27 with cure beliefs, treatment control and positively correlated with risk of recurrence. Identity  
28 beliefs correlated positively with tamoxifen consequences, risk of recurrence, breast cancer  
29 consequences and emotional representations.  
30  
31  
32  
33  
34  
35  
36  
37

38 [TABLE 6 NEAR HERE]

39  
40  
41 To further explore the validity of the constructs of the IPQ-BCS subscales, correlations were  
42 examined with distress using the HADS and treatment beliefs using the BMQ. These  
43 correlations were consistent with hypothesised relationships and supported the construct  
44 validity of the IPQ-R dimensions (Table 7). HADs distress correlated positively with identity,  
45 consequences, risk of recurrence and emotional representations, and negatively with cure  
46 beliefs and treatment control. BMQ Tamoxifen concerns correlated positively with IPQ-BCS  
47 tamoxifen consequences, breast cancer consequences, identity and emotional representations,  
48 and negatively with treatment control and coherence. BMQ Tamoxifen necessity beliefs  
49 correlated positively with IPQ-BCS emotional representations and treatment control.  
50  
51  
52  
53  
54  
55

56 [TABLE 7 NEAR HERE]

## Discussion

This paper developed and validated a modified version of the IPQ-R for use with BCS prescribed tamoxifen. The modified version includes an identity scale which has been modified to assess symptoms attributed to tamoxifen, the original illness consequences scale and a new tamoxifen consequences scale. The timeline acute scale was amended to measure cure beliefs and the timeline cyclical was replaced with a risk of recurrence scale. The personal control, treatment control and emotional representations scales were amended to assess risk of recurrence rather than current cancer. The coherence scale was amended to measure coherence around tamoxifen rather than breast cancer. The 35 item IPQ-BCS showed acceptable model fit, with high factor loadings on the conceptual subscales, and high reliability for all subscales. To decrease participant burden, this was reduced down to a 32 item questionnaire where each subscale has four items. This modification did not affect model fit and the reliability for each scale remained high, demonstrating that the removed items were redundant and that the shortened questionnaire is sufficient to understand these constructs. This modification and validation of the IPQ-R for use in BCS was a vital step in furthering understanding of illness perceptions held by BCS. The qualitative interviews we conducted showed that women would have had difficulty answering questions on the original IPQ-R regarding their current illness and breast cancer symptoms. The think-aloud study showed that items on the modified IPQ-BCS were easy to interpret and to answer.

These results provide support for the CSM and the idea that BCS hold perceptions about their previous breast cancer and ongoing treatment and survivorship. Investigating these illness perceptions will enhance understanding of the psychosocial issues associated with breast cancer survivorship and will help with developing interventions to reduce distress or improve QOL in this population. The modified IPQ-BCS assesses beliefs which would not have been assessed with the original IPQ-R, such as beliefs around risk of recurrence and cure. These beliefs are likely to be relevant to understanding FCR and depression in BCS. The benefit of using the IPQ-BCS to assess FCR is that it allows examination of both perceptions of risk (risk of recurrence scale) and emotional responses to this risk perception (emotional representations scale). Whilst they are correlated, perceptions of the likelihood of a recurrence differ from the emotions (e.g. fear, distress) that women feel in response to this risk perception. Understanding these separate constructs and how they relate to distress or QOL will aid development of interventions to reduce FCR. Furthermore, the IPQ-BCS allows these risk of recurrence beliefs to be measured alongside other illness perceptions, such as

1  
2  
3 control or consequences, which feed into beliefs around risk of recurrence (Fardell et al.,  
4 2016). The IPQ-BCS could be supplemented with a more complex FCR scale which also  
5 assesses hypervigilant checking behaviours, functional impairment of FCR or FCR in relation  
6 to actual risk.  
7  
8

9  
10 The IPQ-BCS also measures beliefs regarding tamoxifen treatment specifically, rather than  
11 the more generalised treatment control scale included in the IPQ-R. The IPQ-BCS assesses  
12 consequences of ongoing tamoxifen treatment as well as breast cancer consequences, and  
13 measures treatment control specifically with regards to tamoxifen treatment. This scale could  
14 therefore be used to identify illness and treatment beliefs related to non-adherence in this  
15 population. Previous studies have found problems with the treatment control subscale of the  
16 IPQ-R, such as low reliability and cross-loading of items (Brzoska, Yilmaz-Aslan,  
17 Sultanoglu, Sultanoglu, & Razum, 2012; Ibrahim, Desa, & Chiew-Tong, 2011; Moss-Morris  
18 et al., 2002). This is likely due to participants being unsure as to which treatment the  
19 questions are referring to. Amending this subscale to specifically assess tamoxifen treatment  
20 may have overcome these issues, as the IPQ-BCS treatment control subscale showed good  
21 reliability and was free from cross-loading. This scale could also be amended to assess  
22 treatment control specific to aromatase inhibitors or hormone therapy in general.  
23  
24  
25  
26  
27  
28  
29  
30  
31

32  
33 The EFA on the cause scale produced three factors. Factor one (psychological attributions)  
34 and factor two (health behaviours) showed good reliability. However some items did not load  
35 onto any factors or had low factor loadings. These results are not consistent with the original  
36 IPQ-R factor structure (Moss-Morris et al., 2002). However, several papers have found a  
37 factor structure which is hard to interpret (Nicholls, Hill, & Foster, 2013; Wittkowski,  
38 Richards, Williams, & Main, 2008). In a sample of Greek cancer patients, Giannousi et al.  
39 (2010) also found that items 2 (hereditary), 5 (chance or bad luck) and 11 (ageing) did not  
40 load onto any factors. Whilst hormonal influence and chance or bad luck did not load onto  
41 any factors in this analysis, they were the most consistently endorsed causes and therefore  
42 they should be considered as individual items in future analysis or larger subscales related to  
43 these constructs should be developed. Whilst attempts were made to amend the cause scale  
44 to enhance its applicability, further modifications may be needed to develop a more robust  
45 factor structure.  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 Correlations between IPQ-BCS subscales were consistent with theory and previous research  
56 and showed good construct validity. The original consequences scale correlated positively  
57  
58  
59  
60

1  
2  
3 with the new tamoxifen consequences subscale, but the correlation was only moderate, which  
4 supports the idea that patients can differentiate symptoms from their breast cancer and their  
5 tamoxifen treatment. Previous research in a cancer setting has found overlap of the  
6 consequences and emotional representations scales, where items from both subscales loaded  
7 onto the same factor (Giannousi et al., 2010). However, the IPQ-BCS correlations between  
8 these subscales were only moderate and the hypothesised factor structure was supported,  
9 suggesting that emotional representations around recurrence are distinct from consequences  
10 of breast cancer. The risk of recurrence scale, which was adapted from the previous timeline  
11 cyclical scale, showed that having high beliefs of a recurrence was associated with higher  
12 consequences, higher emotional representations and lower cure beliefs.  
13  
14  
15  
16  
17  
18  
19

20 The personal and treatment control subscales were positively correlated, but the correlations  
21 were low enough to support the assumption of two distinct constructs, which is consistent  
22 with previous research (Dempster & McCorry, 2012; Giannousi et al., 2010; Moss-Morris et  
23 al., 2002). Women who scored highly on the two control subscales were less likely to believe  
24 they would have a recurrence, more likely to believe their breast cancer had been cured and  
25 more likely to have higher coherence beliefs. Women who attributed a high number of  
26 symptoms to tamoxifen were significantly more likely to believe they would have a risk of  
27 recurrence and less likely to believe they were cured, but these were small correlations. This  
28 is consistent with correlations found in previous research (Hagger & Orbell, 2003) and  
29 suggests that there is a relationship between symptom experience and perceptions of risk.  
30  
31  
32  
33  
34  
35  
36  
37

38 The correlations between IPQ-BCS subscales, HADS distress and BMQ treatment beliefs  
39 provided further support for construct validity. Higher concerns about tamoxifen were  
40 associated with higher tamoxifen consequences, a greater number of symptoms attributed to  
41 tamoxifen, and to a lesser extent, higher breast cancer consequences. This is expected in this  
42 population as tamoxifen concerns focus almost exclusively on side effects (Moon, Moss-  
43 Morris, Hunter, & Hughes, 2016) and are therefore related to beliefs around consequences  
44 and symptom attribution. Understanding the interactions between illness perceptions and  
45 medication beliefs may help to understand medication non-adherence in BCS (Horne &  
46 Weinman, 2002). HADS distress was associated with tamoxifen consequences, breast cancer  
47 consequences and emotional representations. These relationships make theoretical sense, as  
48 greater illness consequences are likely to contribute to levels of distress. However, as this was  
49 cross-sectional data, the direction of the effect cannot be established. It may be that women  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 who experience higher levels of distress perceive greater consequences from their illness or  
4 ongoing treatment.  
5

6  
7 Strengths of this study include the large sample size and robust analysis. The scale was  
8 amended based on interviews with patients, and before being analysed it was subject to think-  
9 aloud analysis. Furthermore, patients were recruited from hospitals throughout England,  
10 which should enhance the generalisability of the results. However, there were several  
11 limitations. Firstly, participants from the same sample were used to develop the items on the  
12 questionnaire and to test the questionnaire in the think aloud studies. Secondly, the factor  
13 structure has only been tested and validated in one sample. Future research could test whether  
14 the IPQ-BCS could be modified further for use in different cancer types with similar  
15 survivorship issues to BCS. Overall, results suggest that the modified IPQ-BCS is a valid and  
16 reliable measure. It is well understood in BCS and has a clear factor structure with ten  
17 distinct constructs (cause, identity, cure, tamoxifen consequences, risk of recurrence, breast  
18 cancer consequences, personal control, treatment control, coherence, emotional  
19 representations). Utilising this scale will help us to understand how women feel about their  
20 illness and their ongoing treatment as they move into survivorship. Illness perceptions have  
21 been shown to be relevant to many of the psychosocial issues inherent to BCS, such as  
22 fatigue, non-adherence, distress and FCR. Using the IPQ-BCS will allow researchers to see  
23 how dimensions such as emotional representations and sense of coherence affect illness  
24 behaviours such as adherence, or outcomes such as QOL or survival, and will help generate  
25 interventions to support these patients. Whilst the scale was developed for tamoxifen  
26 treatment, it is likely it will be equally as applicable for women who have been prescribed  
27 other hormonal therapy such as aromatase inhibitors. It can also be used in other areas, such  
28 as to investigate beliefs around cancer in relation to FCR, fatigue or distress.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 Word Count: 5282

45  
46  
47 Disclosure: No potential conflict of interest was reported by the authors.  
48

49 Acknowledgements: The research was funded by Breast Cancer Now. We are grateful to all  
50 the women who took part in the study. We would also like to thank all the sites who recruited  
51 to the study (Airedale General Hospital, Calderdale and Huddersfield NHS Foundation Trust,  
52 Charing Cross Hospital, Dorset County Hospital, Ealing Hospital, East Lancashire Hospitals  
53 NHS Trust, East Sussex Healthcare NHS Trust, Frimley Park Hospital, Great Western  
54 Hospital, Guy's Hospital, Kingston Hospital, Macclesfield Hospital, Norfolk and Norwich  
55  
56  
57  
58  
59  
60

1  
2  
3 University Hospital, Northwick Park Hospital, Oxford University Hospitals NHS Foundation  
4 Trust, Queen Alexandra Hospital, Royal Albert Edward Infirmary, Royal Blackburn Hospital,  
5 Royal Shrewsbury Hospital, Russells Hall Hospital, Tameside Hospital, The Mid Yorkshire  
6 Hospitals NHS Trust, The Pennine Acute Hospitals NHS Trust, The Princess Alexandra  
7 Hospital, The Whittington Hospital, University Hospital Coventry and Warwickshire, York  
8 Teaching Hospital NHS Foundation Trust) and Breast Cancer Care and Macmillan Cancer  
9 Support for assisting us with online advertisements.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 References:

- 24  
25 American Cancer Society. (2014). Cancer Treatment and Survivorship Facts & Figures 2014-  
26 2015. Atlanta: American Cancer Society.  
27  
28 Barron, T. I., Cahir, C., Sharp, L., & Bennett, K. (2013). A nested case-control study of  
29 adjuvant hormonal therapy persistence and compliance, and early breast cancer  
30 recurrence in women with stage I-III breast cancer. *British Journal of Cancer*, *109*,  
31 1513-1521. doi: 10.1038/bjc.2013.518  
32  
33 Basto, M., & Pereira, J. (2012). An SPSS R-menu for ordinal factor analysis. *Journal of*  
34 *Statistical Software*, *46*, 1-29.  
35  
36 Bender, C. M., Gentry, A. L., Brufsky, A. M., Casillo, F. E., Cohen, S. M., Dailey, M. M., . .  
37 . Sereika, S. M. (2014). Influence of patient and treatment factors on adherence to  
38 adjuvant endocrine therapy in breast cancer. *Oncology Nursing Forum*, *41*, 274-285.  
39 doi: 10.1188/14.ONF.274-285  
40  
41 Broadbent, E., Petrie, K.J., Main, J., & Weinman, J.(2006). The Brief Illness Perception  
42 Questionnaire. *Journal of Psychosomatic Research*, *60*, 631-637. doi:  
43 <http://dx.doi.org/10.1016/j.jpsychores.2005.10.020>  
44  
45 Brzoska, P., Yilmaz-Aslan, Y., Sultanoglu, E., Sultanoglu, B., & Razum, O. (2012). The  
46 factor structure of the Turkish version of the Revised Illness Perception Questionnaire  
47 (IPQ-R) in patients with diabetes and cardiovascular disease. *BMC Public Health*, *12*,  
48 1. doi: 10.1186/1471-2458-12-852  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Byrne, B.M. (2001). Structural equation modeling with AMOS, EQS, and LISREL:  
4 Comparative approaches to testing for the factorial validity of a measuring instrument.  
5 *International Journal of Testing, 1*, 55-86. doi: 10.1207/S15327574IJT0101  
6  
7  
8 Corter, A. L., Findlay, M., Broom, R., Porter, D., & Petrie, K. J. (2013). Beliefs about  
9 medicine and illness are associated with fear of cancer recurrence in women taking  
10 adjuvant endocrine therapy for breast cancer. *British Journal of Health Psychology,*  
11 *18*, 168-181. doi: 10.1111/bjhp.12003  
12  
13 10.1080/08870440701616714  
14  
15  
16 Cvetković, J., & Nenadović, M. (2016). Depression in breast cancer patients. *Psychiatry*  
17 *Research, 240*, 343-347. doi: <http://dx.doi.org/10.1016/j.psychres.2016.04.048>  
18  
19  
20 Dempster, M., & McCorry, N. K. (2012). The factor structure of the revised Illness  
21 Perception Questionnaire in a population of oesophageal cancer survivors.  
22 *Psychooncology, 21*, 524-530. doi: 10.1002/pon.1927  
23  
24  
25 Ericsson, K.A., & Simon, H.A. (1998). How to study thinking in everyday life: Contrasting  
26 think-aloud protocols with descriptions and explanations of thinking. *Mind, Culture,*  
27 *and Activity, 5*, 178-186. doi: 10.1207/s15327884mca0503\_3  
28  
29  
30 Fardell, J. E., Thewes, B., Turner, J., Gilchrist, J., Sharpe, L., Smith, A., . . . Butow, P.  
31 (2016). Fear of cancer recurrence: a theoretical review and novel cognitive processing  
32 formulation. *Journal of Cancer Survivorship*. Advance online publication. doi:  
33 10.1007/s11764-015-0512-5  
34  
35  
36 Ford, J. K., MacCallum, R. C., & Tait, M. (1986). THE APPLICATION OF  
37 EXPLORATORY FACTOR ANALYSIS IN APPLIED PSYCHOLOGY: A  
38 CRITICAL REVIEW AND ANALYSIS. *Personnel Psychology, 39*, 291-314. doi:  
39 10.1111/j.1744-6570.1986.tb00583.x  
40  
41  
42 French, D. P., Cooke, R., McLean, N., Williams, M., & Sutton, S. (2007). What do people  
43 think about when they answer theory of planned behaviour questionnaires? A 'think  
44 aloud' study. *Journal of Health Psychology, 12*, 672-687. doi:  
45 10.1177/1359105307078174  
46  
47  
48 French, D.P., & Weinman, J. (2008). Current issues and new directions in Psychology and  
49 Health : "Assessing illness perceptions: Beyond the IPQ". *Psychology & Health, 23*,  
50 5-9. doi: 10.1080/08870440701616714.  
51  
52  
53  
54 Frostholm, L., Oernboel, E., Christensen, K.S., Toft, T., Olesen, F., Weinman, J., & Fink, P.  
55 (2007). Do illness perceptions predict health outcomes in primary care patients? A 2-  
56  
57  
58  
59  
60

- 1  
2  
3 year follow-up study. *Journal of Psychosomatic Research*, 62, 129-138. doi:  
4 10.1016/j.jpsychores.2006.09.003  
5  
6 Garreau, J. R., Delameleno, T., Walts, D., Karamlou, K., & Johnson, N. (2006). Side effects  
7 of aromatase inhibitors versus tamoxifen: the patients' perspective. *American Journal*  
8 *of Surgery*, 192, 496-498. doi: 10.1016/j.amjsurg.2006.06.018  
9  
10 Giannousi, Z., Manaras, I., Georgoulis, V., & Samonis, G. (2010). Illness perceptions in  
11 Greek patients with cancer: a validation of the Revised-Illness Perception  
12 Questionnaire. *Psycho-Oncology*, 19, 85-92. doi: 10.1002/pon.1538.  
13  
14 Hagger, M.S., & Orbell, S. (2003). A meta-analytic review of the common-sense model of  
15 illness representations. *Psychology and Health*, 18, 141-184. doi:  
16 10.1080/088704403100081321  
17  
18 Hagger, M.S., & Orbell, S. (2005). A confirmatory factor analysis of the revised illness  
19 perception questionnaire (IPQ-R) in a cervical screening context. *Psychology &*  
20 *Health*, 20, 161-173. doi: 10.1080/0887044042000334724  
21  
22 Harrell, J. C., Dye, W. W., Harvell, D. M., Pinto, M., Jedlicka, P., Sartorius, C. A., &  
23 Horwitz, K. B. (2007). Estrogen insensitivity in a model of estrogen receptor positive  
24 breast cancer lymph node metastasis. *Cancer Research*, 67, 10582-10591. doi:  
25 10.1158/0008-5472.can-07-1655  
26  
27 Hershman, D. L., Shao, T., Kushi, L. H., Buono, D., Tsai, W. Y., Fehrenbacher, L., . . .  
28 Neugut, A. I. (2011). Early discontinuation and non-adherence to adjuvant hormonal  
29 therapy are associated with increased mortality in women with breast cancer. *Breast*  
30 *Cancer Research and Treatment*, 126, 529-537. doi: 10.1007/s10549-010-1132-4  
31  
32 Hershman, D.L., Kushi, L. H., Shao, T., Buono, D., Kershenbaum, A., Tsai, W., . . . Neugut,  
33 A. I. (2010). Early Discontinuation and Nonadherence to Adjuvant Hormonal  
34 Therapy in a Cohort of 8,769 Early-Stage Breast Cancer Patients. *Journal of Clinical*  
35 *Oncology*, 28, 4120-4128. doi: 10.1200/jco.2009.25.9655  
36  
37 Horn, J.L. (1965). A rationale and test for the number of factors in factor analysis.  
38 *Psychometrika*, 30, 179-185. doi: 10.1007/bf02289447  
39  
40 Horne, R., Weinman, J., & Hankins, M. (1999). The beliefs about medicines questionnaire:  
41 The development and evaluation of a new method for assessing the cognitive  
42 representation of medication. *Psychology & Health*, 14, 1-24. doi:  
43 10.1080/08870449908407311  
44  
45 Horne, R., & Weinman, J. (2002). Self-regulation and Self-management in Asthma:  
46 Exploring The Role of Illness Perceptions and Treatment Beliefs in Explaining Non-  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 adherence to Preventer Medication. *Psychology & Health*, 17, 17-32. doi:  
4 10.1080/08870440290001502  
5  
6 Hu, L., & Bentler, P.M. (1999). Cutoff criteria for fit indexes in covariance structure analysis:  
7 Conventional criteria versus new alternatives. *Structural equation modeling: a*  
8 *multidisciplinary journal*, 6, 1-55. doi: 10.1080/10705519909540118  
9  
10 Ibrahim, N., Desa, A., & Chiew-Tong, N.K. (2011). Illness perception and depression in  
11 patients with end-stage renal disease on chronic haemodialysis. *Social Sciences*, 6,  
12 221-226. doi: 10.3923/sscience.2011.221.226  
13  
14 Jacob Arriola, K. R., Mason, T. A., Bannon, K. A., Holmes, C., Powell, C. L., Horne, K., &  
15 O'Regan, R. (2014). Modifiable risk factors for adherence to adjuvant endocrine  
16 therapy among breast cancer patients. *Patient Education and Counselling*, 95, 98-103.  
17 doi: 10.1016/j.pec.2013.12.019  
18  
19 Jopson, N.M., & Moss-Morris, R. (2003). The role of illness severity and illness  
20 representations in adjusting to multiple sclerosis. *Journal of Psychosomatic Research*,  
21 54, 503-511. doi: 10.1016/S0022-3999(02)00455-5  
22  
23 Kaptein, A.A., Schoones, J.W., Fischer, M.J., Thong, M.S.Y, Kroep, J.R., & van der Hoeven,  
24 K.J.M. (2015). Illness Perceptions in Women with Breast Cancer—a Systematic  
25 Literature Review. *Current breast cancer reports*, 7, 117-126. doi:10.1007/s12609-  
26 015-0187-y.  
27  
28 Kostev, K., Haas, G., & Hadji, P. (2012). Persistence in patients with breast cancer treated  
29 with tamoxifen or aromatase inhibitors: Results of a retrospective cohort study. *Value*  
30 *in Health*, 15 , A428. doi: 10.1016/j.jval.2012.08.2146  
31  
32 Leventhal, H., Diefenbach, M., & Leventhal, E. A. Illness cognition: Using common sense to  
33 understand treatment adherence and affect cognition interactions. *Cognitive Therapy*  
34 *and Research*, 16, 143-163. doi: 10.1007/bf01173486  
35  
36 Makubate, B., Donnan, P. T., Dewar, J. A., Thompson, A. M., & McCowan, C. (2013).  
37 Cohort study of adherence to adjuvant endocrine therapy, breast cancer recurrence  
38 and mortality. *British Journal of Cancer*, 108, 1515-1524. doi: 10.1038/bjc.2013.116  
39  
40 Matthews, E. E., Berger, A. M., Schmiede, S. J., Cook, P. F., McCarthy, M. S., Moore, C.  
41 M., & Aloia, M. S. (2014). Cognitive Behavioral Therapy for Insomnia Outcomes in  
42 Women After Primary Breast Cancer Treatment: A Randomized, Controlled Trial.  
43 *Oncology Nursing Forum*, 41, 241-253. doi: 10.1188/14.onf.41-03ap  
44  
45 McCorry, N. K., Scullion, L., McMurray, C. M., Houghton, R., & Dempster, M. (2013).  
46 Content validity of the illness perceptions questionnaire--revised among people with  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 type 2 diabetes: a think-aloud study. *Psychology & Health*, 28, 675-685. doi:  
4 10.1080/08870446.2012.746690  
5  
6 Moon, Z., Moss-Morris, R., Hunter, M., & Hughes, L.D. (2016). *Understanding tamoxifen*  
7 *adherence in women with breast cancer: a qualitative study*. Manuscript submitted for  
8 publication.  
9  
10 Moss-Morris, R., Weinman, J., Petrie, K. J., Horne, R., Cameron, L. D., & Buick, D. (2002).  
11 The revised Illness Perception Questionnaire (IPQ-R). *Psychology & Health*, 17, 1-  
12 16. doi: 10.1080/08870440290001494  
13  
14 Nicholls, E. E., Hill, S., & Foster, N. E. (2013). Musculoskeletal pain illness perceptions:  
15 factor structure of the Illness Perceptions Questionnaire-Revised. *Psychology &*  
16 *Health*, 28, 84-102. doi: 10.1080/08870446.2012.714782  
17  
18 Norton, S., Cosco, T., Doyle, F., Done, J., & Sacker, A. (2013). The Hospital Anxiety and  
19 Depression Scale: A meta confirmatory factor analysis. *Journal of Psychosomatic*  
20 *Research*, 74, 74-81. doi: <http://dx.doi.org/10.1016/j.jpsychores.2012.10.010>  
21  
22 Owusu, C., Buist, D. S. M., Field, T. S., Lash, T. L., Thwin, S. S., Geiger, A. M., . . .  
23 Silliman, R. A. (2008). Predictors of tamoxifen discontinuation among older women  
24 with estrogen receptor-positive breast cancer. *Journal of Clinical Oncology*, 26, 549-  
25 555.  
26  
27 Petrie, K.J., Jago, L.A., & Devcich, D.A. (2007). The role of illness perceptions in patients  
28 with medical conditions. *Current opinion in psychiatry*, 20, 163-167. doi:  
29 10.1097/YCO.0b013e328014a871  
30  
31 Servaes, P., Gielissen, M. F. M., Verhagen, S., & Bleijenberg, G. (2007). The course of  
32 severe fatigue in disease-free breast cancer patients: A longitudinal study. *Psycho-*  
33 *Oncology*, 16, 787-795. doi: 10.1002/pon.1120  
34  
35 Simard, S., Thewes, B., Humphris, G., Dixon, M., Hayden, C., Mireskandari, S., & Ozakinci,  
36 G. (2013). Fear of cancer recurrence in adult cancer survivors: a systematic review of  
37 quantitative studies. *Journal of Cancer Survivorship*, 7, 300-322. doi:  
38 10.1007/s11764-013-0272-z  
39  
40 Stanton, A.L., Petrie, K. J., & Partridge, A.H. (2014). Contributors to nonadherence and  
41 nonpersistence with endocrine therapy in breast cancer survivors recruited from an  
42 online research registry. *Breast Cancer Research and Treatment*, 145, 525-534. doi:  
43 10.1007/s10549-014-2961-3  
44  
45 Tabachnick, B. G., & Fidell, L. S. (2007). Using multivariate statistics. Boston:  
46 Pearson/Allyn & Bacon.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Thewes, B., Butow, P., Bell, M. L., Beith, J., Stuart-Harris, R., Grossi, M., . . . Dalley, D.  
4 (2012). Fear of cancer recurrence in young women with a history of early-stage breast  
5 cancer: a cross-sectional study of prevalence and association with health behaviours.  
6 *Supportive Care in Cancer*, *20*, 2651-2659. doi: 10.1007/s00520-011-1371-x  
7  
8  
9  
10 van Oort, L., Schroder, C., & French, D. P. (2011). What do people think about when they  
11 answer the Brief Illness Perception Questionnaire? A 'think-aloud' study. *British*  
12 *Journal of Health Psychology*, *16*, 231-245. doi: 10.1348/135910710x500819  
13  
14 Weinman, J., Petrie, K.J., Moss-Morris, R., & Horne, R. (1996). The illness perception  
15 questionnaire: a new method for assessing the cognitive representation of illness.  
16 *Psychology and health*, *11*, 431-445. doi: 10.1080/08870449608400270  
17  
18  
19  
20 Westerman, M.J., Hak, T., Sprangers, M.A. G., Groen, H.J. M., Wal, G., & The, A. (2008).  
21 Listen to their answers! Response behaviour in the measurement of physical and role  
22 functioning. *Quality of Life Research*, *17*, 549-558. doi: 10.1007/s11136-008-9333-6  
23  
24  
25 Whittaker, R., Kemp, S., & House, A. (2007). Illness perceptions and outcome in mild head  
26 injury: a longitudinal study. *Journal of Neurology, Neurosurgery & Psychiatry*, *78*,  
27  
28 644-646. doi:10.1136/jnnp.2006.101105  
29  
30  
31 Wittkowski, A., Richards, H.L., Williams, J., & Main, C.J. (2008). Factor analysis of the  
32 Revised Illness Perception Questionnaire in adults with atopic dermatitis. *Psychology,*  
33 *Health & Medicine*, *13*, 346-359. doi: 10.1080/13548500701487697  
34  
35  
36 Zigmond, A. S., & Snaith, R. P. (1983). THE HOSPITAL ANXIETY AND DEPRESSION  
37 SCALE. *Acta Psychiatrica Scandinavica*, *67*, 361-370. doi: 10.1111/j.1600-  
38  
39 0447.1983.tb09716.x  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1.** Interview schedule for qualitative interviews

- 
1. Are there any specific side effects that you have experienced?  
- *Ones that your doctor did not tell you about?*
  2. Do you believe that your previous treatment has cured your breast cancer?
  3. Do you still experience ongoing effects from your previous treatment (chemo, surgery, radio)?
  4. Do you still see yourself as having breast cancer?  
- *What is your relationship with breast cancer?*
  5. What do you see as the main consequences of Tamoxifen?
  6. What do you see as the main consequences of breast cancer?
  7. Do you think that tamoxifen is preventing a risk of recurrence?
  8. What else might be impacting a risk of recurrence?
  9. Is there anything else you can do to control this (prevent risk of recurrence)?
-

**Table 2.** Demographics characteristics of participants.

	<i>Interview study</i> <i>n=18</i>	<i>Think aloud study</i> <i>n=8</i>	<i>Factor analysis</i> <i>n=753</i>	<i>Test-retest</i> <i>reliability</i> <i>n=48</i>
Age <i>mean (SD)</i>	53 (10.2) Range 36 – 63	53 (9.2) Range 37 – 63	53 (10.5) Range 30 – 91	56 (10.3) Range 38 - 82
Race <i>n (%)</i>				
White British	13 (72%)	8 (100%)	654 (87%)	44 (94%)
Other	5 (28%)	0 (0%)	99 (13%)	3 (6%)
Age left full time education <i>n (%)</i>				
16 or under			304 (40%)	25 (52%)
Over 16			449 (60%)	23 (48%)
Menopausal status at diagnosis <i>n (%)</i>				
Pre-menopausal	4 (22%)	2 (25%)	414 (55%)	
Menopausal	2 (11%)	1 (12.5%)	86 (11%)	
Post-menopausal	9 (50%)	4 (50%)	202 (27%)	
Unsure	3 (17%)	1 (12.5%)	33 (4%)	
Months since breast cancer diagnosis <i>Mean (SD)</i>	36 (25) Range 1 year – 5.5 years	25 (19) Range 1 year – 6 years	33 (24) Range 2 months – 16 years	45 (25) Range 1 month – 9 years
Stage at diagnosis <i>n (%)</i>				
Stage I			321 (43%)	
Stage II			339 (45%)	
Stage III			93 (12%)	
Previous treatment <i>n (%)</i>				
Lumpectomy	12 (67%)	5 (63%)	483 (64%)	
Single mastectomy	2 (11%)	1 (13%)	249 (33%)	
Double mastectomy	1 (5%)	2 (25%)	44 (6%)	
Chemotherapy	7 (44%)	3 (38%)	384 (51%)	
Radiotherapy	15 (83%)	6 (75%)	557 (74%)	

Note. SD, Standard deviation. Blank spaces indicate incidences where data was not collected.

**Table 3.** Examples of changes made to the original IPQ-R.

	Previous item	New item
<i>Identity scale</i>		Change in libido, hot flushes, leg cramps, loss of concentration, night sweats, joint pain, vaginal dryness/itchiness/discomfort, feeling down, changes to periods, feeling lightheaded
<i>Timeline acute / chronic (cure)</i>	My illness will last for a long time	My breast cancer is cured
<i>Breast cancer consequences</i>	My illness has major consequences on my life	My breast cancer still has major consequences on my life
<i>Tamoxifen consequences</i>	-	I can't function normally whilst taking tamoxifen
<i>Personal control</i>	My actions will have no effect on the outcome of my illness	My actions will have no effect on the risk of cancer coming back
<i>Treatment control</i>	Tamoxifen treatment can control my illness	Tamoxifen treatment can control my risk of recurrence
<i>Coherence</i>	My breast cancer is a mystery to me	Tamoxifen is a mystery to me
<i>Timeline cyclical (risk of recurrence)</i>	I go through cycles in which my breast cancer gets better and worse	There is a good chance my cancer will come back
<i>Emotional representations</i>	I get depressed when I think about my breast cancer	I get depressed when I think about my risk of recurrence
<i>Causes</i>		Hormonal influence, exercise

**Table 4.** Confirmatory Factor Analysis of the 8 factor IPQ-R.

	1	2	3	4	5	6	7	8	Symptoms attributed to tamoxifen	Causes
<i>Cure</i>										
My treatment has been effective in curing my breast cancer	<b>0.74</b>									
I no longer have breast cancer	<b>0.89</b>									
My breast cancer is cured	<b>0.85</b>									
I still see myself as having cancer	<b>0.81</b>									
<i>Breast cancer consequences</i>										
My breast cancer still has major consequences on my life		<b>0.87</b>								
My breast cancer currently does not have much effect on my life		<b>0.66</b>								
I still experience long lasting effects from my original treatment for breast cancer		<b>0.69</b>								
My breast cancer currently causes difficulties for those who are close to me (e.g. emotional difficulties)		<b>0.75</b>								
<i>Tamoxifen consequences</i>										
Tamoxifen has major consequences on my life			<b>0.63</b>							
I can't function normally whilst taking tamoxifen			<b>0.89</b>							
Taking tamoxifen has had an impact on those around me			<b>0.88</b>							
My work / social life has been affected by taking tamoxifen			<b>0.95</b>							
<i>Risk of recurrence</i>										
There's a good chance my cancer will come back				<b>0.91</b>						
I expect to have a recurrence of cancer in the future				<b>0.95</b>						
I am extremely likely to have a recurrence				<b>0.92</b>						
The chance of my cancer coming back is low				<b>0.72</b>						
<i>Personal control</i>										
There are things I can do to stop the cancer coming back					<b>0.79</b>					
What I do has an influence on whether my cancer comes back					<b>0.77</b>					
There is nothing I can do to help my risk of recurrence					<b>0.87</b>					
My actions will have no effect on the risk of cancer coming back					<b>0.81</b>					

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

*Treatment control*

Tamoxifen treatment can reduce my risk of recurrence	<b>0.82</b>
There is very little that can be done to stop the cancer coming back	<b>0.84</b>
Taking tamoxifen will help stop the cancer coming back	<b>0.78</b>
There is nothing that can help my risk of recurrence	<b>0.82</b>

*Coherence*

Tamoxifen is a mystery to me	<b>0.76</b>
I understand how tamoxifen helps prevent cancer recurrence	<b>0.80</b>
I don't understand how much tamoxifen can help me	<b>0.83</b>
I have a good understanding of why I am taking tamoxifen	<b>0.82</b>

*Emotional representations*

I get depressed when I think about my risk of recurrence	<b>0.91</b>
I worry about my risk of recurrence	<b>0.90</b>
When I think about the cancer coming back I get upset	<b>0.90</b>
My risk of recurrence makes me feel afraid	<b>0.94</b>

---

*Cronbach's alpha*

0.81	0.79	0.87	0.90	0.81	0.76	0.81	0.92
------	------	------	------	------	------	------	------

*Test retest reliability (intraclass correlation coefficient)*

0.92	0.92	0.92	0.87	0.77	0.91	0.91	0.94	0.86	0.87
------	------	------	------	------	------	------	------	------	------

---

**Table 5.** Exploratory factor analysis on the causal items.

	Factor 1: Psychological attributions	Factor 2: Health behaviors	Factor 3: External causes
Stress or worry	<b>.771</b>	.066	-.040
Family problems	<b>.907</b>	-.004	-.014
My own emotional state	<b>.818</b>	-.009	.096
Diet or eating habits	-.004	<b>.840</b>	-.146
My own behaviour	.097	<b>.622</b>	.064
Exercise	-.008	<b>.686</b>	.059
Pollution in the environment	.212	<b>.400</b>	.044
Ageing	.043	.004	<b>.788</b>
Hormonal influence	-.097	.209	<b>.330</b>
Runs in the family	.076	.076	.112
Chance or bad luck	.002	.067	.239
<i>Cronbach alpha</i>	<i>0.85</i>	<i>0.71</i>	<i>0.44</i>

**Table 6.** Correlations between IPQ-R subscales.

	1	2	3	4	5	6	7	8	9
1. Cure	1								
2. Tamoxifen consequences	-.14**	1							
3. Risk of recurrence	-.45**	.23**	1						
4. Breast cancer consequences	-.31**	.49**	.42**	1					
5. Personal control	.15**	-.08*	-.24**	-.13**	1				
6. Treatment control	.23**	-.17**	-.35**	-.22**	.58**	1			
7. Coherence	.10**	-.10**	-.15**	-.16**	.26**	.44**	1		
8. Emotional representations	-.24**	.30**	.41**	.54**	-.15**	-.20**	-.16**	1	
9. Symptoms attributed to tamoxifen	-.12**	.56**	.19**	.36**	.04	.00	.05	.25*	1

\*\*p < 0.01, \*p < 0.05

**Table 7.** Correlations between IPQ-R subscales, HADS distress and BMQ necessity and concerns.

	Distress	Concerns	Necessity beliefs
Cure	-.20**	-.18**	-.04
Tamoxifen consequences	.53**	.56**	.10*
Risk of recurrence	.31**	.19**	.12**
Breast cancer consequences	.55**	.40**	.15**
Personal control	-.15**	-.08*	.02
Treatment control	-.21**	-.23**	.15**
Coherence	-.15**	-.28**	.07
Emotional representations	.45**	.36**	.23**
Symptoms attributed to tamoxifen	.35**	.43**	.09*

\*\*p &lt; 0.001