



## King's Research Portal

DOI:

[10.1016/j.yjmcc.2017.12.007](https://doi.org/10.1016/j.yjmcc.2017.12.007)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Dautova, Y., Kapustin, A., Pappert, K., Epple, M., Okkenhaug, H., Cook, S. J., Shanahan, C. M., Bootman, M. D., & Proudfoot, D. (2017). Calcium phosphate particles stimulate interleukin-1 $\beta$  release from human vascular smooth muscle cells: A role for spleen tyrosine kinase and exosome release. *Journal of Molecular and Cellular Cardiology*. <https://doi.org/10.1016/j.yjmcc.2017.12.007>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal


### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## Accepted Manuscript

Calcium phosphate particles stimulate interleukin-1 $\beta$  release from human vascular smooth muscle cells: A role for spleen tyrosine kinase and exosome release

Yana Dautova, Alexander N. Kapustin, Kevin Pappert, Matthias Epple, Hanneke Okkenhaug, Simon J. Cook, Catherine M. Shanahan, Martin D. Bootman, Diane Proudfoot



PII: S0022-2828(17)30363-2  
DOI: doi:[10.1016/j.yjmcc.2017.12.007](https://doi.org/10.1016/j.yjmcc.2017.12.007)  
Reference: YJMCC 8650

To appear in: *Journal of Molecular and Cellular Cardiology*

Received date: 4 October 2017  
Revised date: 18 December 2017  
Accepted date: 19 December 2017

Please cite this article as: Yana Dautova, Alexander N. Kapustin, Kevin Pappert, Matthias Epple, Hanneke Okkenhaug, Simon J. Cook, Catherine M. Shanahan, Martin D. Bootman, Diane Proudfoot, Calcium phosphate particles stimulate interleukin-1 $\beta$  release from human vascular smooth muscle cells: A role for spleen tyrosine kinase and exosome release. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Yjmcc*(2017), doi:[10.1016/j.yjmcc.2017.12.007](https://doi.org/10.1016/j.yjmcc.2017.12.007)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Calcium phosphate particles stimulate interleukin-1 $\beta$  release from human vascular smooth muscle cells: a role for spleen tyrosine kinase and exosome release**

Yana Dautova<sup>1</sup>; Alexander N. Kapustin<sup>2</sup>, Kevin Pappert<sup>3</sup>, Matthias Epple<sup>3</sup>; Hanneke Okkenhaug<sup>1</sup>, Simon J. Cook<sup>1</sup>; Catherine M. Shanahan<sup>2</sup>, Martin D. Bootman<sup>4</sup>; and Diane Proudfoot<sup>1,\*</sup>

<sup>1</sup>Signalling Programme, Babraham Institute, Babraham, Cambridge, CB22 3AT, U.K

<sup>2</sup>Cardiovascular Division, James Black Centre, King's College London, 125 Coldharbour Lane, London SE5 9NU, U.K

<sup>3</sup>Inorganic Chemistry and Center for Nanointegration Duisburg-Essen (CeNIDE), University of Essen-Duisburg, 45117 Essen, Germany

<sup>4</sup>School of Life, Health and Chemical Sciences, The Open University, Milton Keynes, MK7 6AA, U.K

\*Corresponding author:

Diane Proudfoot,  
Signalling Programme,  
Babraham Institute,  
Babraham,  
Cambridge CB22 3AT,  
U.K.

Tel: +441223 496056

Fax: +441223

496043

## Abstract

**Aims:** Calcium phosphate (CaP) particle deposits are found in several inflammatory diseases including atherosclerosis and osteoarthritis. CaP, and other forms of crystals and particles, can promote inflammasome formation in macrophages leading to caspase-1 activation and secretion of mature interleukin-1 $\beta$  (IL-1 $\beta$ ). Given the close association of small CaP particles with vascular smooth muscle cells (VSMCs) in atherosclerotic fibrous caps, we aimed to determine if CaP particles affected pro-inflammatory signalling in human VSMCs.

**Methods and Results:** Using ELISA to measure IL-1 $\beta$  release from VSMCs, we demonstrated that CaP particles stimulated IL-1 $\beta$  release from proliferating and senescent human VSMCs, but with substantially greater IL-1 $\beta$  release from senescent cells; this required caspase-1 activity but not LPS-priming of cells. Potential inflammasome agonists including ATP, nigericin and monosodium urate crystals did not stimulate IL-1 $\beta$  release from VSMCs. Western blot analysis demonstrated that CaP particles induced rapid activation of spleen tyrosine kinase (SYK) (increased phospho-Y525/526). The SYK inhibitor R406 reduced IL-1 $\beta$  release and caspase-1 activation in CaP particle-treated VSMCs, indicating that SYK activation occurs upstream of and is required for caspase-1 activation. In addition, IL-1 $\beta$  and caspase-1 colocalised in intracellular endosome-like vesicles and we detected IL-1 $\beta$  in exosomes isolated from VSMC media. Furthermore, CaP particle treatment stimulated exosome secretion by VSMCs in a SYK-dependent manner, whilst the exosome-release inhibitor spiroepoxide reduced IL-1 $\beta$  release.

**Conclusions:** CaP particles stimulate SYK and caspase-1 activation in VSMCs, leading to the release of IL-1 $\beta$ , at least in part via exosomes. These novel findings in human VSMCs highlight the pro-inflammatory and pro-calcific potential of microcalcification.

## Keywords

Vascular smooth muscle, calcium phosphate particles, cytokines, caspase-1, SYK, exosomes

## Abbreviations

CaP, calcium phosphate; IL-1 $\beta$ , interleukin-1 $\beta$ ; VSMC, vascular smooth muscle cells; SYK, spleen tyrosine kinase; LPS, lipopolysaccharide; BM, basal culture medium; SFM, serum-free medium; MSU, monosodium urate; ATP, adenosine triphosphate; NLRP3: nucleotide-binding domain, leucine-rich repeat/pyrin domain-containing-3.

## 1. Introduction

Vascular calcification is a regulated process occurring in ageing and diseased blood vessels that correlates positively with cardiovascular deaths. The calcific deposits consist of bone-like nanoparticles that often form aggregates of various sizes and degrees of crystallinity. In particular, the very small calcium phosphate (CaP) particles that appear as 'speckled' deposits in human atherosclerotic plaques are associated with plaque instability, increasing the likelihood of plaque rupture and subsequent thrombus formation (1-3). Recent studies suggest that small CaP particles may be damaging due to effects on mechanical stress in the fibrous cap (4, 5), or that CaP particles can be engulfed by macrophages leading to secretion of pro-inflammatory cytokines (6). Our own studies suggest that small CaP particles are taken up into vascular smooth muscle cells (VSMCs) via endocytosis, macropinocytosis and plasma membrane damage, causing transient intracellular  $\text{Ca}^{2+}$  rises and inducing cell death in subsets of VSMCs (7, 8). Serum proteins, particularly fetuin-A, inhibit cell death by binding CaP particles and delaying particle uptake, membrane damage and dissolution. CaP particles are implicated in the pathogenesis of other diseases including severe calcification of skin arterioles seen in calciphylaxis, degenerative arthritis and breast cancer (9-12). The association of calcification with inflammation *in vivo* either in joints or in blood vessels could be explained by several *in vitro* studies where small particles of CaP were found to interact with neutrophils, macrophages and dendritic cells, subsequently activating various signalling mechanisms that lead to the release of pro-inflammatory cytokines (6, 13-15).

IL-1 $\beta$  has been shown to have a key function in the recruitment of neutrophils *in vivo* in response to monosodium urate (MSU) crystals (16), cholesterol crystals (17) and CaP crystals (18, 19). IL-1 $\beta$  secretion generally requires a two-step process where cells first synthesise pro-IL-1 $\beta$ , followed by a second stimulus to cleave pro-IL-1 $\beta$  to its active form by inflammasome-activated caspase-1, although other proteases such as neutral elastase can also cleave pro-IL-1 $\beta$  (20). The steps involved in crystal interaction with cells leading to the activation of caspase-1 are not fully understood, however, SYK activation, intracellular  $\text{Ca}^{2+}$  increases, phosphorylation of ASC (apoptosis associated speck-like protein containing a CARD) and  $\text{K}^+$  efflux have been reported to occur upstream of caspase-1 activation in various cell types (21, 22). SYK is most commonly known for its role in coupling immune recognition receptors bearing cytoplasmic immunoreceptor tyrosine-based activation motifs (ITAMs) to intracellular signalling pathways but also has roles in cancer, autophagy and in pro-inflammatory responses to calcified particles in osteoarthritic joints (23-25).

VSMCs reside in the media of blood vessels and their function is to maintain vascular tone by co-ordinated contraction and dilation. VSMCs are known to display considerable phenotypic plasticity and migrate from the medial layer into the intima of diseased blood vessels where they become proliferative and secrete extracellular matrix that contributes to the strength of the fibrous cap that overlies the soft atherosclerotic plaque, protecting it from rupture. In addition to VSMCs acquiring osteoblast, chondrocyte or adipocyte-like features, VSMCs secrete several pro-inflammatory cytokines constitutively as the cells age and become senescent (26-29). Furthermore, VSMCs were recently shown to differentiate to a macrophage-like phenotype in a mouse *in vivo* model of atherosclerosis (30).

VSMCs are the main cell type within the fibrous cap region that overlies atherosclerotic plaques, which is thought to form as part of the healing process and to protect the underlying necrotic core from exposure to the blood vessel lumen. Small particles of CaP at this location are associated with plaque rupture in man and are thought to reflect active formation of microcalcification (3, 31-35), making the exploration of CaP particle effects on human VSMCs clinically relevant. With reports that human VSMCs express caspase-1 (36) and that various forms of crystals can activate caspase-1 in several cell types, we aimed to determine if CaP particles could influence human VSMC pro-inflammatory mediator production. We identified a

SYK-dependent pathway in VSMCs leading to caspase-1 activation and release of IL-1 $\beta$  via exosomes after exposure to CaP particles. Our study highlights the pro-inflammatory potential of human VSMCs in an environment where cells are exposed to small CaP particles.

ACCEPTED MANUSCRIPT

## 2. Methods

**2.1. Cell culture.** VSMCs were derived from medial layers of normal human aortae and cultured in smooth muscle basal medium (BM, Lonza) containing 5% foetal bovine serum, insulin, human fibroblast growth factor, human epidermal growth factor and gentamycin, buffered with 3.7 mg/mL NaHCO<sub>3</sub> and 5% CO<sub>2</sub>. Some of the medial VSMCs used in this study were kindly provided by Prof. Martin Bennett (Addenbrooke's Hospital, Cambridge). Donors gave written informed consent for tissue samples to be used for research, on a standard hospital consent form. Ethical approval for use of human VSMC cultures was approved by the Cambridgeshire 1 Research Ethics Committee which conforms to the principles outlined in the Declaration of Helsinki. In some experiments, human aortic VSMCs purchased from Lonza or Promocell were used. Cells from 8 different individuals were used in this study; 4 males (aged 22, 28, 30 and 58 years old) and 4 females (aged 35, 43, 51 and 63 years old). Cells were used between passages 6 and 13 and cultured in BM, unless otherwise indicated.

**2.2. Materials.** CaP particles were prepared as described previously (8). Briefly, aqueous solutions of calcium lactate (9.0 mM; Merck) and diammonium hydrogen phosphate (5.4 mM; Merck) were adjusted to pH 8.0 with NaOH and sterile filtered through a Filtropur S plus unit (0.2 µm). The precipitation reaction was achieved by rapidly pumping (5 mL/min) both solutions into a glass vessel under sterile conditions. The prepared suspensions were immediately centrifuged at 900 rpm for 3 min. The supernatant was removed and the particles were resuspended in 200 times less water volume than the initial nanoparticle dispersion. The concentration of Ca<sup>2+</sup> in the synthesised CaP particles was quantified by atomic absorption spectroscopy (AAS; M-Serie, Thermo Electron) and particle concentrations used in this manuscript are expressed as mg/mL in terms of Ca<sup>2+</sup> content, rather than weight of the nanoparticles. Under the assumption of the stoichiometry of hydroxyapatite for calcium phosphate, Ca<sub>5</sub>(PO<sub>4</sub>)<sub>3</sub>OH, the content of calcium phosphate can be computed as  $w(\text{CaP}) = w(\text{Ca}^{2+}) \cdot M(\text{Ca}_5(\text{PO}_4)_3\text{OH}) / 5 \cdot M(\text{Ca}) = w(\text{Ca}^{2+}) \cdot 2.51$ . The morphology of freshly prepared nanoparticles was characterised with scanning electron microscopy (SEM; ESEM Quanta 400 FEG, gold/palladium sputtering). The particle diameter was between 30 and 60 nm. The particles had a spherical shape after the initial synthesis. Nanoparticle preparations were stored in sterile water at 4 °C and under these conditions, within 2 days the particles changed their shape (or 'ripened', presumably by Ostwald ripening) from a spherical to a needle-like shape. The particles were not colloidal dispersible due to the absence of a surface functionalisation (i.e. a bare, non-coated surface) and therefore agglomerated. Dynamic light scattering in pure water is impossible due to agglomeration. However, we expect a considerable enhancement of the dispersibility in cell-culture media due to the presence of proteins (formation of a stabilizing protein corona). An image displaying the needle-shaped (ripened) CaP particles interacting with human VSMCs is shown in the Supplement Figure S1 and further examples are presented in our previous studies (8). The zeta potential of bare calcium phosphate nanoparticles is slightly negative at neutral pH and will become more negative at increasing pH (and more positive at decreasing pH). All assays were performed with particles stored for at least 2 days. Particles were checked for sterility using an LPS/endotoxin testing kit (stocks of particles had <0.5 EU/mL endotoxin, Pierce). Particle solutions were vortexed immediately prior to addition to cells to achieve a moderate degree of dispersion.

LPS, nigericin, ATP, and CaCl<sub>2</sub> were from Sigma and Na<sub>2</sub>HPO<sub>4</sub> was from VWR. Z-YVAD-FMK was from BioVision. MSU crystals were from Adipogen. R406 was from Selleck, and sc-202721 (spiroepoxide inhibitor) was from Insight Biotechnology. All chemicals were prepared and stored following manufacturer recommendations.

**2.3. Propidium iodide uptake.** An assay using propidium iodide (PI) as a measure of cell death was used as described previously (8).

**2.4. ELISA.** Human VSMCs in culture were exposed to various conditions in the presence or absence of CaP particles and cell supernatants were collected and stored at  $-20^{\circ}\text{C}$  until required for analysis. Samples were defrosted, centrifuged at 13,000 rpm before measuring human IL-1 $\beta$  content using a bead-based Alphasisa (Perkin Elmer) or plate-based multiplex ELISA (Mesoscale discovery).

**2.5. Preparation of cell extracts and Western blotting.** Cells were lysed in ice-cold TG lysis buffer (Tris-HCl, pH 7.4, 20 mM, Triton X-100, 1% (v/v), Glycerol 10% (v/v), NaCl (137 mM), EGTA (1 mM) and  $\text{MgCl}_2$  (1.5 mM), NaF (0.05 M),  $\text{Na}_3\text{VO}_4$  (1 mM), Aprotinin (5  $\mu\text{g}/\text{mL}$ ) Leupeptin (10  $\mu\text{g}/\text{mL}$ ) and PMSF (1 mM) and tested for protein content by BCA assay (Thermo Scientific). SDS-loading buffer was added to lysates, vortexed and heated at  $95^{\circ}\text{C}$  for 5 min before centrifugation at 13,000 rpm and fractionation by SDS-PAGE. Cell supernatants (conditioned medium), were collected on ice into tubes containing PMSF (final concentration 1 mmol/L). SDS-loading buffer was then added and samples were incubated at  $95^{\circ}\text{C}$  for 5 min before centrifugation at 13,000 rpm and fractionation by SDS-PAGE. Briefly, SDS-PAGE gels were transferred to Immobilon P membranes (Millipore), which were blocked in 5% (w/v) powdered milk containing 0.1% (v/v) Tween-20-TBS (tris-buffered saline) and probed with either antibodies recognising pro-IL-1 $\beta$ , SYK (total), phosphorylated SYK (Tyr 525/526) (all from Cell Signalling Technologies), IL-1 $\beta$  (R and D, AF-201), caspase-1 p20 (Adipogen), GAPDH or  $\alpha$ -tubulin (both from ProSci). Mouse anti-human CD63 antibody was from BD Pharmingen. HRP-conjugated antibodies recognising primary antibodies were purchased from BioRad. Immune complexes were visualised using the ECL system (GE Healthcare or high sensitivity substrate, Fisher Scientific). GAPDH or  $\alpha$ -tubulin were used as loading controls and were applied directly to re-probe original blots. In some experiments, Western blot signal intensities were quantified using ImageJ.

**2.6.  $\beta$ -galactosidase staining.** VSMCs were cultured in 12-well plates and assessed for  $\beta$ -galactosidase staining following the protocol of Chen et. al.(37)

**2.7. Caspase-1 activity.** Cells were treated with or without CaP particles for up to 16 hours. Culture medium was then replaced with a fluorescent caspase-1 inhibitor peptide, FAM-YVAD-FLICA, for 1 hour at  $37^{\circ}\text{C}$  following the manufacturer's recommended protocol (FAM-FLICA caspase-1 assay kit, ImmunoChemistry Technologies). Cells were then washed and allowed to incubate for 1 hour in basal culture medium to allow non-bound reagent to elute from cells. Cells were then stained for 5 min with Hoechst 33342 and PI (both 1  $\mu\text{g}/\text{mL}$ ) before placing in physiological buffer (7). Active caspase-1 was assessed qualitatively in cells cultured in 12-well plates on a Zeiss Microbeam microscope and images were processed using ImageJ. Quantitative measurement of active caspase-1 was achieved using cells cultured in glass-bottomed black 96-well plates using an IN Cell 6000 high content screening system. Autofluorescence values for each treatment were corrected by measuring fluorescence in cells with no FAM-FLICA or no PI. Scatter plots in GraphPad were used to display levels of active caspase-1 per cell.

## **2.8. Immunofluorescence**

Cells were grown on sterile glass coverslips, treated with CaP particles and exposed to FAM-YVAD-FLICA as above (section 2.7). After allowing non-bound reagent to elute for 1 hour, cells were fixed in 4% paraformaldehyde, permeabilised in 0.2% Triton X/PBS, blocked for 1 hour in 10% Chemibloc/0.1% Triton X/PBS, washed with 0.1% Triton X/PBS before incubating with a mouse anti-human IL-1 $\beta$  antibody (0.25  $\mu\text{g}/\text{mL}$ , R and D Systems MAB601) for 3 hours. Cells were then washed in 0.1% Triton X/PBS before incubation with anti-mouse IgG NL557 conjugate (1:200 dilution, R and D Systems NL007), washed with 0.1% Triton X/PBS followed by a wash with PBS, before finally mounting in DAPI-containing hard-set mounting medium (Vectashield). Cells were imaged using a Nikon A1R confocal microscope and images were processed using ImageJ.

## 2.9. Exosome isolation and quantification

Quantification of exosomes in VSMC conditioned media was performed by using an anti-CD63-coated bead capturing assay, as described previously (38). Briefly, anti-CD63 antibody (BD Bioscience) was immobilised on 4 µm aldehyde-sulphate latex beads (Invitrogen). VSMCs were plated on 24-well plates (10,000 cells /well) and incubated for 16 hours. The cells were then washed with EBSS and incubated in M199 supplemented with 2.5% exosome-free FBS, 100 U/ml penicillin, 100 µg/ml streptomycin and 2mM L-glutamine in the presence or absence of R406 (1 µM) for 2 hours. Media were then replaced with fresh media with or without the addition of CaP particles (12.5 µg/ml) and cells were cultured for a further 16 hours. Conditioned media were collected and centrifuged at 2,500 x g for 5 minutes. The supernatants were incubated with 1 µl of anti-CD63-coated beads on a shaker overnight at 4 °C. Cells were washed twice in EBSS, trypsinised and quantified using a Nucleocounter NC-3000 (ChemoMetec A, Denmark). Anti-CD63-coated beads were washed with blocking buffer (PBS containing 2% BSA) and incubated with CD81-PE antibody (1:50 in blocking buffer) for 1 hour. Next, anti-CD63-coated beads were washed with blocking buffer and analysed by flow cytometry (BD Accuri™ C6, BD Biosciences). Arbitrary units were calculated as (mean fluorescence units x percentage of positive beads) and normalised to the number of live VSMCs.

Exosomes were purified from VSMC-conditioned media by differential centrifugation as previously described (38). In brief, VSMCs were incubated in the absence or presence of CaP particles (12.5 µg/ml) in M199 supplemented with 2.5% exosome-free FBS, 100 U/ml penicillin, 100 µg/ml streptomycin and 2mM L-glutamine for 16 hours. Next, conditioned media were collected and centrifuged for 5 minutes at 2500 rpm (Thermo Scientific Heraeus Multifuge 3SR+ centrifuge, rotor Sorvall 75006441K) and 30 minutes at 10,000 x g (Sorvall RC6). Exosomes were pelleted from the supernatants by ultracentrifugation at 100,000 x g for 40 minutes at 4 °C (Beckman Coulter Optima Max). Exosomes were washed with PBS, centrifuged again at 100,000 x g and resuspended in PBS.

To prepare cell lysates for comparisons with exosome lysates, VSMCs were washed with PBS and lysed in 0.1M Tris buffer (pH 8.1) supplemented with 0.15 M NaCl, 1% triton X-100 and protease inhibitor cocktail (1:100, Sigma). Cell lysates were sonicated for 5 seconds (Branson Sonifier 150) and centrifuged at 16,353 x g for 15 minutes.

**Statistics.** Relevant statistical analyses were performed using GraphPad Prism and specific statistical tests are detailed in each of the Figure legends. For active caspase-1 measurements in live cells, SPSS was used to perform statistical analysis between cells from different individuals.

### 3. Results

#### 3.1. CaP particles induce IL-1 $\beta$ release from human VSMCs

To establish if CaP particles influenced IL-1 $\beta$  release from human VSMCs, cells were exposed to CaP particles for 16 hours and IL-1 $\beta$  levels were measured in cell supernatants. CaP particles stimulated IL-1 $\beta$  release in the presence or absence of LPS, compared with controls (Figure 1A). We confirmed that CaP particles induced release of the mature form of IL-1 $\beta$  (Supplement Figure S2A). IL-1 $\beta$  accumulated in conditioned media after exposure to CaP particles (Figure 1B), while cellular pro-IL-1 $\beta$  levels did not change after CaP particle exposure (Figure 1C and Supplement Figure S2B and S2C). These data suggest that pro-IL-1 $\beta$  is constitutively expressed in human VSMCs and that CaP particles stimulated processing and release of IL-1 $\beta$ , rather than stimulating *de novo* IL-1 $\beta$  synthesis. CaP particles also stimulated release of TNF $\alpha$ , IL-6, IL-8, DPPIV (CD26), GM-CSF and emmprin (CD147) (Supplement Figure S3). However, we focussed the present study on the release of IL-1 $\beta$  from VSMCs.

During the course of this study, it became clear that VSMCs derived from different individuals had variations in levels of IL-1 $\beta$  release (Figure 1A and B). Additionally, we observed an increase in basal and CaP particle-stimulated IL-1 $\beta$  levels as cells matured in culture, i.e. from early passage (3-5) to mid-passage (6-13) to cells that had undergone replicative senescence (passage 14-18) (Figure 2A); senescence was confirmed by  $\beta$ -galactosidase staining (Figure 2B). No changes in cellular levels of pro-IL-1 $\beta$  were observed in response to CaP particles or between proliferating and senescent cells (Figure 2C, Supplement Figure S2C and S2D). Together, these results suggest that (i) CaP particles stimulate the processing and release of IL-1 $\beta$  from VSMCs and (ii) senescent VSMCs constitutively release IL-1 $\beta$  and exhibit a greater release of IL-1 $\beta$  than proliferating VSMCs in response to CaP particles. Thus, variations observed in IL-1 $\beta$  release from VSMCs were not only donor-dependent but also due to replicative ageing in culture. Nevertheless, addition of CaP particles increased IL-1 $\beta$  release.

To assess the effects of CaP particles compared with other potential inflammasome activators, VSMCs were incubated with monosodium urate (MSU) crystals, nigericin or ATP. However, none of these stimulated IL-1 $\beta$  release from VSMCs (Figure 3 and Supplement Figure S4). Furthermore, exposing cells to high levels of extracellular Ca<sup>2+</sup> or PO<sub>4</sub><sup>3-</sup> did not stimulate IL-1 $\beta$  release (Figure 3 and Supplement Figure S4).

#### 3.2. CaP particles induce IL-1 $\beta$ release via caspase-1

Stimulation of IL-1 $\beta$  release from VSMCs by CaP particles was inhibited by treatment with the caspase-1 inhibitor YVAD (Figure 4A). Live cell imaging of active caspase-1 using a fluorogenic caspase-1 probe (FAM-YVAD-FLICA) revealed active caspase-1 in a low percentage of VSMCs after exposure to CaP particles (Figure 4B, and for quantitative analysis see Figure 6D and Supplement Figure 8). Active caspase-1 was detected either as a single complex, similar to observations of single inflammasome foci seen in macrophages (39), or in a diffuse pattern. Thus, CaP particles stimulated activation of caspase-1 in a small proportion of VSMCs.

A minority of VSMCs contained PI after CaP particle exposure, indicating that some cells underwent necrosis (Figure 4B and for quantitative analysis see Figure 6D and Supplement Figure 8). We therefore tested if conditions that enhanced necrosis affected CaP particle-induced IL-1 $\beta$  release. In the absence of serum, CaP particles induced higher levels of necrosis than when serum was present (Figure 4C), consistent with our previous studies (7). However, serum-free conditions did not enhance the ability of CaP particles to stimulate IL-1 $\beta$  release, despite higher levels of necrosis (Figure 4D). These results suggest that necrosis is

not the main mechanism involved in CaP particle-induced IL-1 $\beta$  release and that serum factors enhance IL-1 $\beta$  release.

We investigated the potential roles of ROS and lysosomal damage in CaP particle-induced IL-1 $\beta$  release, but found that the ROS scavenger Trolox and the pan-cathepsin inhibitor e64 had no effect on IL-1 $\beta$  release (Supplement Figure S5 and S6).

Using confocal analysis to image active caspase-1 and IL-1 $\beta$ , some areas of colocalisation were observed within VSMCs (Figure 5), suggesting that active caspase-1 and IL-1 $\beta$  interact after CaP particle stimulation. The colocalisation foci resembled endosomes, both in the perinuclear area and cell periphery.

### 3.3. SYK phosphorylation is required for caspase-1 activation and IL-1 $\beta$ release

Next, we investigated potential signalling mechanisms activated by CaP particles upstream of caspase-1. We have previously shown by electron microscopy that CaP particles are rapidly endocytosed by VSMCs (8). Inhibitors of endocytosis, wortmannin and chlorpromazine, reduced IL-1 $\beta$  release, while nystatin and Y27632 did not reduce IL-1 $\beta$  levels (Figure 6A). These results suggest that clathrin-mediated endocytosis was involved in particle uptake and IL-1 $\beta$  release.

As SYK activation has been linked with crystal-membrane interactions, endocytosis and inflammasome activation, we investigated SYK activation after CaP particle exposure. CaP particle treatment did not change SYK protein levels but increased phosphorylation at Y525/Y526, in the SYK activation loop (Figure 6B and Supplement Figure S7A and B). In addition, the SYK inhibitor R406 reduced CaP particle-induced IL-1 $\beta$  release in a concentration-dependent manner (Figure 6C). Full inhibition of CaP-induced IL-1 $\beta$  release was observed with R406 (1 or 3  $\mu$ M), and this was the case for each VSMC isolate tested, regardless of whether cells had low or high levels of basal IL-1 $\beta$  release. Furthermore, R406 inhibited caspase-1 activity after CaP particle exposure, as assessed by quantitative live cell imaging (Figure 6D), with no effect on VSMC death (Supplement Figure S8 and S9). This suggests that SYK activation occurs prior to and is required for caspase-1 activation and thence IL-1 $\beta$  in response to CaP particles.

### 3.4. Exosomes as a route for IL-1 $\beta$ release

To assess whether vesicle release may be a mechanism for IL-1 $\beta$  release from VSMCs, cells were incubated with an exosome release inhibitor, spiroepoxide (38). Spiroepoxide reduced CaP particle-induced IL-1 $\beta$  release, suggesting that exosomes are involved in the release of IL-1 $\beta$  (Figure 7A). We next measured exosome release from VSMCs in response to CaP particles and found that CaP particles stimulated exosome release (Figure 7B). Additionally, the SYK inhibitor R406 inhibited CaP particle-induced exosome release, suggesting that SYK phosphorylation is involved in CaP particle-stimulated exosome release (Figure 7B).

To confirm if IL-1 $\beta$  is secreted in exosomes, we isolated exosomes from conditioned media of VSMCs treated with or without CaP particles. We previously showed that VSMC exosomes are enriched with tetraspanin CD63 (38) and correspondingly, Western blot analysis revealed that exosomes enriched with CD63 contained the pro-form of IL-1 $\beta$  (31 kDa) (Figure 7C). Notably, CaP particle-treated and control cells released exosomes with similar levels of IL-1 $\beta$ , which suggests that the increase in released IL-1 $\beta$  detected by ELISA after CaP particle treatment may be due to increased numbers of released vesicles, rather than selective loading of released exosomes with IL-1 $\beta$ .

The mature, active, 17 kDa form of IL-1 $\beta$  was not detected in cells or exosomes but a 28 kDa band was observed in cells in addition to the 31 kDa pro-IL-1 $\beta$  (Figure 7C, Supplement Figure

S2B). The 28 kDa form of IL-1 $\beta$  has been reported in other studies as an intermediate form, which is generated upon the first step in caspase-1-mediated cleavage of pro-IL-1 $\beta$  (40). Although we found active caspase-1 at foci within cells, indicating intracellular caspase-1 activity, active caspase-1 was also detected in cell supernatants *by Western Blotting* (Supplement Figure S10), suggesting that some IL-1 $\beta$  processing may also occur extracellularly.

ACCEPTED MANUSCRIPT

#### 4. Discussion

Here we report that CaP particles induce the release of IL-1 $\beta$  from human VSMCs via activation of SYK and caspase-1. The levels of basal and CaP particle-stimulated IL-1 $\beta$  release varied between VSMCs isolated from different donors and also with stage of cell culture; from early passage cells, where IL-1 $\beta$  levels were lowest, to cells that had undergone replicative senescence, where IL-1 $\beta$  levels were highest. This suggests that senescent VSMCs exposed to CaP particles have the greatest pro-inflammatory potential, compared with proliferating cells, which is particularly relevant to atherosclerotic plaque caps where VSMCs that have undergone senescence are thought to weaken plaques and encourage plaque instability, in aged populations in particular (41, 42). Why VSMCs are already 'primed' to express pro-IL-1 $\beta$  is not known but may be related to their de-differentiated, non-contractile phenotype in culture that is commonly observed in diseased blood vessels *in vivo*. Various mechanisms have been proposed for the basal, unstimulated cytokine release by senescent VSMCs, including altered signalling pathways and genomic instability (28, 29, 43, 44).

Our studies revealed that human VSMCs do not show enhanced release of IL-1 $\beta$  in response to several of the common inflammasome agonists reported to activate caspase-1 in macrophages, including ATP, nigericin and MSU crystals. This suggests that either: (1) human VSMCs do not share the same upstream inflammasome-activating mechanisms as other cell types (e.g. VSMCs lose purinergic receptor expression in culture (45)), and that CaP particle composition or physical parameters such as shape, size and charge may be important in inflammasome activation in VSMCs; or (2) that CaP particles may have additional stimulatory effects on release mechanisms for IL-1 $\beta$ . Potential dissolution of the particles being responsible for the effects on IL-1 $\beta$  release was unlikely, since elevated extracellular levels of either Ca<sup>2+</sup> or PO<sub>4</sub><sup>3-</sup> did not mimic the effects of CaP particles. Thus, VSMCs appeared to be particularly sensitive to CaP particles in terms of stimulating IL-1 $\beta$  release.

Using live cell imaging, active caspase-1 was detected in a minority of VSMCs exposed to CaP particles, perhaps reflecting transient activity of caspase-1. Alternatively, since VSMCs are well known to be heterogeneous, certain phenotypes within the population of cells may be intrinsically primed to activate caspase-1 in response to CaP particles. Further investigation including phenotyping of cells would help to elucidate why some VSMCs have higher potential for activating caspase-1. The finding of high levels of active caspase-1 in subsets of cells and the assumption that IL-1 $\beta$  is activated within these cells may explain why we observed relatively low levels of IL-1 $\beta$  release from VSMCs (mostly in the picomolar range), compared with other cell types such as macrophages, where agonists activate caspase-1 in almost the whole cell population, depending on the potency of the stimulus (46). The different primary cell isolates used in the current study were from 8 healthy donors (4 males and 4 females) and in general, we found that responses to CaP particles were similar between males and females. However, as we used a relatively small number of donors, this is a limitation of the study. It would be interesting to investigate if CaP particles stimulated caspase-1 activation and IL-1 $\beta$  release at higher levels or in the majority of VSMCs derived from a larger cohort of aged or diseased blood vessels. A recent study of over 100 individuals identified a sub-group of older people with hypertension that had elevated inflammasome gene expression profiles and constitutive IL-1 $\beta$  expression (47). Since calcification is closely linked with hypertension, ageing and cardiovascular mortality (48), it would be particularly interesting to investigate caspase-1 activation and IL-1 $\beta$  release from VSMCs derived from hypertensive patients.

Having established that IL-1 $\beta$  was released by VSMCs in response to CaP particles, we sought to establish which mechanisms control its release. IL-1 $\beta$  is a leaderless protein and is not secreted by the classical secretory pathway. Several mechanisms for its release have been proposed including microvesicle shedding from the plasma membrane, secretion via exosomes, exocytosis of secretory lysosomes, secretion across the plasma membrane during pyroptotic death, secretory autophagy and passive loss during necrosis (49-51). Confocal images of VSMCs suggest IL-1 $\beta$  may be released via vesicle-like structures, most likely late

endosomes/multivesicular bodies, and this concept was supported by the finding that inhibition of exosome secretion reduced CaP particle-induced IL-1 $\beta$  release. It is also possible that necrosis contributed to the release of IL-1 $\beta$ , since PI positive cells were detected after 2 hours of treatment with CaP particles. However, in serum-free conditions where levels of necrosis were higher than when serum was present, CaP particle-induced IL-1 $\beta$  levels were found to be lower, arguing against a necrosis-dependent mechanism causing IL-1 $\beta$  loss. The reason for lower IL-1 $\beta$  release in serum-free conditions could be due to several mechanisms such as: (1) serum being required for efficient IL-1 $\beta$  release; (2) because of cell death, fewer cells are available to respond to CaP particles and activate signalling leading to IL-1 $\beta$  release; or (3) that cells entering quiescence release less IL-1 $\beta$ . However, the latter possibility is less likely as we observed that highly proliferative cells from very early passages released the lowest levels of IL-1 $\beta$ . Additionally, we found that SYK inhibition reduced caspase-1 activation but did not reduce numbers of PI-positive cells, suggesting different mechanisms for CaP particle-induced activation of SYK and CaP particle-induced necrosis. Furthermore, quantitative analysis of PI levels and caspase-1 levels in VSMCs revealed that cells with high PI (dead cells) had relatively low levels of active caspase-1, suggesting that pyroptosis is not involved. However, the kinetics and extent of CaP particle damage to VSMCs and its relation to SYK activation require further investigation, especially in regard to studies reporting that slow-onset necrosis stimulates the NLRP3 inflammasome, while rapid necrosis does not (52). Our results suggest that in an environment with low serum, CaP particles will induce substantial VSMC death and IL-1 $\beta$  release, while in a serum-containing environment VSMC death is negligible, but signalling leading to IL-1 $\beta$  release is greater. In either scenario, inflammation is the expected result.

SYK activation is known to be linked with crystal-plasma membrane interactions and endocytosis, and may have a large range of effects on downstream signalling such as PI3K and Akt activation (53-56). We previously found that CaP particles are taken up by VSMCs via endocytosis or macropinocytosis within 5-10 minutes of exposure to CaP particles, as demonstrated by electron microscopy (8) and here we found that inhibition of clathrin-mediated endocytosis reduced the effects of CaP particles on IL-1 $\beta$  release. This suggested that particle uptake, rather than membrane surface activation, is important in initiating signalling leading to IL-1 $\beta$  release. However, since endocytosis inhibitors such as wortmannin can have off-target effects, further studies are required to investigate the mechanisms involved in the endocytosis of CaP particles and how this might lead to SYK activation. SYK was activated rapidly in VSMCs, consistent with kinetics of activation reported in other cell types (57) and the highly selective SYK inhibitor, R406, inhibited CaP-induced IL-1 $\beta$  release. SYK is involved in various biological processes including focal plasma membrane damage-induced necroptosis in red blood cells (58), while overexpression of SYK in tumour cells leads to senescence (59). However, it appeared that in VSMCs, SYK activation, rather than raised SYK protein levels, was important in IL-1 $\beta$  release. *Inhibition of SYK phosphorylation with R406 also inhibited caspase-1 activation, indicating that SYK phosphorylation occurs upstream of caspase-1 and is required for caspase-1 activation. It will be interesting in future studies to investigate the intermediate steps that link SYK phosphorylation to caspase-1 activation, such as other kinases and which inflammasome is involved.*

We found that the highly selective inhibitor of neutral sphingomyelinase, spiroepoxide, which inhibits exosome release, inhibited CaP particle-induced IL-1 $\beta$  release. Further investigation found that CaP particles stimulated exosome release, suggested that IL-1 $\beta$  may be exported from VSMCs via exosomes. Supporting this hypothesis, we found that R406 inhibited CaP particle-induced exosome release and that exosomes contained IL-1 $\beta$ . SYK involvement in exosome release has been suggested by others (60) and raises the possibility that R406 inhibits IL-1 $\beta$  release by more than one mechanism, i.e. inhibition of caspase-1 activation and inhibition of exosome release. We demonstrated a role for exosomes in IL-1 $\beta$  release but secretion via other extracellular bodies and/or microvesicles cannot be excluded and require further investigation. Exosome release is known to be increased in senescent cells, which may

help to explain why higher basal release of IL-1 $\beta$  was observed in senescent VSMCs. In conditions such as those found in the environment of the atherosclerotic plaque, exosomes released from VSMCs have been shown to initiate and propagate mineralisation (38). Thus, a reduction in exosome release by R406 would also be expected to reduce calcification, although this remains to be tested experimentally.

Intriguingly, the mature, active, 17 kDa form of IL-1 $\beta$  was detected in VSMC supernatants but not in cells or exosomes. The pro-form of IL-1 $\beta$  (31 kDa) and an additional band at 28 kDa was observed. A lack of detection of mature IL-1 $\beta$  may have been due to low levels in cells or exosomes, or perhaps due to rapid release of active IL-1 $\beta$ . The presence of a 28 kDa form of IL-1 $\beta$  has been described as an intermediate, first step in caspase-1 cleavage (40, 61). Thus, IL-1 $\beta$  was detected intracellularly in VSMCs in a pro- and intermediate cleaved form, in exosomes in a pro-form, while IL-1 $\beta$  in its pro-, intermediate cleaved and mature form was detected in cell supernatants. It should be noted that levels of cleaved forms of IL-1 $\beta$  were low, relative to pro-IL-1 $\beta$  levels. This observation raises the question of how and where IL-1 $\beta$  is cleaved to generate the mature form. Clearly, the initial step in pro-IL-1 $\beta$  cleavage was detected within cells and the presence of active caspase-1 within foci in VSMCs suggests intracellular caspase-1 activity. However, active caspase-1 was also found in cell supernatants, raising the possibility that some IL-1 $\beta$  processing may occur extracellularly in VSMC cultures. Although most studies in macrophages describe IL-1 $\beta$  cleavage by caspase-1 within cells before IL-1 $\beta$  is released, *the cleaved form of caspase-1 is often detected in cell supernatants* (56, 62) and a recent study highlighted the detection of extracellular active caspase-1 released from human monocyte/macrophages stimulated with LPS and ATP (63). The idea of extracellular processing of pro-IL-1 $\beta$  is not new and other proteases, such as neutrophil and mast cell-derived proteases have been implicated in pro-inflammatory environments (reviewed in (61)). This highlights that even if inactive forms of IL-1 $\beta$  are released by cells, they may be important in generating active IL-1 $\beta$  *in vivo*. *Additionally, several studies have reported that exosomes were implicated in the transfer of fully functional cytokines and growth factors (64-66). Further studies are required to determine how IL-1 $\beta$  is packaged and released from VSMCs exosomes, and to what extent it is functional.*

Mature IL-1 $\beta$  is a potent cytokine known to stimulate neutrophil migration, cause vascular permeability and induce further calcification, all important disease-amplifying mechanisms. Blockade of IL-1 $\beta$  activation in atherosclerotic plaques is desirable, as reported in a recent clinical trial that successfully targeted IL-1 $\beta$  and reduced cardiovascular event rates (67). The SYK inhibitor used in this study, R406 is the orally active metabolite of fostamatinib, which is currently in clinical trials for autoimmune thrombocytopenia, haemolytic anaemia and IgA nephropathy. In previous clinical trials fostamatinib was found to be ineffective in treating rheumatoid arthritis but it was well tolerated. In a mouse model of atherosclerosis, fostamatinib was effective in reducing atherosclerotic lesion size and macrophage infiltration (68). Thus, SYK inhibition may have therapeutic potential in the vasculature and perhaps also at other locations where CaP particles appear to be damaging, such as in osteoarthritis.

In conclusion, our studies in primary human VSMCs suggest that CaP particles activate SYK and caspase-1 leading to IL-1 $\beta$  release via exosomes. If similar mechanisms occur *in vivo*, inhibition of SYK could potentially interfere with the damaging effects of microcalcification in VSMC-rich fibrous caps in addition to other proinflammatory targets in the diseased blood vessel wall. Further work is now required to decipher: how CaP particles activate SYK in VSMCs; exactly how this leads to caspase-1 activation, exosome and IL-1 $\beta$  release; and to explore the importance of CaP particle-induced cytokine release *in vivo*.

**5. Funding**

This work was funded by the British Heart Foundation (Career Re-entry Fellowship to D.P. FS/11/21/28691 and a project grant to AK PG/17/37/33023) and supported by an Institute Strategic Programme Grant from the Biotechnology and Biological Sciences Research Council (BB/P013384/1).

**6. Acknowledgments**

We wish to thank Dr. Diana Kozlova for preparation of particles (University of Essen-Duisburg), Dr. Simon Walker (Babraham Institute) for assistance with imaging, Dr. Anne Segonds-Pichon (Babraham Institute) for assistance with statistics, Dr. Hannah Johnson (Babraham Institute) for advice on SYK activation and Dr. Jeremy Skepper (Cambridge University) for assistance with electron microscopy.

**7. Conflict of Interest**

Conflict of Interest: none declared.

## 8. References

1. Virmani R, Burke AP, Kolodgie FD, Farb A. Pathology of the thin-cap fibroatheroma: a type of vulnerable plaque. *J Interv Cardiol.* 2003;16(3):267-72.
2. Ehara S, Kobayashi Y, Yoshiyama M, Shimada K, Shimada Y, Fukuda D, et al. Spotty calcification typifies the culprit plaque in patients with acute myocardial infarction: an intravascular ultrasound study. *Circulation.* 2004;110(22):3424-9.
3. Vengrenyuk Y, Carlier S, Xanthos S, Cardoso L, Ganatos P, Virmani R, et al. A hypothesis for vulnerable plaque rupture due to stress-induced debonding around cellular microcalcifications in thin fibrous caps. *Proc Natl Acad Sci U S A.* 2006;103(40):14678-83.
4. Hutcheson JD, Maldonado N, Aikawa E. Small entities with large impact: microcalcifications and atherosclerotic plaque vulnerability. *Current opinion in lipidology.* 2014;25(5):327-32.
5. Maldonado N, Kelly-Arnold A, Vengrenyuk Y, Laudier D, Fallon JT, Virmani R, et al. A mechanistic analysis of the role of microcalcifications in atherosclerotic plaque stability: potential implications for plaque rupture. *American journal of physiology Heart and circulatory physiology.* 2012;303(5):H619-28.
6. Nadra I, Mason JC, Philippidis P, Florey O, Smythe CD, McCarthy GM, et al. Proinflammatory activation of macrophages by basic calcium phosphate crystals via protein kinase C and MAP kinase pathways: a vicious cycle of inflammation and arterial calcification? *Circ Res.* 2005;96(12):1248-56.
7. Ewence AE, Bootman M, Roderick HL, Skepper JN, McCarthy G, Epple M, et al. Calcium phosphate crystals induce cell death in human vascular smooth muscle cells: a potential mechanism in atherosclerotic plaque destabilization. *Circ Res.* 2008;103(5):e28-34.
8. Dautova Y, Kozlova D, Skepper JN, Epple M, Bootman MD, Proudfoot D. Fetuin-A and albumin alter cytotoxic effects of calcium phosphate nanoparticles on human vascular smooth muscle cells. *PLoS one.* 2014;9(5):e97565.
9. Molloy ES, McCarthy GM. Hydroxyapatite deposition disease of the joint. *Current rheumatology reports.* 2003;5(3):215-21.
10. Fuerst M, Niggemeyer O, Lammers L, Schafer F, Lohmann C, Ruther W. Articular cartilage mineralization in osteoarthritis of the hip. *BMC musculoskeletal disorders.* 2009;10:166.
11. Hendriks EJ, de Jong PA, van der Graaf Y, Mali WP, van der Schouw YT, Beulens JW. Breast arterial calcifications: a systematic review and meta-analysis of their determinants and their association with cardiovascular events. *Atherosclerosis.* 2015;239(1):11-20.
12. Scimeca M, Giannini E, Antonacci C, Pistolesse CA, Spagnoli LG, Bonanno E. Microcalcifications in breast cancer: an active phenomenon mediated by epithelial cells with mesenchymal characteristics. *BMC cancer.* 2014;14:286.
13. Morgan MP, McCarthy GM. Signaling mechanisms involved in crystal-induced tissue damage. *Curr Opin Rheumatol.* 2002;14(3):292-7.
14. Pazar B, Ea HK, Narayan S, Kolly L, Bagnoud N, Chobaz V, et al. Basic calcium phosphate crystals induce monocyte/macrophage IL-1 $\beta$  secretion through the NLRP3 inflammasome in vitro. *J Immunol.* 2011;186(4):2495-502.
15. Smith ER, Hanssen E, McMahon LP, Holt SG. Fetuin-A-containing calciprotein particles reduce mineral stress in the macrophage. *PLoS one.* 2013;8(4):e60904.
16. Martin WJ, Herst PM, Chia EW, Harper JL. Sesquiterpene dialdehydes inhibit MSU crystal-induced superoxide production by infiltrating neutrophils in an in vivo model of gouty inflammation. *Free radical biology & medicine.* 2009;47(5):616-21.
17. Duewell P, Kono H, Rayner KJ, Sirois CM, Vladimer G, Bauernfeind FG, et al. NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature.* 464(7293):1357-61.
18. Jin C, Frayssinet P, Pelker R, Cwirka D, Hu B, Vignery A, et al. NLRP3 inflammasome plays a critical role in the pathogenesis of hydroxyapatite-associated arthropathy. *Proc Natl Acad Sci U S A.* 2011;108(36):14867-72.

19. Narayan S, Pazar B, Ea HK, Kolly L, Bagnoud N, Chobaz V, et al. Octacalcium phosphate crystals induce inflammation in vivo through interleukin-1 but independent of the NLRP3 inflammasome in mice. *Arthritis and rheumatism*. 2011;63(2):422-33.
20. Alfaidi M, Wilson H, Daigneault M, Burnett A, Ridger V, Chamberlain J, et al. Neutrophil elastase promotes interleukin-1 $\beta$  secretion from human coronary endothelium. *The Journal of biological chemistry*. 2015;290(40):24067-78.
21. Gross O, Poeck H, Bscheider M, Dostert C, Hanneschlager N, Endres S, et al. Syk kinase signalling couples to the Nlrp3 inflammasome for anti-fungal host defence. *Nature*. 2009;459(7245):433-6.
22. Kistowska M, Fenini G, Jankovic D, Feldmeyer L, Kerl K, Bosshard P, et al. *Malassezia* yeasts activate the NLRP3 inflammasome in antigen-presenting cells via Syk-kinase signalling. *Experimental dermatology*. 2014;23(12):884-9.
23. Krisenko MO, Higgins RL, Ghosh S, Zhou Q, Trybula JS, Wang WH, et al. Syk is Recruited to Stress Granules and Promotes their Clearance through Autophagy. *The Journal of biological chemistry*. 2015.
24. Cunningham CC, Mills E, Mielke LA, O'Farrell LK, Lavelle E, Mori A, et al. Osteoarthritis-associated basic calcium phosphate crystals induce pro-inflammatory cytokines and damage-associated molecules via activation of Syk and PI3 kinase. *Clin Immunol*. 2012;144(3):228-36.
25. Coopman PJ, Do MT, Barth M, Bowden ET, Hayes AJ, Basyuk E, et al. The Syk tyrosine kinase suppresses malignant growth of human breast cancer cells. *Nature*. 2000;406(6797):742-7.
26. Iyemere VP, Proudfoot D, Weissberg PL, Shanahan CM. Vascular smooth muscle cell phenotypic plasticity and the regulation of vascular calcification. *J Intern Med*. 2006;260(3):192-210.
27. Burton DG, Giles PJ, Sheerin AN, Smith SK, Lawton JJ, Ostler EL, et al. Microarray analysis of senescent vascular smooth muscle cells: A link to atherosclerosis and vascular calcification. *Experimental gerontology*. 2009;44(10):659-65.
28. Ragnauth CD, Warren DT, Liu Y, McNair R, Tajsic T, Figg N, et al. Prelamin A acts to accelerate smooth muscle cell senescence and is a novel biomarker of human vascular aging. *Circulation*. 2010;121(20):2200-10.
29. Song Y, Shen H, Schenten D, Shan P, Lee PJ, Goldstein DR. Aging enhances the basal production of IL-6 and CCL2 in vascular smooth muscle cells. *Arteriosclerosis, thrombosis, and vascular biology*. 2012;32(1):103-9.
30. Feil S, Fehrenbacher B, Lukowski R, Essmann F, Schulze-Osthoff K, Schaller M, et al. Transdifferentiation of vascular smooth muscle cells to macrophage-like cells during atherogenesis. *Circ Res*. 2014;115(7):662-7.
31. Ruiz JL, Hutcheson JD, Aikawa E. Cardiovascular calcification: current controversies and novel concepts. *Cardiovascular pathology : the official journal of the Society for Cardiovascular Pathology*. 2015;24(4):207-12.
32. Cilla M, Pena E, Martinez MA. 3D computational parametric analysis of eccentric atheroma plaque: influence of axial and circumferential residual stresses. *Biomechanics and modeling in mechanobiology*. 2012;11(7):1001-13.
33. Nakahara T, Dweck MR, Narula N, Pisapia D, Narula J, Strauss HW. Coronary Artery Calcification: From Mechanism to Molecular Imaging. *JACC Cardiovascular imaging*. 2017;10(5):582-93.
34. Pugliese G, Iacobini C, Blasetti Fantauzzi C, Menini S. The dark and bright side of atherosclerotic calcification. *Atherosclerosis*. 2015;238(2):220-30.
35. Irkle A, Vesey AT, Lewis DY, Skepper JN, Bird JL, Dweck MR, et al. Identifying active vascular microcalcification by (18)F-sodium fluoride positron emission tomography. *Nature communications*. 2015;6:7495.
36. Young JL, Sukhova GK, Foster D, Kisiel W, Libby P, Schonbeck U. The serpin proteinase inhibitor 9 is an endogenous inhibitor of interleukin 1 $\beta$ -converting enzyme (caspase-1) activity in human vascular smooth muscle cells. *J Exp Med*. 2000;191(9):1535-44.

37. Chen J-H, Ozanne SE, Hales CN. Methods of Cellular Senescence Induction Using Oxidative Stress. In: Tollesbolto, editor. *Biological Aging: Methods and Protocols*. Totowa, NJ: Humana Press; 2007. p. 179-89.
38. Kapustin AN, Chatrou ML, Drozdov I, Zheng Y, Davidson SM, Soong D, et al. Vascular smooth muscle cell calcification is mediated by regulated exosome secretion. *Circ Res*. 2015;116(8):1312-23.
39. Baroja-Mazo A, Martin-Sanchez F, Gomez AI, Martinez CM, Amores-Iniesta J, Compan V, et al. The NLRP3 inflammasome is released as a particulate danger signal that amplifies the inflammatory response. *Nature immunology*. 2014;15(8):738-48.
40. Swaan PW, Knoell DL, Helder F, Wewers MD. Sequential processing of human ProIL-1beta by caspase-1 and subsequent folding determined by a combined in vitro and in silico approach. *Pharmaceutical research*. 2001;18(8):1083-90.
41. Wang J, Uryga AK, Reinhold J, Figg N, Baker L, Finigan A, et al. Vascular Smooth Muscle Cell Senescence Promotes Atherosclerosis and Features of Plaque Vulnerability. *Circulation*. 2015;132(20):1909-19.
42. Bennett MR, Sinha S, Owens GK. Vascular Smooth Muscle Cells in Atherosclerosis. *Circ Res*. 2016;118(4):692-702.
43. Bautista Nino PK, Durik M, Danser AH, de Vries R, Musterd-Bhaggoe UM, Meima ME, et al. Phosphodiesterase 1 regulation is a key mechanism in vascular aging. *Clin Sci (Lond)*. 2015;129(12):1061-75.
44. Liu R, Lo L, Lay AJ, Zhao Y, Ting KK, Robertson EN, et al. ARHGAP18 Protects Against Thoracic Aortic Aneurysm Formation by Mitigating the Synthetic and Pro-Inflammatory Smooth Muscle Cell Phenotype. *Circ Res*. 2017.
45. Erlinge D. Extracellular ATP: a growth factor for vascular smooth muscle cells. *General pharmacology*. 1998;31(1):1-8.
46. Liu T, Yamaguchi Y, Shirasaki Y, Shikada K, Yamagishi M, Hoshino K, et al. Single-cell imaging of caspase-1 dynamics reveals an all-or-none inflammasome signaling response. *Cell reports*. 2014;8(4):974-82.
47. Furman D CJ, Lartrigue L, Bolen CR, Haddad F, Gaudilliere B, Ganio EA, Fragiadakis GK, Spitzer MH, Douchet I, Daburon S, Moreau J-F, Nolan GP, Blanco P, Dechanet-Merville J, Dekker CL, Jojic V, Kuo CJ, Davis MM, Faustin B. Expression of specific inflammasome gene modules stratifies older individuals into two extreme clinical and immunological states. *Nature*. 2017;23(2):174-84. E48.
48. Kalra SS. Vascular calcification and hypertension: cause and effect. *Ann Med*. 2012;44(Suppl 1):S85-92.
49. Piccioli P, Rubartelli A. The secretion of IL-1beta and options for release. *Seminars in immunology*. 2013;25(6):425-9.
50. Cullen SP, Kearney CJ, Clancy DM, Martin SJ. Diverse Activators of the NLRP3 Inflammasome Promote IL-1beta Secretion by Triggering Necrosis. *Cell reports*. 2015;11(10):1535-48.
51. Qu Y, Franchi L, Nunez G, Dubyak GR. Nonclassical IL-1 beta secretion stimulated by P2X7 receptors is dependent on inflammasome activation and correlated with exosome release in murine macrophages. *J Immunol*. 2007;179(3):1913-25.
52. Li H, Ambade A, Re F. Cutting edge: Necrosis activates the NLRP3 inflammasome. *J Immunol*. 2009;183(3):1528-32.
53. Mocsai A, Ruland J, Tybulewicz VL. The SYK tyrosine kinase: a crucial player in diverse biological functions. *Nature reviews Immunology*. 2010;10(6):387-402. Epub 2010/05/15.
54. Ng G, Sharma K, Ward SM, Desrosiers MD, Stephens LA, Schoel WM, et al. Receptor-independent, direct membrane binding leads to cell-surface lipid sorting and Syk kinase activation in dendritic cells. *Immunity*. 2008;29(5):807-18. Epub 2008/11/11.
55. Lau C, Wang X, Song L, North M, Wiehler S, Proud D, et al. Syk associates with clathrin and mediates phosphatidylinositol 3-kinase activation during human rhinovirus internalization. *J Immunol*. 2008;180(2):870-80. Epub 2008/01/08.
56. Migita K, Izumi Y, Jiuchi Y, Kozuru H, Kawahara C, Nakamura M, et al. Serum amyloid A induces NLRP-3-mediated IL-1beta secretion in neutrophils. *PLoS one*. 2014;9(5):e96703.

57. Vigano E, Diamond CE, Spreafico R, Balachander A, Sobota RM, Mortellaro A. Human caspase-4 and caspase-5 regulate the one-step non-canonical inflammasome activation in monocytes. *Nature communications*. 2015;6:8761.
58. LaRocca TJ, Stivison EA, Mal-Sarkar T, Hooven TA, Hod EA, Spitalnik SL, et al. CD59 signaling and membrane pores drive Syk-dependent erythrocyte necroptosis. *Cell death & disease*. 2015;6:e1773.
59. Bailet O, Fenouille N, Abbe P, Robert G, Rocchi S, Gonthier N, et al. Spleen tyrosine kinase functions as a tumor suppressor in melanoma cells by inducing senescence-like growth arrest. *Cancer research*. 2009;69(7):2748-56.
60. Mfunyi CM, Vaillancourt M, Vitry J, Nsimba Batomene TR, Posvanzic A, Lambert AA, et al. Exosome release following activation of the dendritic cell immunoreceptor: a potential role in HIV-1 pathogenesis. *Virology*. 2015;484:103-12.
61. Afonina IS, Muller C, Martin SJ, Beyaert R. Proteolytic Processing of Interleukin-1 Family Cytokines: Variations on a Common Theme. *Immunity*. 2015;42(6):991-1004.
62. Laliberte RE, Egger J, Gabel CA. ATP treatment of human monocytes promotes caspase-1 maturation and externalization. *The Journal of biological chemistry*. 1999;274(52):36944-51.
63. Shamaa OR, Mitra S, Gavrilin MA, Wewers MD. Monocyte Caspase-1 Is Released in a Stable, Active High Molecular Weight Complex Distinct from the Unstable Cell Lysate-Activated Caspase-1. *PLoS one*. 2015;10(11):e0142203.
64. Webber J, Steadman R, Mason MD, Tabi Z, Clayton A. Cancer exosomes trigger fibroblast to myofibroblast differentiation. *Cancer research*. 2010;70(23):9621-30.
65. Luga V, Zhang L, Vitoria-Petit AM, Ogunjimi AA, Inanlou MR, Chiu E, et al. Exosomes mediate stromal mobilization of autocrine Wnt-PCP signaling in breast cancer cell migration. *Cell*. 2012;151(7):1542-56.
66. Cossetti C, Iraci N, Mercer TR, Leonardi T, Alpi E, Drago D, et al. Extracellular vesicles from neural stem cells transfer IFN-gamma via Ifngr1 to activate Stat1 signaling in target cells. *Molecular cell*. 2014;56(2):193-204.
67. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, et al. Antiinflammatory Therapy with Canakinumab for Atherosclerotic Disease. *The New England journal of medicine*. 2017.
68. Hilgendorf I, Eisele S, Remer I, Schmitz J, Zeschky K, Colberg C, et al. The oral spleen tyrosine kinase inhibitor fostamatinib attenuates inflammation and atherogenesis in low-density lipoprotein receptor-deficient mice. *Arteriosclerosis, thrombosis, and vascular biology*. 2011;31(9):1991-9.

## 9. Figure Legends

### Figure 1. CaP particles stimulate IL-1 $\beta$ release.

**A.** ELISA measurement of IL-1 $\beta$  release from 6 different human VSMC isolates (indicated by 6 different symbols) over 16 hours with or without CaP particles (12.5  $\mu$ g/ml) and with or without a 24-hour pre-treatment with LPS (0.1  $\mu$ g/ml). The control contained equivalent PBS. Values with means are displayed and one-way ANOVA followed by Holm-Sidak's multiple comparisons test was used to compare groups with the control. Significant differences are indicated as \* ( $P < 0.05$ ) or \*\* ( $P < 0.001$ ). **B.** IL-1 $\beta$  release from 6 different human VSMC isolates exposed to CaP particles (12.5  $\mu$ g/ml) at time points up to 24 hours or LPS (0.1  $\mu$ g/ml) for 24 hours. Values with means are displayed and were analysed by one-way ANOVA followed Holm-Sidak's multiple comparisons test; significant differences from the no additions '0' control are indicated as \* ( $P < 0.01$ ). **C.** Western analysis of pro-IL-1 $\beta$  levels in VSMC lysates treated as in B. In C, blots are representative of 4 independent experiments in cell isolates from different individuals. Results from 3 further VSMC isolates are displayed in the Supplement Figure S1B.

### Figure 2. IL-1 $\beta$ release at different levels at different stages of VSMC culture.

**A.** ELISA measurement of IL-1 $\beta$  release from early passage (3-5), mid-passage (6-13) or senescent VSMCs (14-18) with or without CaP particles (12.5  $\mu$ g/ml) incubated over 16 hours. Data are displayed on a log-scale and mean levels for each treatment are indicated. Data were log transformed prior to statistical analysis. A two-way ANOVA followed by a Holm-Sidak's multiple comparisons test was used to compare groups. Significant differences are indicated by \* ( $P < 0.01$ ) or \*\* ( $P < 0.001$ ). 4 different VSMC isolates were used for each cell culture stage. **B.**  $\beta$ -galactosidase staining of proliferating (mid-passage) or senescent VSMCs, showing higher expression in senescent cells. **C.** Western analysis of pro-IL-1 $\beta$  levels in proliferating or senescent VSMC lysates treated with CaP particles (12.5  $\mu$ g/ml) for indicated times. Blots showing results from 2 different cell isolates are shown in the Supplement Figure S1C and S1D.

### Figure 3. Inflammasome activators MSU, nigericin (N) and ATP do not induce IL-1 $\beta$ release from human VSMCs.

ELISA measurement of IL-1 $\beta$  release from 5 different VSMCs isolates (indicated by 5 different symbols), displayed as fold-change related to '0' control values for each cell isolate. Cells were treated with either CaP particles (12.5  $\mu$ g/ml), MSU crystals (12.5  $\mu$ g/ml), nigericin (25  $\mu$ M), ATP (10 mM), CaCl<sub>2</sub> (5.4 mM, 'Ca'), Na<sub>2</sub>HPO<sub>4</sub> (2 mM, 'P'), vehicle control (DMSO) or no additions (0) for 16 hours. Mean levels for each treatment are indicated. Raw data are displayed in Supplement Figure S4. A one-way ANOVA on log-transformed raw data followed by a Holm-Sidak's multiple comparisons test determined that CaP treatment significantly differed from each of the other treatments (\* $P < 0.05$ ).

#### Figure 4. CaP particles stimulate caspase-1 activity

**A.** ELISA measurement of IL-1 $\beta$  release from 3 different VSMC isolates over 16 hours with or without CaP particles (12.5  $\mu$ g/ml) and with or without YVAD (20  $\mu$ M). A two-way ANOVA followed by Holm-Sidak's multiple comparisons test was used to determine differences between means. Results are presented as mean  $\pm$  SEM, \* $P$ <0.05. **B.** VSMCs were treated with or without CaP particles at indicated times then labelled with FAM-YVAD-FLICA (active caspase-1 reagent), PI and Hoechst and imaged live. Arrows indicate foci/cells containing active caspase-1. Images are representative of 5 experiments in different cell isolates, quantitation of active caspase-1 and PI in live cells is displayed in Figure 6D and Supplement Figure S8 and S9. **C** and **D.** PI uptake (C) or IL-1 $\beta$  release (D) from 3 different VSMCs isolates in serum-free conditions (SFM) or in basal medium containing 5% FBS (BM) with or without CaP particles (12.5  $\mu$ g/ml) over 16 hours. Mean values  $\pm$  SEM are displayed. In C, a two-way ANOVA followed by Holm-Sidak's multiple comparisons test was performed and significant differences between corresponding no CaP and CaP-treated samples are indicated by \* ( $P$ <0.05). In D, a two-way ANOVA was performed on log-transformed data followed by Holm-Sidak's multiple comparison's test and significant differences between corresponding no CaP and CaP-treated samples are indicated by \* ( $P$ <0.05).

#### Figure 5. Caspase-1 activity colocalises with IL-1 $\beta$

VSMCs were treated with or without CaP particles (12.5  $\mu$ g/ml) for indicated times, labelled with FAM-YVAD-FLICA (active caspase-1 reagent), then fixed and incubated with antibodies recognising IL-1 $\beta$ . Arrows indicate areas in merged images where active caspase-1 and IL-1 $\beta$  overlap. Arrowheads indicate some cells treated with CaP particles that did not contain IL-1 $\beta$ . Confocal images are representative of experiments in 3 different cell isolates.

#### Figure 6. CaP uptake and SYK phosphorylation

**A.** ELISA measurement of IL-1 $\beta$  release from VSMCs that were exposed to chlorpromazine (1  $\mu$ g/ml), wortmannin (100 ng/ml), nystatin (10  $\mu$ g/ml), Y27632 (10  $\mu$ mol/L) or vehicle control (DMSO) for 1 hour, then exposed to CaP particles (12.5  $\mu$ g/ml) for 16 hours. A two-way ANOVA followed by Sidak's multiple comparisons test was used to compare groups. Significant differences are indicated by \* ( $P$ <0.01),  $n$ =3, i.e. VSMC isolates from 3 different individuals (indicated by 3 different symbols). **B.** VSMCs were treated with or without CaP particles (12.5  $\mu$ g/ml) for 10 minutes and analysed for SYK or phosphorylated SYK (525/526) by Western analysis. Western blots shown are representative of experiments from 3 different cell isolates (see Supplement Figure S3.) Ratios of SYK-P to loading control were higher in CaP-treated cells, compared with non-treated controls (paired t-test of log-transformed data,  $P$ <0.01,  $n$ =3 different isolates). **C.** ELISA measurement of IL-1 $\beta$  release from 5 different VSMC isolates exposed to CaP particles (12.5  $\mu$ g/ml) and R406 at concentrations indicated or vehicle control (DMSO). One-way ANOVA of log-transformed data was performed followed by Holm-Sidak's multiple comparisons tests; indicated treatments differed significantly from control values, \*  $P$ <0.05. **D.** Quantitation of active caspase-1 in live VSMCs after treatment with CaP particles (12.5  $\mu$ g/ml) for 0, 2 or 16 hours with either a 1 hour pre-treatment with R406 (1  $\mu$ M) or equivalent DMSO. Results shown are representative of 4 independent experiments in different cell isolates. Scatter plots display fluorescence levels of the caspase-1 substrate (FAM-YVAD-FLICA) for approximately 10000 live VSMCs for each treatment group. Each dot represents the active caspase-1 level for 1 cell. As indicated, when comparing VSMCs from 4 different individuals, CaP particles increased active caspase-1 activity and this effect was reduced in the presence of R406, when comparing the top 5<sup>th</sup> percentile of VSMCs at the 16 hour time point ( $P$ =0.0215).

**Figure 7. Release of IL-1 $\beta$  via exosomes**

A. VSMCs were treated with or without CaP particles (12.5  $\mu\text{g}/\text{ml}$ ) for 16 hours with or without a pre-treatment with spiroepoxide (10  $\mu\text{M}$ ). A two-way ANOVA followed by Sidak's multiple comparisons test was used to compare groups. Significant differences are indicated by \* ( $P < 0.05$ ),  $n = 3$ , i.e. VSMC isolates from 3 different individuals (indicated by 3 different symbols).

B. Exosome secretion by VSMCs. VSMCs were incubated in 2.5%FBS/M199 media with or without CaP particles (12.5  $\mu\text{g}/\text{ml}$ ) and with or without R406 (1  $\mu\text{M}$ ) for 16 hours. Conditioned media were harvested and exosomes were quantified using an anti-CD63 bead-capturing assay which detects CD63/CD81 exosomes. Statistical significance was tested by one-way ANOVA followed by Sidak's multiple comparisons test. Significant differences are indicated by \*\* ( $P < 0.01$ ), \*\*\* ( $P < 0.001$ ),  $n = 10$ .

C. Western analysis of exosomes isolated by differential ultracentrifugation from conditioned media of VSMCs treated with or without CaP particles for 16 hours. Exo, exosomes; VSMC, whole cell lysates. The membrane was probed with IL-1 $\beta$  antibodies (R and D), CD63 (BD Pharmingen), with vinculin [38] and Coomassie brilliant blue staining to demonstrate protein loading.




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5




Figure 6





Figure 7.

## Graphical Abstract



**Highlights**

- CaP particles induce IL-1 $\beta$  release from human VSMCs
- Senescent cells display higher basal and CaP-stimulated IL-1 $\beta$  release
- Inflammasome agonists ATP, nigericin and MSU crystals do not induce IL-1 $\beta$  release from human VSMCs
- CaP particle-induced IL-1 $\beta$  release is dependent on SYK, caspase-1 and exosome release