



King's Research Portal

DOI:

[10.1016/j.schres.2017.12.011](https://doi.org/10.1016/j.schres.2017.12.011)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Quednow, B. B., Ejebe, K., Wagner, M., Giakoumaki, S. G., Bitsios, P., Kumari, V., & Roussos, P. (2017). Meta-analysis on the association between genetic polymorphisms and prepulse inhibition of the acoustic startle response. *Schizophrenia Research*. <https://doi.org/10.1016/j.schres.2017.12.011>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Meta-analysis on the association between genetic polymorphisms and prepulse inhibition of the acoustic startle response

Boris B. Quednow^{1,2,*}, Kenechi Ejebe³, Michael Wagner⁴, Stella G. Giakoumaki⁵, Panos Bitsios⁶, Veena Kumari⁷, Panos Roussos^{3,8,9,*}

¹Experimental and Clinical Pharmacopsychology, Psychiatric Hospital, University of Zurich, Switzerland

²Neuroscience Center Zurich, Research Group Disorders of the Nervous System, University and ETH Zurich, Zurich, Switzerland

³Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, USA

⁴Department of Psychiatry and Psychotherapy, University of Bonn, Germany

⁵Department of Psychology, Gallos University campus, University of Crete, Rethymno, Greece

⁶Department of Psychiatry and Behavioral Sciences, Faculty of Medicine, Voutes University campus, University of Crete, Heraklion, Greece

⁷Department of Psychology, Institute of Psychiatry, King's College London, United Kingdom

⁸Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, USA

⁹Mental Illness Research, Education, and Clinical Center (VISN 2), James J. Peters VA Medical Center, New York, USA

Abbreviated Running Title:

Meta-analysis on genetics of sensorimotor gating

Manuscript Category:

Original Article

Revision resubmitted:

December 13th, 2017

Manuscript Characteristics:

Number of words in the abstract: 234

Number of words in the text: 4458

Number of references: 114

Number of tables: 2

Number of figures: 0

*Corresponding authors:

Boris B. Quednow, Ph.D., Dipl.-Psych.

Experimental and Clinical Pharmacopsychology

Psychiatric Hospital of the University of Zurich

Lenggstrasse 31

CH-8032 Zurich, Switzerland

Tel.: 0041-44-384-2777

Fax: 0041-44-384-2499

E-Mail: quednow@bli.uzh.ch

Panos Roussos, M.D., Ph.D.

Department of Psychiatry and Department of Genetics and

Genomic Science and Institute for Multiscale Biology

Icahn School of Medicine at Mount Sinai

1470 Madison Ave,

New York, NY, 10029, USA

Tel.: 001-212-824-8982

Fax: 001-646-537-9583

E-Mail: Panagiotis.roussos@mssm.edu

Abstract

Sensorimotor gating measured by prepulse inhibition (PPI) of the acoustic startle response (ASR) has been proposed as one of the most promising electrophysiological endophenotypes of schizophrenia. During the past decade, a number of publications have reported significant associations between genetic polymorphisms and PPI in samples of schizophrenia patients and healthy volunteers. However, an overall evaluation of the robustness of these results has not been published so far. Therefore, we performed the first meta-analysis of published and unpublished associations between gene polymorphisms and PPI of ASR. Unpublished associations between genetic polymorphisms and PPI were derived from three independent samples. In total, 120 single observations from 16 independent samples with 2,660 study participants and 43 polymorphisms were included. After correction for multiple testing based on false discovery rate and considering the number of analyzed polymorphisms, significant associations were shown for four variants, even though none of these associations survived a genome-wide correction ($P < 5 \times 10^{-8}$). These results imply that PPI might be modulated by four genotypes – *COMT* rs4680 (primarily in males), *GRIK3* rs1027599, *TCF4* rs9960767, and *PRODH* rs385440 – indicating a role of these gene variations in the development of early information processing deficits in schizophrenia. However, the overall impact of single genes on PPI is still rather small suggesting that PPI is – like the disease phenotype – highly polygenic. Future genome-wide analyses studies with large sample sizes will enhance our understanding on the genetic architecture of PPI.

Keywords

Prepulse inhibition; startle; sensorimotor gating; polymorphism; SNP; genotype; gene; mutation; schizophrenia; psychosis; endophenotype; intermediate phenotype; meta-analysis

Introduction

Prepulse inhibition (PPI) of the acoustic startle response (ASR) is defined as a substantial reduction of the startle amplitude that occurs when a startling stimulus is preceded within a timeframe of 20-500 ms by a stimulus of lower intensity than the startling stimulus (Graham, 1975). PPI has been shown to occur across species ranging from mollusks and fishes to mammals including non-human and human primates (Burgess and Granato, 2007; Frost et al., 2003; Hoffman and Searle, 1965; Ison and Leonhard, 1970; Krauter et al., 1973; Linn and Javitt, 2001). Animal studies carried out predominantly in rodents suggested that PPI is regulated by a cortico-striato-pallido-pontine (CSPP) circuitry including frontal and mediotemporal regions, ventral striatum, ventral pallidum, and pontine regions of the brainstem (Fendt et al., 2001; Swerdlow et al., 2001). Within the CSPP circuit, several neurotransmitters have been demonstrated to play a major role in the mediation of PPI such as dopamine, noradrenaline, serotonin, acetylcholine, glutamate, and γ -aminobutyric acid (GABA) (Geyer et al., 2001; Koch, 1999). Consequently, PPI has been proposed as a “window” into brain chemistry potentially allowing the identification of neuropharmacological alterations in specific psychiatric disorders with PPI abnormalities (Braff, 2010). In fact, lower PPI levels have been reported for several neuropsychiatric disorders (Braff et al., 2001; Kohl et al., 2013; Quednow, 2008), but were replicated best for schizophrenia spectrum disorders (e.g., Braff et al., 1992; Cadenhead et al., 1993; Kumari et al., 2000; Ludewig et al., 2003; Parwani et al., 2000; Quednow et al., 2006; Swerdlow et al., 2014). Given that PPI shows such a robust association with schizophrenia and because it is heritable (Anokhin et al., 2003; Greenwood et al., 2007; 2016; Willott et al., 1994; 2003), reduced in unaffected relatives of schizophrenia patients (Cadenhead et al., 2000; Kumari et al., 2005), and already decreased in early (prodromal) stages of the disease (Quednow et al., 2008a; Ziermans et al., 2011; 2012), PPI was suggested as an promising candidate of an intermediate or endophenotypic marker in genetic studies of schizophrenia (Braff and Light, 2005; Gottesman and Gould, 2003). Specifically, the substantial heritability of PPI in humans – ranging from 29% to 50% across a number of studies – suggests PPI as a favorable target for genetic analyses (Anokhin et al., 2003; Greenwood et al., 2007; Hasenkamp et al., 2010; Seidman et al., 2015). Additionally, it was recently shown that PPI revealed substantially increased heritability (47%) in 97 families multiply affected by schizophrenia

when compared with a 96 families, in which only a single individual was affected (2%). This finding further promotes the assumption that a commonality of genes underlies both schizophrenia and PPI (Greenwood et al., 2016).

The endophenotype concept assumes that an endophenotypic marker is a heritable, quantifiable, and stable trait, which is determined by a smaller number of genes compared to the respective complex disease phenotype (Braff et al., 2007). Accordingly, in the last decade several research groups aimed to identify gene effects on the expression of PPI. The first positive findings were published in 2008, where associations of PPI with single nucleotide polymorphisms (SNPs) of the neuregulin-1 (*NRG1*, rs3924999) (Hong et al., 2008), catechol-O-methyltransferase (*COMT*, rs4680) (Roussos et al., 2008b), dopamine D3 receptor (*DRD3*, rs6280) (Roussos et al., 2008a), and the serotonin-2A receptor (*5-HT2AR*, rs6311/6313) gene (Quednow et al., 2008b) have been reported from hypothesis-driven single association studies. Since then a number of further single association studies reported significant associations of PPI with numerous SNPs, which mainly have been identified as schizophrenia risk genes previously (for a list of studies and SNPs see **Table 1** and **2**, respectively). The only SNPs that have been replicated in at least two independent samples so far are *CHRNA3* (rs1051730) (Petrovsky et al., 2010), *5-HT2AR* (rs6311/6313) (Quednow et al., 2008b; 2009), *COMT* (rs4680) (Liu et al., 2013; Quednow et al., 2009; 2010; Roussos et al., 2008b), *NRG1* (rs3924999) (Hong et al., 2008), *TCF4* (rs9960767) (Quednow et al., 2011), and *DRD2* (rs1800497) (Volter et al., 2012). However, for most of these SNPs also negative findings have been reported. Most recently, the first explorative genome-wide association study (GWAS) identified two non-coding loci (rs61810702 and rs4718984) that were co-localized with expression quantitative trait loci related with the gene expression of nerve growth factor (*NGF*) and calneuron 1 (*CALN1*) genes. Additionally, a higher polygenic risk score for schizophrenia was associated with lower PPI (Roussos et al., 2016).

The heterogeneity of these genetic results implies i) that there are probably no single genes with a high impact on PPI and ii) that there are likely also some false positive results considering that – from today’s perspective – many of the previous studies are strongly underpowered and lack replication samples (Button et al., 2013). One way to reduce the number of false positive (but also false negative) results in psychiatric genetics is the application of a meta-analysis, in which all

available data are included (Levinson, 2005; Lohmueller et al., 2003). Therefore, we performed a systematic meta-analysis of all genotype-SNP associations published so far using a weighted Z-method approach (Stouffer et al., 1949). We additionally included three independent data sets coming from samples that have been published before for reporting of genotype-PPI associations but for which also yet unpublished genotype or GWAS data existed (Petrovsky et al., 2010; Quednow et al., 2008b; 2009; 2010; 2011; Roussos et al., 2016). Genetic variants were included into the analysis if they were available in at least two independent samples. In order to control for multiple comparisons, a false discovery rate (FDR) method based on estimation of tail area-based FDR was applied (Strimmer, 2008). The aim of this systematic meta-analysis was the identification of the most robust i.e., significant genotype-PPI associations.

Methods

Eligibility criteria

Human association studies (single associations and GWAS) with genetic variants reporting p-values and direction of effect for PPI-genotype associations in samples of healthy controls or patients with schizophrenia spectrum disorders. To be included, a gene variant must have been available in at least two independent samples.

Information sources and search strategy

With the search term (("prepulse inhibition" OR "sensorimotor gating") AND (mutation OR polymorphism OR polymorphisms OR snp OR snps OR gene OR genotype) NOT (rats OR rat OR mice OR mouse)) NOT review[ptyp]), 63 articles have been identified initially by a MEDLINE search (PubMed.gov). Only studies reporting p-values of genetic effects on PPI in healthy volunteers and patients with schizophrenia spectrum disorders were included. Studies investigating PPI-gene associations in pregnant women or individuals with developmental disorders (e.g., with 22q11 syndrome) were excluded. With this procedure, we identified 19 original articles reporting SNP-PPI associations (**Table 1**). After checking, if at least two p-values of a specific genotype-PPI and from independent samples are available (e.g., in two publications or in a publication and an unpublished data set), 120 single observations from 16 independent samples with 2,660 study participants and 43 polymorphisms were included in the meta-analysis (**Table 2**).

Inclusion of unpublished data

The authors included unpublished genotype-PPI association data from three samples that have been published before: (1) the LOGOS sample recruited in Crete, Greece, consisting of healthy volunteers (max. n=686) (Roussos et al., 2016), (2) a sample of healthy volunteers (max. n=100) from London, UK (Petrovsky et al., 2010; Quednow et al., 2009; 2011), and (3) a sample of patients with schizophrenia spectrum disorders (max. n=107) from Bonn, Germany (Petrovsky et al., 2010; Quednow et al., 2008b; 2010; 2011). P-values of yet unpublished genotype-PPI associations have been calculated using the same statistical approach as previously reported, e.g., using the same covariates

such as sex, smoking status, and antipsychotic medication (London and Bonn samples: Quednow et al., 2011). The final studies, cohorts and number of samples per study included in the meta-analysis is shown in **Table 1** and **2**.

Statistical Analysis

We sum the association evidence across studies using a weighted Z -method approach where every study is weighted by the effective sample size (Stouffer et al., 1949). Briefly, in each study, we convert P_i P -values into Z_i signed Z -scores based on the Δ_i direction of effect using the following equation:

$$Z_i = \phi^{-1} \left(\frac{P_i}{2} \right) * \text{sign}(\Delta_i).$$

We then estimate overall Z score by summing Z -scores for each genetic polymorphism using weights proportional to the square-root of the sample size for each study:

$$Z = \frac{\sum_i Z_i w_i}{\sqrt{\sum_i w_i^2}},$$

where $w_i = \sqrt{N_i}$ and N_i is the sample size of study i . Finally, the overall Z score is converted back to overall P -value for a given genotype based on:

$$P = 2 \Phi(|-Z|).$$

For multiple testing correction, we applied FDR at combined P -values estimated based on `fdrTool` in R (Strimmer, 2008). The Cohen's d effect sizes for the strengths of association were estimated from Z -values according to Rosenthal (1984). For estimation of effect sizes in schizophrenia GWAS study we used the average odds ratio of 1.1 across all genome-wide significant SNPs and estimated Cohen's d based on:

$$d = \text{LogOddsRatio} \times \frac{\sqrt{3}}{\pi}$$

As sex effects are well-documented for the rs4680 *COMT* SNP (Tunbridge and Harrison, 2011), and as strong sex differences in rs4680 *COMT* effects on PPI were recently reported in a non-trivial sample of healthy subjects (Swerdlow et al., 2017), we additionally analyzed sex effects for this SNP only. Based on the existing studies, we are well powered to study the *COMT* association in males only as the majority of the included samples with known gender (92% or 870 out of 943) were males.

Results

In this study, we examined the associations of SNPs and PPI of ASR. We evaluated a total of 43 SNP-PPI associations, while for each SNP two to six samples were available (mean 2.8, standard deviation 1.1, median 2.0 samples). Statistical significance was defined as an FDR-adjusted p-value of 0.05 or smaller. At this threshold, four index SNPs were significant. The strongest association was between *COMT* rs4680 and PPI ($P = 5.5 \times 10^{-5}$, FDR = 0.002, Cohen's $d = 0.28$) in male samples only (**Table 2**). The other three loci with significant association with PPI were: *GRIK3* rs1027599 ($P = 0.004$, FDR = 0.045, Cohen's $d = 0.19$), *TCF4* rs9960767 ($P = 0.006$, FDR = 0.050, Cohen's $d = 0.19$) and *PRODH* rs385440 ($P = 0.006$, FDR = 0.050, Cohen's $d = 0.27$). As expected, the detected Cohen's d effect sizes are small according to the definition of Cohen (1988) and none of these associations survived a genome-wide correction ($P < 5 \times 10^{-8}$). Notably, we also found a significant effect for *COMT* rs4680 in both males and females, but with smaller effect size that did not survive multiple testing corrections ($P = 0.01$, FDR = 0.057, Cohen's $d = 0.15$), indicating a putative gender-specific effect of *COMT* on PPI.

Discussion

The aim of this meta-analysis was the identification of robust associations between gene polymorphisms and PPI of the ASR. Given that PPI was repeatedly proposed as an endophenotype of schizophrenia (Braff et al., 2007), this investigation might elucidate which genes may robustly contribute to the low PPI levels of schizophrenia patients and presumably to the disease itself. We thus analyzed published and unpublished data resulting in the inclusion of 120 single observations from 16 independent samples with 2,660 study participants and 43 polymorphisms. After multiple testing corrections based on an FDR approach, four polymorphisms were significantly associated with PPI. In males, the *COMT* rs4680 (Val158Met) polymorphism showed the strongest association with PPI ($P = 5.5 \times 10^{-5}$, FDR = 0.002, Cohen's $d = 0.28$), while the *COMT* gene effect was barely not significant ($P = 0.01$, FDR = 0.057, Cohen's $d = 0.15$) in the mixed-gender sample. Carriers of the *COMT* rs4680 G (valine) allele showed lower PPI levels than carriers of the A (methionine) allele. Moreover, *GRIK3* rs1027599 ($P = 0.004$, FDR = 0.045, Cohen's $d = 0.19$), *TCF4* rs9960767 ($P = 0.006$, FDR = 0.050, Cohen's $d = 0.19$), and *PRODH* rs385440 ($P = 0.006$, FDR = 0.050, Cohen's $d = 0.27$) showed significant association with PPI. Regarding *GRIK3* rs1027599, the T allele carriers showed lower PPI, while the same was true for *TCF4* rs9960767 C and *PRODH* rs385440 A allele carriers. In fact, when the widely accepted threshold for genome-wide multiple corrections ($P < 5 \times 10^{-8}$) is applied, none of the results is powerful enough to survive. However, it has to be taken into account that the power of this meta-analysis (sample size range: $N=436$ to 1224) was much lower, e.g., compared to previously published mega-analyses in schizophrenia genetics, including ten thousands of participants (Ripke et al., 2013; Steinberg et al., 2011). Nevertheless, the effect sizes of the significant SNPs in the present meta-analysis are small (Cohen's $d = 0.19$ to 0.28) in the sense of Cohen's definition (Cohen, 1988) but relatively strong considering the strengths of genetic associations commonly shown in meta-analysis of GWAS in schizophrenia (Cohen's $d = 0.05$ based on Schizophrenia Working Group of the Psychiatric Genomics, 2014).

The *COMT* gene effect on PPI is highly plausible as the *COMT* enzyme is involved in the metabolic inactivation of dopamine and norepinephrine in regions with a low density of dopamine and

noradrenaline transporters specifically in the frontal cortex (Rivett et al., 1982). The rs4680 polymorphism leads to an amino acid substitution of methionine for valine changing the metabolic rate of the enzyme. Consequently, compared to methionine homozygotes, valine homozygotes have a 3 to 4-fold stronger COMT enzyme activity and, thus, an increased catabolism of catecholamines in the frontal cortex (Lachman et al., 1996). Animal and human studies suggest that PPI is critically modulated by dopamine and noradrenaline neurotransmission at several stages of the CSPP circuit (Geyer et al., 2001; Swerdlow et al., 2001). Moreover, *COMT* messenger RNA is highly expressed in the prefrontal cortex and the hippocampus and less in the striatum, the ventral tegmental area, or the substantia nigra (Tunbridge et al., 2006). Considering that these structures are participating in the CSPP circuit, it is likely that the *COMT* polymorphism influences PPI at the prefrontal or hippocampal level (Swerdlow et al. 2001; Quednow et al, 2010). Specifically, lower prefrontal dopamine concentrations probably contribute to reduced PPI levels in valine allele carriers as is was shown that reduced dopamine activity in the prefrontal cortex goes along with a disruption of PPI (Bubser and Koch, 1994; Ellenbroek et al., 1996; Zavitsanou et al., 1999). In addition, drug-induced inhibition of the COMT enzyme with tolcapone leads to PPI enhancement only in the *COMT* Val carriers (Bitsios and Roussos, 2011; Giakoumaki et al., 2008; Roussos et al., 2009b). In line with these human data, it was shown that the effects of *COMT* inhibition by tolcapone on amphetamine-modified PPI were categorically different in rat strains exhibiting low vs. high levels *Comt* expression in the forebrain (Swerdlow et al., 2013). Moreover, genetic associations between *COMT* and psychiatric phenotypes frequently display sex differences (Tunbridge and Harrison, 2011). Accordingly, *COMT* rs4680 genotype effects on PPI have been demonstrated particularly in males, while the gene effects were absent, less strong, or even reversed in females (Montag et al., 2008; Quednow et al., 2009; 2010; Roussos et al., 2008b; Swerdlow et al., 2017). Finally, *COMT* rs4680 has been proposed to be a risk gene for schizophrenia for a long time but recent meta-analytical results are conflicting (Gonzalez-Castro et al., 2016; Taylor, 2017). Again, associations between *COMT* rs4680 and schizophrenia or schizotypy have been demonstrated primarily in males but not females (for reviews see Tunbridge and Harrison, 2011). Thus, the sex-specific effect of the *COMT* rs4680 on PPI shown here is in agreement with a variety of previous findings.

The *GRIK3* gene encodes the glutamate ionotropic receptor kainate type subunit 3, which is one of three principal subunits (GRIK1-3) of the tetrameric kainate/AMPA receptors. Two further auxiliary subunits of these receptors exist (GRIK4-5) (Fernandez et al 2009, Neuron). GRIK3-containing receptors differ from other kainate/AMPA receptors as they can only be activated by fast and strong glutamate releases. Moreover, they have some special electrophysiological properties suggesting that GRIK3-containing receptors have specialized presynaptic functions (Perrais et al., 2009). Decreased expression of GRIK3 in the prefrontal cortex and hippocampus of schizophrenia patients have been shown (Hu et al., 2015): While there are no published positive associations of the *GRIK3* rs1027599 SNP with psychiatric disorders so far, other *GRIK3* SNPs have been associated with schizophrenia (Begni et al., 2002; Kilic et al., 2010) and major depression (Schiffer and Heinemann, 2007), previously. Moreover, sporadic deletion in chromosome 1p34.3 involving *GRIK3* was reported for a girl with developmental delay (Takenouchi et al., 2014). Interestingly, in a recent twin study, a *GRIK3* gene set was associated with overall startle responses (Vaidyanathan et al., 2014), raising the question if the association with PPI shown here is rather an epiphenomenon of changes in startle reactivity (Csomor et al., 2008). Moreover, the kainate antagonist LY382884 reduced both exaggerated ASR and PPI deficits observed in mice with genetically reduced NMDA receptor expression (Duncan et al., 2010). However, these results might be explained by the changes in ASR again, as strongly elevated startle responses can result in low PPI levels also in mice (Csomor et al., 2008).

TCF4 belongs to the superfamily of basic Helix-Loop-Helix (bHLH) transcription factors that can act as a transcriptional repressor or activator in a context specific fashion. It can be considered as an integrator ('hub') of several bHLH networks controlling critical steps of various developmental and plasticity related transcriptional programs in neurons (Quednow et al., 2014). Haploinsufficiency of the *TCF4* gene causes the Pitt-Hopkins syndrome – a severe neurodevelopmental disease characterized by mental retardation, microcephaly, epilepsy, facial dysmorphisms, and intermittent hyperventilation –, suggesting that *TCF4* is critical for the development of the mammalian nervous system (de Pontual

et al., 2009; Zweier et al., 2007). Three large but also partially overlapping meta-analyses of GWAS consistently identified that common *TCF4* polymorphisms are associated with the risk of schizophrenia (Schizophrenia Psychiatric Genome-Wide Association Study, 2011; Stefansson et al., 2009; Steinberg et al., 2011). In these analyses, two SNPs at the intron located between the internal exon 4 and internal exon 5 of human *TCF4* gene on chromosome 18q21.2 (rs9960767, rs17512836) and an intragenic SNP near the *TCF4* gene (rs4309482) have shown the strongest association with the disease (Schizophrenia Psychiatric Genome-Wide Association Study, 2011; Stefansson et al., 2009; Steinberg et al., 2011). Several subsequent studies replicated schizophrenia-*TCF4* gene associations in independent samples: (1) Two studies in Han Chinese (in which the rs9960767 SNP is not polymorphic) identified *TCF4* rs2958182 (Li et al., 2010) as well as rs9320010, rs7235757, and rs1452787 (Li et al., 2016) to be associated with schizophrenia. (2) In a discovery sample from Ireland and a replication sample including non-overlapping samples from the Psychiatric GWAS Consortium (PGC), two intronic *TCF4* SNPs (again rs9960767 and rs17594526) passed the genome-wide significance threshold of $p < 5 \times 10^{-8}$ (Irish Schizophrenia Genomics and the Wellcome Trust Case Control, 2012). (3) In a recent family-based linkage meta-analysis a further *TCF4* SNP was identified (rs1261117) as linked to schizophrenia (Aberg et al., 2013). These results have contributed to the view that *TCF4* SNPs belong to the best-replicated schizophrenia susceptibility genes. A potential disturbance of *TCF4* function in schizophrenia would be in line with the assumption that schizophrenia is a neurodevelopmental disorder (Quednow et al., 2014); however, the exact functional impact of the discussed *TCF4* SNPs has still to be elucidated. Finally, transgenic mice moderately overexpressing *Tcf4* in the brain as well as *Tcf4* haploinsufficient mice both display profound reductions in PPI (Brzozka et al., 2010; Kennedy et al., 2016). Beyond PPI, *TCF4* SNPS including rs9960767 have been shown to be associated with another proposed endophenotype of schizophrenia: P50 suppression also called “sensory gating”; however, this association was only present in smoking individuals but not in never-smokers (Quednow et al., 2012). Taken together, the significant association of *TCF4* rs9960767 and PPI in human cohorts reported here is neurobiologically and pathophysiologically plausible as well.

PRODH encodes the enzyme proline oxidase, which is involved, among other functions, in the conversion of proline to D-1-pyrroline-5-carboxylate (P5C) in mitochondria. P5C is subsequently converted to glutamate or GABA, two neurotransmitters critically implicated in the pathophysiology of schizophrenia (Roussos et al., 2009a). Together with *COMT*, *PRODH* is located at 22q11 and, thus, both genes belong to the ~30 genes affected by the 22q11 deletion syndrome (Velo-Cardio-Facial or DiGeorge syndrome). Individuals with a 22q11 deletion syndrome show a strongly elevated risk for schizophrenia symptoms (Prasad et al., 2008) as well as reductions of PPI (Sobin et al., 2005a, b). In addition, rodent models of the 22q11 syndrome are associated with deficits in PPI (Diamantopoulou et al., 2017; Didriksen et al., 2017; Gogos et al., 1999). Previous studies suggested a possible role of the *PRODH* gene variations, in the pathogenesis of schizophrenia. Specifically, *PRODH* haplotypes consisting of rs372055 (1945T>C), rs450046 (1766A>G), and rs385440 (1852G>A) SNPs were significantly associated with schizophrenia, indicating that the alleles 1945C, 1766G, and 1852A are overtransmitted in schizophrenia patients (Li et al., 2004; Liu et al., 2002). The haplotypes, which included rs385440, is associated with *PRODH* hyperactivity potentially resulting in reduced proline levels and increased P5C/Glu availability in the central nervous system (Phang et al., 2008) and these changes might induce changes in the modulation of PPI (Roussos et al., 2009a). As both *COMT* and *PRODH* SNPs seem to have an impact on PPI and given that these genes are within 22q11 locus, further studies should investigate gene-gene interactions of both variants with regard to PPI.

Limitations of our meta-analysis include the small number of included studies and sample sizes making the estimates of effect sizes less reliable. Moreover, for most of the SNPs only two independent samples were available. It should be noticed, that the *PRODH* effect on PPI was indeed similarly strong as the *COMT* effect in males but that the *PRODH* sample size was nevertheless the lowest amongst all analyzed SNPs. Thus, the *PRODH* effect is likely less reliable than the *COMT* effect. Moreover, similar as for *PRODH*, the *GRIK* effect was calculated from only two studies, while three studies were included for *TCF4* and four regarding *COMT* (in males, and six for *COMT* including females). Consequently, the *GRIK* effects might also be less robust compared to *TCF4* and *COMT* (in males). Finally, we used a weighted Z-method approach using reported p-values to

calculate the meta-analysis and consequently, statistical differences between studies (e.g., regarding the inclusion of different sets of covariates) have not been considered. Accordingly, as we did not use the original data sets of previously published work we were not able to adjust the results for the common confounding factors affecting PPI, such as gender and menstrual cycle, smoking, startle magnitude, as well as medication (Quednow, 2008). Particularly in patient samples, PPI-gene associations could have been masked by the potential PPI-enhancing impact of antipsychotic medication (Quednow et al., 2006; 2008a). However, several previous studies have controlled for such confounding factors and reported adjusted p-values, but not all. The additional analyses of so far unpublished data from patients with schizophrenia spectrum disorders reported in the present paper in fact considered smoking, sex, and antipsychotic medication status as covariates and only the adjusted p-values were used for the meta-analysis. Nevertheless, the overall strength of the gene effects might have been biased by confounding factors such as antipsychotic medication. Another limitation is that the included studies employed a variety of different PPI setups. Given that it has not been investigated systematically yet, which PPI parameters (e.g., prepulse and pulse intensities, stimulus-onset asynchrony, number of stimuli, intertrial intervals) might provide an optimal signal-to-noise ratio with regard to gene effects (or if genes might specifically influence PPI at various conditions), it remains unclear how the heterogeneity of PPI setups used in the included studies might have affected our results.

Beyond PPI, several further electrophysiological “gating” or filtering” paradigms have been proposed as potential endophenotypes and clinical biomarkers of schizophrenia, such as P50 suppression, P300 and P3a, and mismatch negativity (MMN) (Light and Swerdlow, 2015). Many studies have investigated the associations between polymorphisms and these electroencephalographic measures but no meta-analysis has been performed for any of these markers yet (Owens et al., 2016). Specifically, the MMN and the P3a event related potential component might be highly promising targets for future endophenotype-related genetic meta-analyses given that both are strongly, robustly, and stably affected in schizophrenia patients and both show considerable heritability (Light and Swerdlow, 2015; Light et al., 2015). However, a recent well-powered study demonstrated significant

difficulty in the identification of genetic associations between 108 schizophrenia risk loci, a polygenic risk score for schizophrenia, and 17 electrophysiological schizophrenia endophenotypes, challenging the endophenotype concept (Liu et al., 2017) – albeit neither PPI, MMN, nor P3a (but P300) have been included in this analysis. Thus, future studies have to reveal if these most promising endophenotype candidates are more useful to decompose the genetic basis of schizophrenia or if they have at least a clinical relevance (Light and Swerdlow, 2015).

Taken together, the present meta-analysis revealed that four SNPs located on genes that have been associated with schizophrenia previously, are robustly correlated with the expression of PPI in humans as well. These associations are neurobiologically plausible as the respective gene products have been shown to be involved in the CSPP circuit processing PPI (Geyer et al., 2001; Koch, 1999; Swerdlow et al., 2001). Although the effect sizes of the gene effects on PPI are relatively strong (at least when compared to the extremely weak gene effects shown in meta-analysis of GWAS in schizophrenia) none of the results would have survived a strict genome-wide correction of the alpha-level. This indicates that PPI – like schizophrenia itself – is highly polygenic. Thus, the question that still has to be answered in the future is whether the hitherto proposed schizophrenia endophenotypes are in fact less polygenic as the complex disease phenotype, so that endophenotypes can be used to predict the risk for schizophrenia or if they can be used to discover new neurobiological treatment targets.

References

- Aberg, K.A., Liu, Y., Bukszar, J., McClay, J.L., Khachane, A.N., Andreassen, O.A., Blackwood, D., Corvin, A., Djurovic, S., Gurling, H., Ophoff, R., Pato, C.N., Pato, M.T., Riley, B., Webb, T., Kendler, K., O'Donovan, M., Craddock, N., Kirov, G., Owen, M., Rujescu, D., St Clair, D., Werge, T., Hultman, C.M., Delisi, L.E., Sullivan, P., van den Oord, E.J., 2013. A comprehensive family-based replication study of schizophrenia genes. *JAMA Psychiatry* 70(6), 573-581.
- Anokhin, A.P., Heath, A.C., Myers, E., Ralano, A., Wood, S., 2003. Genetic influences on prepulse inhibition of startle reflex in humans. *Neurosci Lett* 353(1), 45-48.
- Begni, S., Popoli, M., Moraschi, S., Bignotti, S., Tura, G.B., Gennarelli, M., 2002. Association between the ionotropic glutamate receptor kainate 3 (GRIK3) ser310ala polymorphism and schizophrenia. *Mol Psychiatry* 7(4), 416-418.
- Bitsios, P., Roussos, P., 2011. Tolcapone, COMT polymorphisms and pharmacogenomic treatment of schizophrenia. *Pharmacogenomics* 12(4), 559-566.
- Braff, D.L., Grillon, C., Geyer, M.A., 1992. Gating and habituation of the startle reflex in schizophrenic patients. *Arch Gen Psychiatry* 49(3), 206-215.
- Braff, D.L., Geyer, M.A., Swerdlow, N.R., 2001. Human studies of prepulse inhibition of startle: normal subjects, patient groups, and pharmacological studies. *Psychopharmacology (Berl)* 156(2-3), 234-258.
- Braff, D.L., Light, G.A., 2005. The use of neurophysiological endophenotypes to understand the genetic basis of schizophrenia. *Dialogues Clin Neurosci* 7(2), 125-135.
- Braff, D.L., Freedman, R., Schork, N.J., Gottesman, II, 2007. Deconstructing schizophrenia: an overview of the use of endophenotypes in order to understand a complex disorder. *Schizophr Bull* 33(1), 21-32.
- Braff, D.L., 2010. Prepulse inhibition of the startle reflex: a window on the brain in schizophrenia. *Curr Top Behav Neurosci* 4, 349-371.
- Brauer, D., Strobel, A., Hensch, T., Diers, K., Lesch, K.P., Brocke, B., 2009. Genetic variation of serotonin receptor function affects prepulse inhibition of the startle. *J Neural Transm (Vienna)* 116(5), 607-613.
- Brzozka, M.M., Radyushkin, K., Wichert, S.P., Ehrenreich, H., Rossner, M.J., 2010. Cognitive and sensorimotor gating impairments in transgenic mice overexpressing the schizophrenia susceptibility gene *Tcf4* in the brain. *Biol Psychiatry*, In press.
- Bubser, M., Koch, M., 1994. Prepulse inhibition of the acoustic startle response of rats is reduced by 6-hydroxydopamine lesions of the medial prefrontal cortex. *Psychopharmacology (Berl)* 113(3-4), 487-492.
- Burgess, H.A., Granato, M., 2007. Sensorimotor gating in larval zebrafish. *J Neurosci* 27(18), 4984-4994.
- Button, K.S., Ioannidis, J.P., Mokrysz, C., Nosek, B.A., Flint, J., Robinson, E.S., Munafò, M.R., 2013. Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci* 14(5), 365-376.
- Cadenhead, K.S., Geyer, M.A., Braff, D.L., 1993. Impaired startle prepulse inhibition and habituation in patients with schizotypal personality disorder. *Am J Psychiatry* 150(12), 1862-1867.
- Cadenhead, K.S., Swerdlow, N.R., Shafer, K.M., Diaz, M., Braff, D.L., 2000. Modulation of the startle response and startle laterality in relatives of schizophrenic patients and in subjects with schizotypal personality disorder: evidence of inhibitory deficits. *Am J Psychiatry* 157(10), 1660-1668.
- Cohen, J., 1988. *Statistical Power Analysis for the Behavioral Sciences*, 2nd ed. Lawrence Erlbaum, Hillsdale.
- Csomor, P.A., Yee, B.K., Vollenweider, F.X., Feldon, J., Nicolet, T., Quednow, B.B., 2008. On the influence of baseline startle reactivity on the indexation of prepulse inhibition. *Behav Neurosci* 122(4), 885-900.

- de Pontual, L., Mathieu, Y., Golzio, C., Rio, M., Malan, V., Boddaert, N., Soufflet, C., Picard, C., Durandy, A., Dobbie, A., Heron, D., Isidor, B., Motte, J., Newbury-Ecob, R., Pasquier, L., Tardieu, M., Viot, G., Jaubert, F., Munnich, A., Colleaux, L., Vekemans, M., Etchevers, H., Lyonnet, S., Amiel, J., 2009. Mutational, functional, and expression studies of the TCF4 gene in Pitt-Hopkins syndrome. *Hum Mutat* 30(4), 669-676.
- Diamantopoulou, A., Sun, Z., Mukai, J., Xu, B., Fenelon, K., Karayiorgou, M., Gogos, J.A., 2017. Loss-of-function mutation in *Mirta22/Emc10* rescues specific schizophrenia-related phenotypes in a mouse model of the 22q11.2 deletion. *Proc Natl Acad Sci U S A* 114(30), E6127-E6136.
- Didriksen, M., Fejgin, K., Nilsson, S.R., Birknow, M.R., Grayton, H.M., Larsen, P.H., Lauridsen, J.B., Nielsen, V., Celada, P., Santana, N., Kallunki, P., Christensen, K.V., Werge, T.M., Stensbol, T.B., Egebjerg, J., Gastambide, F., Artigas, F., Bastlund, J.F., Nielsen, J., 2017. Persistent gating deficit and increased sensitivity to NMDA receptor antagonism after puberty in a new mouse model of the human 22q11.2 microdeletion syndrome: a study in male mice. *J Psychiatry Neurosci* 42(1), 48-58.
- Duncan, G.E., Inada, K., Koller, B.H., Moy, S.S., 2010. Increased sensitivity to kainic acid in a genetic model of reduced NMDA receptor function. *Brain Res* 1307, 166-176.
- Ellenbroek, B.A., Budde, S., Cools, A.R., 1996. Prepulse inhibition and latent inhibition: the role of dopamine in the medial prefrontal cortex. *Neuroscience* 75(2), 535-542.
- Fendt, M., Li, L., Yeomans, J.S., 2001. Brain stem circuits mediating prepulse inhibition of the startle reflex. *Psychopharmacology (Berl)* 156(2-3), 216-224.
- Frost, W.N., Tian, L.M., Hoppe, T.A., Mongeluzi, D.L., Wang, J., 2003. A cellular mechanism for prepulse inhibition. *Neuron* 40(5), 991-1001.
- Geyer, M.A., Krebs-Thomson, K., Braff, D.L., Swerdlow, N.R., 2001. Pharmacological studies of prepulse inhibition models of sensorimotor gating deficits in schizophrenia: a decade in review. *Psychopharmacology (Berl)* 156(2-3), 117-154.
- Giakoumaki, S.G., Roussos, P., Bitsios, P., 2008. Improvement of prepulse inhibition and executive function by the COMT inhibitor tolcapone depends on COMT Val158Met polymorphism. *Neuropsychopharmacology* 33(13), 3058-3068.
- Gogos, J.A., Santha, M., Takacs, Z., Beck, K.D., Luine, V., Lucas, L.R., Nadler, J.V., Karayiorgou, M., 1999. The gene encoding proline dehydrogenase modulates sensorimotor gating in mice. *Nat Genet* 21(4), 434-439.
- Gonzalez-Castro, T.B., Hernandez-Diaz, Y., Juarez-Rojop, I.E., Lopez-Narvaez, M.L., Tovilla-Zarate, C.A., Fresan, A., 2016. The Role of a Catechol-O-Methyltransferase (COMT) Val158Met Genetic Polymorphism in Schizophrenia: A Systematic Review and Updated Meta-analysis on 32,816 Subjects. *Neuromolecular Med* 18(2), 216-231.
- Gottesman, I.I., Gould, T.D., 2003. The endophenotype concept in psychiatry: etymology and strategic intentions. *Am J Psychiatry* 160(4), 636-645.
- Graham, F.K., 1975. Presidential Address, 1974. The more or less startling effects of weak prestimulation. *Psychophysiology* 12(3), 238-248.
- Greenwood, T.A., Braff, D.L., Light, G.A., Cadenhead, K.S., Calkins, M.E., Dobie, D.J., Freedman, R., Green, M.F., Gur, R.E., Gur, R.C., Mintz, J., Nuechterlein, K.H., Olincy, A., Radant, A.D., Seidman, L.J., Siever, L.J., Silverman, J.M., Stone, W.S., Swerdlow, N.R., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Schork, N.J., 2007. Initial heritability analyses of endophenotypic measures for schizophrenia: the consortium on the genetics of schizophrenia. *Arch Gen Psychiatry* 64(11), 1242-1250.
- Greenwood, T.A., Light, G.A., Swerdlow, N.R., Radant, A.D., Braff, D.L., 2012. Association analysis of 94 candidate genes and schizophrenia-related endophenotypes. *PLoS One* 7(1), e29630.
- Greenwood, T.A., Light, G.A., Swerdlow, N.R., Calkins, M.E., Green, M.F., Gur, R.E., Gur, R.C., Lazzeroni, L.C., Nuechterlein, K.H., Olincy, A., Radant, A.D., Seidman, L.J., Siever, L.J., Silverman, J.M., Stone, W.S., Sugar, C.A., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Freedman, R., Braff, D.L., 2016. Gating Deficit

- Heritability and Correlation With Increased Clinical Severity in Schizophrenia Patients With Positive Family History. *Am J Psychiatry* 173(4), 385-391.
- Hasenkamp, W., Epstein, M.P., Green, A., Wilcox, L., Boshoven, W., Lewison, B., Duncan, E., 2010. Heritability of acoustic startle magnitude, prepulse inhibition, and startle latency in schizophrenia and control families. *Psychiatry Res* 178(2), 236-243.
- Hoffman, H.S., Searle, J.L., 1965. Acoustic variables in the modification of the startle reaction in the rat. *J Comp Physiol Psychol* 60, 53-58.
- Hokyo, A., Kanazawa, T., Uenishi, H., Tsutsumi, A., Kawashige, S., Kikuyama, H., Glatt, S.J., Koh, J., Nishimoto, Y., Matsumura, H., Motomura, N., Yoneda, H., 2010. Habituation in prepulse inhibition is affected by a polymorphism on the NMDA receptor 2B subunit gene (GRIN2B). *Psychiatr Genet* 20(5), 191-198.
- Hong, L.E., Wonodi, I., Stine, O.C., Mitchell, B.D., Thaker, G.K., 2008. Evidence of missense mutations on the neuregulin 1 gene affecting function of prepulse inhibition. *Biol Psychiatry* 63(1), 17-23.
- Hu, W., MacDonald, M.L., Elswick, D.E., Sweet, R.A., 2015. The glutamate hypothesis of schizophrenia: evidence from human brain tissue studies. *Ann N Y Acad Sci* 1338, 38-57.
- Irish Schizophrenia Genomics, C., the Wellcome Trust Case Control, C., 2012. Genome-wide association study implicates HLA-C*01:02 as a risk factor at the major histocompatibility complex locus in schizophrenia. *Biol Psychiatry* 72(8), 620-628.
- Ison, J.R., Leonhard, D.W., 1970. Effects of auditory stimuli on the amplitude of the nictitating membrane reflex of the rabbit (*Oryctolagus cuniculus*). *J Comp Physiol Psychol* 75, 157-164.
- Kennedy, A.J., Rahn, E.J., Paulukaitis, B.S., Savell, K.E., Kordasiewicz, H.B., Wang, J., Lewis, J.W., Posey, J., Strange, S.K., Guzman-Karlsson, M.C., Phillips, S.E., Decker, K., Motley, S.T., Swayze, E.E., Ecker, D.J., Michael, T.P., Day, J.J., Sweatt, J.D., 2016. Tcf4 Regulates Synaptic Plasticity, DNA Methylation, and Memory Function. *Cell Rep* 16(10), 2666-2685.
- Kilic, G., Ismail Kucukali, C., Orhan, N., Ozkok, E., Zengin, A., Aydin, M., Kara, I., 2010. Are GRIK3 (T928G) gene variants in schizophrenia patients different from those in their first-degree relatives? *Psychiatry Res* 175(1-2), 43-46.
- Koch, M., 1999. The neurobiology of startle. *Prog Neurobiol* 59(2), 107-128.
- Kohl, S., Heekeren, K., Klosterkotter, J., Kuhn, J., 2013. Prepulse inhibition in psychiatric disorders--apart from schizophrenia. *J Psychiatr Res* 47(4), 445-452.
- Krauter, E.E., Leonhard, D.W., Ison, J.R., 1973. Inhibition of the human eye blink by brief acoustic stimulus. *J Comp Physiol Psychol* 84, 246-251.
- Kumari, V., Soni, W., Mathew, V.M., Sharma, T., 2000. Prepulse inhibition of the startle response in men with schizophrenia: effects of age of onset of illness, symptoms, and medication. *Arch Gen Psychiatry* 57(6), 609-614.
- Kumari, V., Das, M., Zachariah, E., Ettinger, U., Sharma, T., 2005. Reduced prepulse inhibition in unaffected siblings of schizophrenia patients. *Psychophysiology* 42(5), 588-594.
- Lachman, H.M., Papolos, D.F., Saito, T., Yu, Y.M., Szumlanski, C.L., Weinshilboum, R.M., 1996. Human catechol-O-methyltransferase pharmacogenetics: description of a functional polymorphism and its potential application to neuropsychiatric disorders. *Pharmacogenetics* 6(3), 243-250.
- Levinson, D.F., 2005. Meta-analysis in psychiatric genetics. *Curr Psychiatry Rep* 7(2), 143-151.
- Li, J., Chen, Z., Wang, F., Ouyang, Y., Zhang, N., Yang, M., Yan, M., Zhu, X., He, X., Yuan, D., Jin, T., 2016. Polymorphisms of the TCF4 gene are associated with the risk of schizophrenia in the Han Chinese. *Am J Med Genet B Neuropsychiatr Genet* 171(8), 1006-1012.

- Li, T., Ma, X., Sham, P.C., Sun, X., Hu, X., Wang, Q., Meng, H., Deng, W., Liu, X., Murray, R.M., Collier, D.A., 2004. Evidence for association between novel polymorphisms in the PRODH gene and schizophrenia in a Chinese population. *Am J Med Genet B Neuropsychiatr Genet* 129B(1), 13-15.
- Li, T., Li, Z., Chen, P., Zhao, Q., Wang, T., Huang, K., Li, J., Li, Y., Liu, J., Zeng, Z., Feng, G., He, L., Shi, Y., 2010. Common variants in major histocompatibility complex region and TCF4 gene are significantly associated with schizophrenia in Han Chinese. *Biol Psychiatry* 68(7), 671-673.
- Light, G.A., Swerdlow, N.R., 2015. Future clinical uses of neurophysiological biomarkers to predict and monitor treatment response for schizophrenia. *Ann N Y Acad Sci* 1344, 105-119.
- Light, G.A., Swerdlow, N.R., Thomas, M.L., Calkins, M.E., Green, M.F., Greenwood, T.A., Gur, R.E., Gur, R.C., Lazzeroni, L.C., Nuechterlein, K.H., Pela, M., Radant, A.D., Seidman, L.J., Sharp, R.F., Siever, L.J., Silverman, J.M., Sprock, J., Stone, W.S., Sugar, C.A., Tsuang, D.W., Tsuang, M.T., Braff, D.L., Turetsky, B.I., 2015. Validation of mismatch negativity and P3a for use in multi-site studies of schizophrenia: characterization of demographic, clinical, cognitive, and functional correlates in COGS-2. *Schizophr Res* 163(1-3), 63-72.
- Linn, G.S., Javitt, D.C., 2001. Phencyclidine (PCP)-induced deficits of prepulse inhibition in monkeys. *Neuroreport* 12(1), 117-120.
- Liu, H., Heath, S.C., Sobin, C., Roos, J.L., Galke, B.L., Blundell, M.L., Lenane, M., Robertson, B., Wijsman, E.M., Rapoport, J.L., Gogos, J.A., Karayiorgou, M., 2002. Genetic variation at the 22q11 PRODH2/DGCR6 locus presents an unusual pattern and increases susceptibility to schizophrenia. *Proc Natl Acad Sci U S A* 99(6), 3717-3722.
- Liu, M., Malone, S.M., Vaidyanathan, U., Keller, M.C., Abecasis, G., McGue, M., Iacono, W.G., Vrieze, S.I., 2017. Psychophysiological endophenotypes to characterize mechanisms of known schizophrenia genetic loci. *Psychol Med* 47(6), 1116-1125.
- Liu, X., Hong, X., Chan, R.C., Kong, F., Peng, Z., Wan, X., Wang, C., Cheng, L., 2013. Association study of polymorphisms in the alpha 7 nicotinic acetylcholine receptor subunit and catechol-o-methyl transferase genes with sensory gating in first-episode schizophrenia. *Psychiatry Res* 209(3), 431-438.
- Lohmueller, K.E., Pearce, C.L., Pike, M., Lander, E.S., Hirschhorn, J.N., 2003. Meta-analysis of genetic association studies supports a contribution of common variants to susceptibility to common disease. *Nat Genet* 33(2), 177-182.
- Ludewig, K., Geyer, M.A., Vollenweider, F.X., 2003. Deficits in prepulse inhibition and habituation in never-medicated, first-episode schizophrenia. *Biol Psychiatry* 54(2), 121-128.
- Montag, C., Hartmann, P., Merz, M., Burk, C., Reuter, M., 2008. D2 receptor density and prepulse inhibition in humans: negative findings from a molecular genetic approach. *Behav Brain Res* 187(2), 428-432.
- Owens, E.M., Bachman, P., Glahn, D.C., Bearden, C.E., 2016. Electrophysiological Endophenotypes for Schizophrenia. *Harv Rev Psychiatry* 24(2), 129-147.
- Parwani, A., Duncan, E.J., Bartlett, E., Madonick, S.H., Efferen, T.R., Rajan, R., Sanfilippo, M., Chappell, P.B., Chakravorty, S., Gonzenbach, S., Ko, G.N., Rotrosen, J.P., 2000. Impaired prepulse inhibition of acoustic startle in schizophrenia. *Biol Psychiatry* 47(7), 662-669.
- Perrais, D., Coussen, F., Mulle, C., 2009. Atypical functional properties of GluK3-containing kainate receptors. *J Neurosci* 29(49), 15499-15510.
- Petrovsky, N., Quednow, B.B., Ettinger, U., Schmechtig, A., Mossner, R., Collier, D.A., Kuhn, K.U., Maier, W., Wagner, M., Kumari, V., 2010. Sensorimotor gating is associated with CHRNA3 polymorphisms in schizophrenia and healthy volunteers. *Neuropsychopharmacology* 35(7), 1429-1439.
- Petrovsky, N., Ettinger, U., Kessler, H., Mossner, R., Wolfsgruber, S., Dahmen, N., Maier, W., Wagner, M., Quednow, B.B., 2013. The effect of nicotine on sensorimotor gating is modulated by a CHRNA3 polymorphism. *Psychopharmacology (Berl)* 229(1), 31-40.

- Phang, J.M., Donald, S.P., Pandhare, J., Liu, Y., 2008. The metabolism of proline, a stress substrate, modulates carcinogenic pathways. *Amino Acids* 35(4), 681-690.
- Prasad, S.E., Howley, S., Murphy, K.C., 2008. Candidate genes and the behavioral phenotype in 22q11.2 deletion syndrome. *Dev Disabil Res Rev* 14(1), 26-34.
- Quednow, B.B., Wagner, M., Westheide, J., Beckmann, K., Bliesener, N., Maier, W., Kuhn, K.U., 2006. Sensorimotor gating and habituation of the startle response in schizophrenic patients randomly treated with amisulpride or olanzapine. *Biol Psychiatry* 59(6), 536-545.
- Quednow, B.B., 2008. Defizite der sensomotorischen Filterleistung bei psychiatrischen Erkrankungen. *Z Neuropsychol* 19(3), 139-163.
- Quednow, B.B., Frommann, I., Berning, J., Kuhn, K.U., Maier, W., Wagner, M., 2008a. Impaired sensorimotor gating of the acoustic startle response in the prodrome of schizophrenia. *Biol Psychiatry* 64(9), 766-773.
- Quednow, B.B., Kuhn, K.U., Mossner, R., Schwab, S.G., Schuhmacher, A., Maier, W., Wagner, M., 2008b. Sensorimotor gating of schizophrenia patients is influenced by 5-HT_{2A} receptor polymorphisms. *Biol Psychiatry* 64(5), 434-437.
- Quednow, B.B., Schmechtig, A., Ettinger, U., Petrovsky, N., Collier, D.A., Vollenweider, F.X., Wagner, M., Kumari, V., 2009. Sensorimotor gating depends on polymorphisms of the serotonin-2A receptor and catechol-O-methyltransferase, but not on neuregulin-1 Arg38Gln genotype: a replication study. *Biol Psychiatry* 66(6), 614-620.
- Quednow, B.B., Wagner, M., Mossner, R., Maier, W., Kuhn, K.U., 2010. Sensorimotor gating of schizophrenia patients depends on Catechol O-methyltransferase Val158Met polymorphism. *Schizophr Bull* 36(2), 341-346.
- Quednow, B.B., Ettinger, U., Mossner, R., Rujescu, D., Giegling, I., Collier, D.A., Schmechtig, A., Kuhn, K.U., Moller, H.J., Maier, W., Wagner, M., Kumari, V., 2011. The schizophrenia risk allele C of the TCF4 rs9960767 polymorphism disrupts sensorimotor gating in schizophrenia spectrum and healthy volunteers. *J Neurosci* 31(18), 6684-6691.
- Quednow, B.B., Brinkmeyer, J., Mobascher, A., Nothnagel, M., Musso, F., Grunder, G., Savary, N., Petrovsky, N., Frommann, I., Lennertz, L., Spreckelmeyer, K.N., Wienker, T.F., Dahmen, N., Thuerauf, N., Clepce, M., Kiefer, F., Majic, T., Mossner, R., Maier, W., Gallinat, J., Diaz-Lacava, A., Toliat, M.R., Thiele, H., Nurnberg, P., Wagner, M., Winterer, G., 2012. Schizophrenia risk polymorphisms in the TCF4 gene interact with smoking in the modulation of auditory sensory gating. *Proc Natl Acad Sci U S A* 109(16), 6271-6276.
- Quednow, B.B., Brzozka, M.M., Rossner, M.J., 2014. Transcription factor 4 (TCF4) and schizophrenia: integrating the animal and the human perspective. *Cell Mol Life Sci* 71(15), 2815-2835.
- Ripke, S., O'Dushlaine, C., Chambert, K., Moran, J.L., Kahler, A.K., Akterin, S., Bergen, S.E., Collins, A.L., Crowley, J.J., Fromer, M., Kim, Y., Lee, S.H., Magnusson, P.K., Sanchez, N., Stahl, E.A., Williams, S., Wray, N.R., Xia, K., Bettella, F., Borglum, A.D., Bulik-Sullivan, B.K., Cormican, P., Craddock, N., de Leeuw, C., Durmishi, N., Gill, M., Golimbet, V., Hamshere, M.L., Holmans, P., Hougaard, D.M., Kendler, K.S., Lin, K., Morris, D.W., Mors, O., Mortensen, P.B., Neale, B.M., O'Neill, F.A., Owen, M.J., Milovancevic, M.P., Posthuma, D., Powell, J., Richards, A.L., Riley, B.P., Ruderfer, D., Rujescu, D., Sigurdsson, E., Silagadze, T., Smit, A.B., Stefansson, H., Steinberg, S., Suvisaari, J., Tosato, S., Verhage, M., Walters, J.T., Multicenter Genetic Studies of Schizophrenia, C., Levinson, D.F., Gejman, P.V., Kendler, K.S., Laurent, C., Mowry, B.J., O'Donovan, M.C., Owen, M.J., Pulver, A.E., Riley, B.P., Schwab, S.G., Wildenauer, D.B., Dudbridge, F., Holmans, P., Shi, J., Albus, M., Alexander, M., Campion, D., Cohen, D., Dikeos, D., Duan, J., Eichhammer, P., Godard, S., Hansen, M., Lerer, F.B., Liang, K.Y., Maier, W., Mallet, J., Nertney, D.A., Nestadt, G., Norton, N., O'Neill, F.A., Papadimitriou, G.N., Ribble, R., Sanders, A.R., Silverman, J.M., Walsh, D., Williams, N.M., Wormley, B., Psychosis Endophenotypes International, C., Arranz, M.J., Bakker, S., Bender, S., Bramon, E., Collier, D., Crespo-Facorro, B., Hall, J., Iyegbe, C., Jablensky, A., Kahn, R.S., Kalaydjieva, L., Lawrie, S., Lewis, C.M., Lin, K., Linszen, D.H., Mata, I., McIntosh, A., Murray, R.M., Ophoff, R.A., Powell, J., Rujescu, D., Van Os, J., Walshe, M., Weisbrod, M., Wiersma, D., Wellcome Trust Case Control, C., Donnelly, P., Barroso, I., Blackwell, J.M., Bramon, E., Brown, M.A., Casas, J.P., Corvin, A.P., Deloukas, P., Duncanson, A., Jankowski, J., Markus, H.S., Mathew, C.G., Palmer, C.N., Plomin, R., Rautanen, A., Sawcer, S.J., Trembath, R.C., Viswanathan, A.C., Wood, N.W., Spencer, C.C., Band, G., Bellenguez, C., Freeman, C., Hellenthal, G.,

- Giannoulatou, E., Pirinen, M., Pearson, R.D., Strange, A., Su, Z., Vukcevic, D., Donnelly, P., Langford, C., Hunt, S.E., Edkins, S., Gwilliam, R., Blackburn, H., Bumpstead, S.J., Dronov, S., Gillman, M., Gray, E., Hammond, N., Jayakumar, A., McCann, O.T., Liddle, J., Potter, S.C., Ravindrarajah, R., Ricketts, M., Tashakkori-Ghanbaria, A., Waller, M.J., Weston, P., Widaa, S., Whittaker, P., Barroso, I., Deloukas, P., Mathew, C.G., Blackwell, J.M., Brown, M.A., Corvin, A.P., McCarthy, M.I., Spencer, C.C., Bramon, E., Corvin, A.P., O'Donovan, M.C., Stefansson, K., Scolnick, E., Purcell, S., McCarroll, S.A., Sklar, P., Hultman, C.M., Sullivan, P.F., 2013. Genome-wide association analysis identifies 13 new risk loci for schizophrenia. *Nat Genet* 45(10), 1150-1159.
- Rivett, A.J., Eddy, B.J., Roth, J.A., 1982. Contribution of sulfate conjugation, deamination, and O-methylation to metabolism of dopamine and norepinephrine in human brain. *J Neurochem* 39(4), 1009-1016.
- Rosenthal, R., 1984. *Meta-Analytic Procedures for Social Research*. Sage, Newbury Park.
- Roussos, P., Giakoumaki, S.G., Bitsios, P., 2008a. The dopamine D(3) receptor Ser9Gly polymorphism modulates prepulse inhibition of the acoustic startle reflex. *Biol Psychiatry* 64(3), 235-240.
- Roussos, P., Giakoumaki, S.G., Rogdaki, M., Pavlakis, S., Frangou, S., Bitsios, P., 2008b. Prepulse inhibition of the startle reflex depends on the catechol O-methyltransferase Val158Met gene polymorphism. *Psychol Med* 38(11), 1651-1658.
- Roussos, P., Giakoumaki, S.G., Bitsios, P., 2009a. A risk PRODH haplotype affects sensorimotor gating, memory, schizotypy, and anxiety in healthy male subjects. *Biol Psychiatry* 65(12), 1063-1070.
- Roussos, P., Giakoumaki, S.G., Bitsios, P., 2009b. Tolcapone effects on gating, working memory, and mood interact with the synonymous catechol-O-methyltransferase rs4818c/g polymorphism. *Biol Psychiatry* 66(11), 997-1004.
- Roussos, P., Giakoumaki, S.G., Adamaki, E., Bitsios, P., 2011. The influence of schizophrenia-related neuregulin-1 polymorphisms on sensorimotor gating in healthy males. *Biol Psychiatry* 69(5), 479-486.
- Roussos, P., Giakoumaki, S.G., Zouraraki, C., Fullard, J.F., Karagiorga, V.E., Tsapakis, E.M., Petraki, Z., Siever, L.J., Lencz, T., Malhotra, A., Spanaki, C., Bitsios, P., 2016. The Relationship of Common Risk Variants and Polygenic Risk for Schizophrenia to Sensorimotor Gating. *Biol Psychiatry* 79(12), 988-996.
- Schiffer, H.H., Heinemann, S.F., 2007. Association of the human kainate receptor GluR7 gene (GRIK3) with recurrent major depressive disorder. *Am J Med Genet B Neuropsychiatr Genet* 144B(1), 20-26.
- Schizophrenia Psychiatric Genome-Wide Association Study, C., 2011. Genome-wide association study identifies five new schizophrenia loci. *Nat Genet* 43(10), 969-976.
- Schizophrenia Working Group of the Psychiatric Genomics, C., 2014. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 511(7510), 421-427.
- Seidman, L.J., Helleman, G., Nuechterlein, K.H., Greenwood, T.A., Braff, D.L., Cadenhead, K.S., Calkins, M.E., Freedman, R., Gur, R.E., Gur, R.C., Lazzaroni, L.C., Light, G.A., Olincy, A., Radant, A.D., Siever, L.J., Silverman, J.M., Sprock, J., Stone, W.S., Sugar, C., Swerdlow, N.R., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Green, M.F., 2015. Factor structure and heritability of endophenotypes in schizophrenia: findings from the Consortium on the Genetics of Schizophrenia (COGS-1). *Schizophr Res* 163(1-3), 73-79.
- Shi, J., Wang, Z., Tan, Y., Fan, H., An, H., Zuo, L., Yang, F., Tan, S., Li, J., Zhang, X., Zhou, D., Luo, X., 2016. CHRNA4 was associated with prepulse inhibition of schizophrenia in Chinese: a pilot study. *Cogn Neuropsychiatry* 21(2), 156-167.
- Sobin, C., Kiley-Brabeck, K., Karayiorgou, M., 2005a. Associations between prepulse inhibition and executive visual attention in children with the 22q11 deletion syndrome. *Mol Psychiatry* 10(6), 553-562.
- Sobin, C., Kiley-Brabeck, K., Karayiorgou, M., 2005b. Lower prepulse inhibition in children with the 22q11 deletion syndrome. *Am J Psychiatry* 162(6), 1090-1099.

Stefansson, H., Ophoff, R.A., Steinberg, S., Andreassen, O.A., Cichon, S., Rujescu, D., Werge, T., Pietilainen, O.P., Mors, O., Mortensen, P.B., Sigurdsson, E., Gustafsson, O., Nyegaard, M., Tuulio-Henriksson, A., Ingason, A., Hansen, T., Suvisaari, J., Lonnqvist, J., Paunio, T., Borglum, A.D., Hartmann, A., Fink-Jensen, A., Nordentoft, M., Hougaard, D., Norgaard-Pedersen, B., Bottcher, Y., Olesen, J., Breuer, R., Moller, H.J., Giegling, I., Rasmussen, H.B., Timm, S., Mattheisen, M., Bitter, I., Rethelyi, J.M., Magnusdottir, B.B., Sigmundsson, T., Olason, P., Masson, G., Gulcher, J.R., Haraldsson, M., Fossdal, R., Thorgeirsson, T.E., Thorsteinsdottir, U., Ruggeri, M., Tosato, S., Franke, B., Strengman, E., Kiemenev, L.A., Melle, I., Djurovic, S., Abramova, L., Kaleda, V., Sanjuan, J., de Frutos, R., Bramon, E., Vassos, E., Fraser, G., Ettinger, U., Picchioni, M., Walker, N., Touloupoulou, T., Need, A.C., Ge, D., Yoon, J.L., Shianna, K.V., Freimer, N.B., Cantor, R.M., Murray, R., Kong, A., Golimbet, V., Carracedo, A., Arango, C., Costas, J., Jonsson, E.G., Terenius, L., Agartz, I., Petursson, H., Nothen, M.M., Rietschel, M., Matthews, P.M., Muglia, P., Peltonen, L., St Clair, D., Goldstein, D.B., Stefansson, K., Collier, D.A., 2009. Common variants conferring risk of schizophrenia. *Nature* 460(7256), 744-747.

Steinberg, S., de Jong, S., Irish Schizophrenia Genomics, C., Andreassen, O.A., Werge, T., Borglum, A.D., Mors, O., Mortensen, P.B., Gustafsson, O., Costas, J., Pietilainen, O.P., Demontis, D., Papiol, S., Huttenlocher, J., Mattheisen, M., Breuer, R., Vassos, E., Giegling, I., Fraser, G., Walker, N., Tuulio-Henriksson, A., Suvisaari, J., Lonnqvist, J., Paunio, T., Agartz, I., Melle, I., Djurovic, S., Strengman, E., Group, Jurgens, G., Glenthøj, B., Terenius, L., Hougaard, D.M., Orntoft, T., Wiuf, C., Didriksen, M., Hollegaard, M.V., Nordentoft, M., van Winkel, R., Kenis, G., Abramova, L., Kaleda, V., Arrojo, M., Sanjuan, J., Arango, C., Sperling, S., Rossner, M., Ribolsi, M., Magni, V., Siracusano, A., Christiansen, C., Kiemenev, L.A., Veldink, J., van den Berg, L., Ingason, A., Muglia, P., Murray, R., Nothen, M.M., Sigurdsson, E., Petursson, H., Thorsteinsdottir, U., Kong, A., Rubino, I.A., De Hert, M., Rethelyi, J.M., Bitter, I., Jonsson, E.G., Golimbet, V., Carracedo, A., Ehrenreich, H., Craddock, N., Owen, M.J., O'Donovan, M.C., Wellcome Trust Case Control, C., Ruggeri, M., Tosato, S., Peltonen, L., Ophoff, R.A., Collier, D.A., St Clair, D., Rietschel, M., Cichon, S., Stefansson, H., Rujescu, D., Stefansson, K., 2011. Common variants at VRK2 and TCF4 conferring risk of schizophrenia. *Hum Mol Genet* 20(20), 4076-4081.

Stouffer, S.A., Suchman, E.A., DeVinney, L.C., Star, S.A., Williams, R.M., 1949. *The American Soldier: Adjustment During Army Life - Volume 1*. Princeton University Press, Princeton, NJ.

Strimmer, K., 2008. A unified approach to false discovery rate estimation. *BMC Bioinformatics* 9, 303.

Swerdlow, N.R., Geyer, M.A., Braff, D.L., 2001. Neural circuit regulation of prepulse inhibition of startle in the rat: current knowledge and future challenges. *Psychopharmacology (Berl)* 156(2-3), 194-215.

Swerdlow, N.R., Hines, S.R., Herrera, S.D., Weber, M., Breier, M.R., 2013. Opposite effects of tolcapone on amphetamine-disrupted startle gating in low vs. high COMT-expressing rat strains. *Pharmacol Biochem Behav* 106, 128-131.

Swerdlow, N.R., Light, G.A., Sprock, J., Calkins, M.E., Green, M.F., Greenwood, T.A., Gur, R.E., Gur, R.C., Lazzeroni, L.C., Nuechterlein, K.H., Radant, A.D., Ray, A., Seidman, L.J., Siever, L.J., Silverman, J.M., Stone, W.S., Sugar, C.A., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Braff, D.L., 2014. Deficient prepulse inhibition in schizophrenia detected by the multi-site COGS. *Schizophr Res* 152(2-3), 503-512.

Swerdlow, N.R., Bhakta, S.G., Rana, B.K., Kei, J., Chou, H.H., Talledo, J.A., 2017. Sensorimotor gating in healthy adults tested over a 15 year period. *Biol Psychol* 123, 177-186.

Takenouchi, T., Hashida, N., Torii, C., Kosaki, R., Takahashi, T., Kosaki, K., 2014. 1p34.3 deletion involving GRIK3: Further clinical implication of GRIK family glutamate receptors in the pathogenesis of developmental delay. *Am J Med Genet A* 164A(2), 456-460.

Taylor, S., 2017. Association between COMT Val158Met and psychiatric disorders: A comprehensive meta-analysis. *Am J Med Genet B Neuropsychiatr Genet*.

Tunbridge, E.M., Harrison, P.J., Weinberger, D.R., 2006. Catechol-o-methyltransferase, cognition, and psychosis: Val158Met and beyond. *Biol Psychiatry* 60(2), 141-151.

Tunbridge, E.M., Harrison, P.J., 2011. Importance of the COMT gene for sex differences in brain function and predisposition to psychiatric disorders. *Curr Top Behav Neurosci* 8, 119-140.

- Vaidyanathan, U., Malone, S.M., Miller, M.B., McGue, M., Iacono, W.G., 2014. Heritability and molecular genetic basis of acoustic startle eye blink and affectively modulated startle response: a genome-wide association study. *Psychophysiology* 51(12), 1285-1299.
- Volter, C., Riedel, M., Wostmann, N., Aichert, D.S., Lobo, S., Costa, A., Schmechtig, A., Collier, D.A., Hartmann, A.M., Giegling, I., Moller, H.J., Quednow, B.B., Rujescu, D., Kumari, V., Ettinger, U., 2012. Sensorimotor gating and D2 receptor signalling: evidence from a molecular genetic approach. *Int J Neuropsychopharmacol* 15(10), 1427-1440.
- Willott, J.F., Carlson, S., Chen, H., 1994. Prepulse inhibition of the startle response in mice: relationship to hearing loss and auditory system plasticity. *Behav Neurosci* 108(703-13).
- Willott, J.F., Tanner, L., O'Steen, J., Johnson, K.R., Bogue, M.A., Gagnon, L., 2003. Acoustic startle and prepulse inhibition in 40 inbred strains of mice. *Behav Neurosci* 117(4), 716-727.
- Zavitsanou, K., Cranney, J., Richardson, R., 1999. Dopamine antagonists in the orbital prefrontal cortex reduce prepulse inhibition of the acoustic startle reflex in the rat. *Pharmacol Biochem Behav* 63(1), 55-61.
- Ziermans, T., Schothorst, P., Magnee, M., van Engeland, H., Kemner, C., 2011. Reduced prepulse inhibition in adolescents at risk for psychosis: a 2-year follow-up study. *J Psychiatry Neurosci* 36(2), 127-134.
- Ziermans, T.B., Schothorst, P.F., Sprong, M., Magnee, M.J., van Engeland, H., Kemner, C., 2012. Reduced prepulse inhibition as an early vulnerability marker of the psychosis prodrome in adolescence. *Schizophr Res* 134(1), 10-15.
- Zweier, C., Peippo, M.M., Hoyer, J., Sousa, S., Bottani, A., Clayton-Smith, J., Reardon, W., Saraiva, J., Cabral, A., Gohring, I., Devriendt, K., de Ravel, T., Bijlsma, E.K., Hennekam, R.C., Orrico, A., Cohen, M., Dreweke, A., Reis, A., Nurnberg, P., Rauch, A., 2007. Haploinsufficiency of TCF4 causes syndromal mental retardation with intermittent hyperventilation (Pitt-Hopkins syndrome). *Am J Hum Genet* 80(5), 994-1001.

Tables

Table 1. Summary list of published and unpublished studies included in meta-analysis. HC = healthy controls, SCZ = patients with schizophrenia.

Reference	Sample (location)	Sample number	Analyzed N
(Brauer et al., 2009)	HC (Giessen)	1	81
(Greenwood et al., 2012)	SCZ (San Diego)	2	219
(Hong et al., 2008)	HC (Maryland)	3	63
	SCZ (Maryland)	4	113
(Hokyo et al., 2010)	HC (Osaka)	5	71
	SCZ (Osaka)	6	81
(Liu et al., 2013)	SCZ (Guangdong)	7	140
(Montag et al., 2008)	HC (Bonn-Montag)	8	96
(Petrovsky et al., 2010)	HC (London)	9	96
	SCZ (Bonn)	10	68
(Petrovsky et al., 2013)	HC (Bonn)	11	63
(Quednow, Ettinger, Kumari, unpublished data)	HC (London)	9	100
(Quednow and Wagner, unpublished data)	SCZ (Bonn)	10	107
(Quednow et al., 2008b)	SCZ (Bonn)	10	68
(Quednow et al., 2009)	HC (London)	9	99
(Quednow et al., 2010)	SCZ (Bonn)	10	71
(Quednow et al., 2011)	HC (London)	9	98
	SCZ (Bonn)	10	105
(Roussos et al., 2008a)	HC (Crete)	12	101
(Roussos et al., 2008b)	HC (Crete)	12	93
(Roussos et al., 2009a)	HC (Crete)	12	217
(Roussos et al., 2011)	HC (LOGOS)	13	445
(Roussos et al., 2016)	HC (LOGOS GWAS)	14	686
(Shi et al., 2016)	SCZ (Beijing)	15	77
(Volter et al., 2012)	HC (London)	9	96
	HC (Munich)	16	101

Varying sample sizes among single samples (e.g., sample nr. 10: range n=68-107) are explained by different schizophrenia spectrum diagnosis included or because genotype was only available in a subsample (e.g., due to genotyping failures).

Table 2. Combined and Z-scores and P-values across genotypes where data were available at least in two independent samples. The signed Z-score indicates the direction of association among genotype and PPI. Negative Z-scores indicate decreased PPI in the reference compared to the alternative allele; Positive Z-scores indicate increased PPI in the reference compared to the alternative allele. Results with false discovery rate (FDR) ≤ 0.05 are in bold.

SNP	Gene	Number of studies	Total sample	Reference allele	Combined Z-score	Cohen's d	Combined P-value	FDR
rs4680 (only males)	COMT	4	870	A	4.03	0.28	5.51E-05	0.002
rs1027599	GRIK3	2	905	C	2.87	0.19	0.004	0.045
rs9960767	TCF4	3	889	C	-2.77	0.19	0.006	0.050
rs385440	PRODH	2	436	A	-2.75	0.27	0.006	0.050
rs533337	GRIK3	2	905	A	2.64	0.18	0.008	0.055
rs4680	COMT	6	1,179	A	2.58	0.15	0.010	0.057
rs13101891	GRID2	2	905	G	-2.41	0.16	0.016	0.067
rs1800497	DRD2	5	1,047	A2	-2.40	0.15	0.017	0.067
rs4782262	GRIN2A	2	905	C	2.36	0.16	0.018	0.069
rs876848	GAD2	2	905	T	-2.27	0.15	0.023	0.074
rs1044396	CHRNA4	3	863	T	-2.26	0.15	0.024	0.075
rs1923292	SLC1A2	2	905	G	-2.15	0.14	0.032	0.097
rs40184	SLC6A3	3	998	A	2.14	0.14	0.033	0.100
rs4852550	CTNNA2	2	905	T	-2.06	0.14	0.039	0.118
rs308787	GRM5	2	905	G	2.04	0.14	0.041	0.122
rs3924999	NRG1	5	1,180	A	-2.03	0.12	0.042	0.126
rs8068673	PAFAH1B1	2	905	C	2.01	0.13	0.045	0.132
rs6277	DRD2	4	905	G	-2.00	0.13	0.046	0.135
rs701567	DAOA	2	905	G	1.91	0.13	0.057	0.162
rs1587526	GRIK3	2	905	A	1.90	0.13	0.057	0.163
rs894829	CTNNA2	2	905	G	-1.89	0.13	0.059	0.167
rs1266475	PAFAH1B1	2	905	C	1.88	0.13	0.060	0.169
rs6311	5HT2AR	3	851	A	1.87	0.13	0.062	0.173
rs1051730	CHRNA3	4	902	T	-1.79	0.12	0.074	0.200
rs6994992	NRG1	3	1,224	T	-1.79	0.10	0.074	0.202

rs10095556	<i>NRG1</i>	2	905	G	1.72	0.11	0.086	0.226
rs12685902	<i>GRIN3A</i>	2	905	C	-1.68	0.11	0.094	0.242
rs11782671	<i>NRG1</i>	2	905	T	1.66	0.11	0.098	0.250
rs9297186	<i>NRG1</i>	2	905	A	1.59	0.11	0.111	0.275
rs778294	<i>DAOA</i>	3	1,003	A	1.36	0.09	0.175	0.373
rs6280	<i>DRD3</i>	4	1,070	G	-1.34	0.08	0.181	0.381
rs8081803	<i>PAFAH1B1</i>	2	905	C	1.21	0.08	0.225	0.433
rs6313	<i>5HT2AR</i>	5	1,151	T	1.12	0.07	0.265	0.474
rs2619539	<i>DTNBP1</i>	4	1,105	C	-0.99	0.06	0.324	0.524
rs2008626	<i>NRG1</i>	2	905	C	-0.97	0.06	0.334	0.532
rs1368909	<i>CTNNA2</i>	2	905	C	-0.93	0.06	0.350	0.544
rs165599	<i>COMT</i>	3	928	G	-0.81	0.05	0.417	0.586
rs3814614	<i>GRID1</i>	2	905	C	0.79	0.05	0.431	0.595
rs7555221	<i>GRIK3</i>	2	905	C	-0.56	0.04	0.578	0.663
rs4648317	<i>DRD2</i>	4	948	G	-0.52	0.03	0.601	0.672
rs2619528	<i>DTNBP1</i>	4	941	G	0.49	0.03	0.625	0.680
rs1486009	<i>DRD3</i>	2	905	G	-0.48	0.03	0.634	0.683
rs4584372	<i>NOS1AP</i>	2	905	C	-0.34	0.02	0.736	0.715
rs1011313	<i>DTNBP1</i>	4	1,108	A	0.08	0.01	0.933	0.760