

Secondary Stroke Prevention following Embolic Stroke of Unknown Source in the Absence of Documented Atrial Fibrillation: A Clinical Review

Kotadia et al.

Authors:

Irum D. Kotadia^{1,2} MBBS, BSc, MRCP

Iain Sim¹ MBBS, BSc, MRCP

Rahul Mukherjee¹ MBBS, PhD, MRCP

Daniel O'Hare¹ MBBS, BEng, MRCP

Amedeo Chiribiri¹ MBBS, MD, PhD, FSCMR

Jonathan Birns² MBBS, PhD, FRCP

Ajay Bhalla² MBBS, MD, FRCP

Mark O'Neill^{1,2} MB BCh, DPhil, FRCP, FHRS

Steven E. Williams^{1,2,3} MB ChB, PhD, MRCP

¹ King's College London, United Kingdom

² Guy's and St Thomas' NHS Foundation Trust, London, United Kingdom

³ Centre for Cardiovascular Science, The University of Edinburgh

Address for correspondence: Dr. I D Kotadia, Division of Imaging Sciences and Biomedical Engineering, King's College London, 4th Floor North Wing, St. Thomas' Hospital, 249 Westminster Bridge Road, London, SE1 7EH; irum.kotadia@kcl.ac.uk; Telephone 020 7188 7188; Fax 020 7188 5442

Manuscript type: Review Article

Word Count: 5608

Conflict of Interest Statement:

Professor O'Neill has received research support and honoraria from Biosense Webster and has received consultation fees from Medtronic, Biosense Webster, St. Jude/Abbott and Siemens.

Dr Williams has received research support from Biosense Webster.

Abstract

Approximately one third of ischaemic strokes are classed as cryptogenic strokes. The risk of stroke recurrence in these patients is significantly elevated with up to one third of cryptogenic stroke patients experiencing a further stroke within 10 years. Whilst anticoagulation is the mainstay of treatment for secondary stroke prevention in the context of documented atrial fibrillation, it is estimated that up to 25% of cryptogenic stroke patients have undiagnosed atrial fibrillation.

Furthermore, the historical acceptance of a causal relationship between atrial fibrillation and stroke has recently come under scrutiny, with evidence to suggest that embolic stroke risk may be elevated even in the absence of documented atrial fibrillation due to the presence of electrical and structural changes constituting an atrial cardiomyopathy.

More recently, the term 'Embolic Stroke of Unknown Source' has garnered increasing interest as a subset of cryptogenic stroke patients in whom a minimum set of diagnostic investigations has been performed and a non-lacunar infarct highly suspicious of embolic aetiology is suspected but in the absence of an identifiable secondary cause of stroke. The ongoing ARCADIA and ATTICUS studies seek to further define this novel term.

This review summarises the relationship between AF, embolic stroke and atrial cardiomyopathy and provides an overview of the clinical relevance of cardiac imaging, electrocardiographic and serum biomarkers in the assessment of AF and secondary stroke risk. The implications of these findings on therapeutic considerations is considered and gaps in the literature identified as areas for future study in risk stratifying this cohort of patients.

Keywords

Ischaemic Stroke; Embolic Stroke of Unknown Source; Atrial Fibrillation; Atrial Cardiomyopathy; Cardiac magnetic resonance imaging; Electrocardiogram

Acknowledgements

SW acknowledges support from the British Heart Foundation (FS/20/26/34952). This work was supported by the Wellcome/EPSRC Centre for Medical Engineering (WT203148/Z/16/Z) at King's College London and the British Heart Foundation (PG/19/44/34368).

Non-standard Abbreviations and Acronyms

AF	Atrial Fibrillation
AT	Atrial Tachycardia
CI	Confidence Interval
CMR	Cardiac Magnetic Resonance Imaging
ECG	Electrocardiography
ECM	Extracellular Matrix
EHRA	European Heart Rhythm Association
ESUS	Embolic Stroke of Unknown Source
HR	Hazard Ratio
HRS	Heart Rhythm Society
LA	Left atrium
LAA	left atrial appendage
LGE	Late Gadolinium Enhancement
NT-proBNP	N-terminal pro-brain natriuretic peptide
OR	Odds Ratio
PAF	Paroxysmal Atrial Fibrillation
PsAF	Persistent Atrial Fibrillation
PTFV ₁	P-wave terminal fove in lead V ₁
RR	Relative Risk
TE	Thromboembolic Stroke

1. Introduction

Ischaemic stroke remains a leading cause of mortality and morbidity.¹ Atrial fibrillation (AF) is a well-recognised risk factor for stroke, accounting for up to 20% of ischaemic events and AF-related strokes are associated with a significantly higher risk of disability or fatality compared with non-AF strokes.¹⁻³ Whilst anticoagulation is effective in the secondary prevention of thromboembolic stroke, current guidelines recommend documentation of AF prior to initiation of anticoagulation therapy.⁴ However, a diagnosis of AF can be elusive due to its paroxysmal and often asymptomatic nature, resulting in a proportion of thromboembolic strokes being erroneously classified as cryptogenic strokes.^{5,6} There is no clear consensus on the appropriate length of investigation required for detecting AF or duration of AF that is diagnostic, but recent trials have suggested that 30-day outpatient event-triggered monitoring or implantation of a loop recorder may increase the detection rate of AF after cryptogenic stroke by almost 30% over 3 years⁷⁻⁹ and 30 seconds is often used as an arbitrary cut-off for diagnosis.^{4,10}

A more recent term, 'embolic stroke of unknown source' (ESUS), describes a subset of cryptogenic stroke patients in whom a non-lacunar infarct highly suspicious of embolic aetiology is suspected but in the absence of an identifiable secondary cause of stroke such as embolism, arteritis, dissection and coagulopathy.¹¹ It has been hypothesised that anticoagulation therapy may benefit this group, particularly those with evidence of an atrial cardiomyopathy.^{11,12}

Separately, there is growing evidence of a more complex interaction between AF and ischaemic stroke than traditionally thought. Inconsistencies regarding temporal association and biological gradient between AF burden and ischaemic stroke suggest that a conceptual shift may be required to understand the risk of stroke beyond traditional scoring methods.¹³ Co-morbidities

or risk factors leading to underlying atrial disease may provide an atrial thromboembolic substrate (*figure 1*).^{13,14} Such models acknowledge that the risk of embolic stroke may be elevated even in the absence of diagnosed AF. A post-stroke pathway consisting of clinical investigations aimed at detecting an atrial cardiomyopathy could potentially identify a cohort of patients at high risk for further cardioembolic strokes but without diagnosed AF and who may therefore arguably benefit from anticoagulation therapy.

Previous studies using non-invasive investigations such as atrial imaging, electrocardiography (ECG) and serum biomarkers have been highlighted as potential methods to identify and/or quantify structural and electrical atrial remodelling suggestive of an atrial cardiomyopathy. Atrial cardiac magnetic resonance imaging provides non-invasive characterisation of atrial tissue and has been shown to identify changes in atrial geometry, contractile function and fibrosis which together constitute structural remodelling and may reflect a pro-thrombotic atrial cardiomyopathy.¹⁵ Alternatively, electrocardiographic and implantable device parameters such as atrial-high-rate episodes, atrial ectopy burden, P-wave terminal force in lead V₁ and PR interval have all been associated with increased ischaemic stroke risk in the absence of diagnosed AF. Many of these markers have been further shown to independently correlate with structural atrial remodelling including fibrosis, hypertrophy and dilatation and are thought to constitute electrical remodelling in the context of an atrial cardiomyopathy.¹²

This review will consider the relationship between AF, stroke and atrial cardiomyopathy. Evidence pertaining to the clinical relevance of imaging, electrocardiographic and serum biomarkers in the diagnosis of atrial cardiomyopathy and future stroke risk will be summarised. Impact on future therapeutic considerations will also be explored.

2. Antiplatelet vs. Anticoagulation Therapy for Secondary Stroke Prevention

Anticoagulation in Cryptogenic Stroke

The Warfarin-Aspirin Recurrent Stroke Study (WARSS) was the first large randomised-control study to perform a sub-group analysis that assessed the use of warfarin versus aspirin in secondary stroke prevention of cryptogenic stroke patients. No change was found in the primary outcome of stroke or mortality between treatment groups. However, a significant limitation of the study was the wide range of diagnostic inclusion criteria in the cryptogenic stroke cohort including ischaemic stroke patients in whom more than one plausible cause was found or incomplete diagnostic assessment had been performed.^{16–18} The breadth of underlying aetiology of disease arguably resulted in limited clinical value of the study and reinforced the need for further sub-categorisation of cryptogenic stroke patients.

Anticoagulation in Embolic Stroke of Unknown Source

The term ‘embolic stroke of unknown source’ (ESUS) has since been defined as a separate entity that requires a minimum diagnostic assessment to be performed prior to categorisation. Over 30% of ischaemic stroke patients are thought to meet this criteria.^{11,19}

The importance of this sub-category stems from the recognition that a sub-group of cryptogenic stroke patients in whom a primary embolic cause is suspected despite an unidentified source, may arguably benefit from anticoagulation. Support for this hypothesis comes from data showing that clots extracted from patients with cryptogenic stroke were similar in histology and composition to those with cardioembolic stroke.²⁰ However, real world data from two large randomised-control studies recently conducted, NAVIGATE ESUS (Rivaroxiban Versus Aspirin in Secondary Prevention of Stroke and Prevention of Systemic Embolism in Patients With Recent Embolic Stroke of Undetermined Source Trial) and RE-SPECT ESUS

(Randomised, Double-Blind, Evaluation in Secondary Stroke Prevention Comparing the Efficacy and Safety of the Oral Thrombin Inhibitor Dabigatran Etexilate Versus Acetylsalicylic Acid in Patients With Embolic Stroke of Undetermined Source Trial) failed to support this hypothesis.²¹ The NAVIGATE ESUS Trial was halted after an interim analysis of 7213 patients due to a higher risk of haemorrhage and no discernible difference in the primary outcome of secondary stroke or systemic embolism between patients randomised to Rivaroxaban or Aspirin.²¹ However, limitations of the study include the use of 15mg rivaroxaban rather than 20mg as recommended for stroke prophylaxis. Furthermore, only 20 hours of cardiac rhythm monitoring was required for inclusion and over 6 minutes of AF for diagnosis.²² The RE-SPECT ESUS Trial enrolled 5390 patients and also found no difference between the treatment groups of Dabigatran and Aspirin.^{23,24} However, a sub-group analysis suggested that Dabigatran may be effective for patients over 75 years of age, which may reflect the increasing incidence of undiagnosed AF within this sub-group.²⁴

Postulated limitations of these studies explaining the neutral results include the breadth of the current ESUS definition.²⁴ Furthermore, although clot histology in atherosclerotic plaque rupture and thrombus formation is similar to that of cardio-embolic strokes, current guidelines for secondary stroke prevention continue to advocate the use of antiplatelet agents over anticoagulation in this setting. The results of NAVIGATE-ESUS and RE-SPECT ESUS have both highlighted the need for further revision of the ESUS definition to identify those patients in whom anticoagulation would be of therapeutic benefit.

Anticoagulation in Embolic Stroke of Unknown Source with Evidence of Atrial Cardiomyopathy

Currently, there are two further ongoing trials: ARCADIA (The AtRial Cardiopathy and Antithrombotic Drugs In prevention After cryptogenic stroke randomized trial) and ATTICUS (Apixiban for Treatment of Embolic Stroke of Undetermined Source), both of which are large

randomised control studies assessing anticoagulation therapy and reduction in stroke recurrence in ESUS patients. Both trials further sub-categorise ESUS patients in to those with high-risk features suggestive of cardio-embolism^{12,25} (see *Table 1*). ARCADIA will recruit patients with specific biomarkers of atrial cardiomyopathy including elevated P-wave terminal force velocity in lead V₁, serum NT-proBNP and left atrial diameter on echocardiogram. The inclusion criteria for ATTICUS are broader and include non-major risk factors for cardio-embolism. As with NAVIGATE ESUS, a significant limitation of the ARCADIA study design is that the minimum period of cardiac rhythm monitoring required is 24 hours. Previous studies have suggested longer periods of monitoring are required for detection of AF.^{7,8} Although this study will likely include patients in whom AF would have been detected and therefore excluded following a longer period of monitoring, further analysis of this cohort may provide data regarding the impact of early anticoagulation which as yet remains unknown. All patients in the ATTICUS study will have an implantable loop recorder inserted and cross over to the Apixiban treatment arm should more than 2 minutes of AF be detected (minimum detection time required by Medtronic Linq). Additionally, the primary endpoint will be new ischaemic lesions identified on imaging rather than clinical stroke. The results of these studies will provide further evidence on the relevance of the ESUS sub-group in influencing treatment strategies.

The study design of both the ARCADIA and ATTICUS trials suggests that non-invasive markers of atrial cardiomyopathy may be the key to identifying specific subgroups of ischaemic stroke patients that could benefit from early anticoagulation.²⁴ Whilst many of these patients may go on to develop AF in the future, there may also be a cohort of individuals in whom biomarkers of atrial cardiomyopathy exists with elevated risk of stroke, but in the absence of AF.

3. Is there a Causal Relationship between AF and Stroke?

The association between ischaemic stroke and AF has been established in several studies with AF patients experiencing on average a 3-5 fold increased risk of stroke after adjustment for risk factors.²⁶⁻²⁸ Whilst there is clearly a strong association between the two with proven consistency amongst different cohorts, a true causal relationship has not been established as outlined below.¹³

Increased duration of atrial fibrillation does not increase stroke risk

Traditional theories suggest that during periods of AF, thromboembolic risk is increased due to ineffective atrial contraction leading to blood stasis and therefore, as per Virchow's triad, increased thromboembolic risk.¹³ In the long-term, left atrial remodelling occurs in the context of longstanding AF leading to atrial dilatation, fibrosis and increased blood stasis as a result of ineffective atrial contractility.²⁹⁻³¹

It has been theorised that these remodelling changes contribute to an elevated stroke risk, and therefore patients with persistent AF are at higher risk of thrombo-embolic stroke. This was demonstrated in the ROCKET-AF Trial (Rivaroxaban Once Daily Oral Direct Factor Xa Inhibition Compared with Vitamin K Antagonism for Prevention of Stroke and Embolism Trial in Atrial Fibrillation) where the adjusted rates of stroke or systemic embolism were 2.18 in the persistent AF group, compared with 1.73 in the paroxysmal AF group. However, other studies including the RE-LY Trial (Randomised Evaluation of Long Term Anticoagulant Therapy) found no significant change in thromboembolic risk between paroxysmal, persistent and permanent AF.³²⁻³⁶ Failure of trials to consistently demonstrate a clear relationship between AF duration and stroke risk suggests lack of causal effect between AF burden and ischaemic stroke.¹³

Moreover, remodelling changes begin to occur after 1-week of AF and progress over time, whereas several studies have demonstrated increased stroke risk with AF episodes of shorter duration suggesting that alternative/additional pathological processes increasing prothrombotic risk must be at play.^{32,36,37}

Embolic Stroke can occur in advance of Atrial Fibrillation

Inconsistencies in the temporal relationship between stroke and AF have also been demonstrated in previous studies (*Figure 2*). If AF alone increases the risk of thromboembolic disease, it stands to reason that AF should occur in advance of stroke. The ASSERT Trial (Asymptomatic Atrial Fibrillation and Stroke Evaluation in Pacemaker Patients and the Atrial Fibrillation Reduction Atrial Pacing) assessed patients with a pacemaker or implantable cardiac defibrillator device and reported a 2.5-fold increase in the risk of stroke or systemic embolism in those in whom sub-clinical AF was detected. However, of those who had a stroke, only 8% had documented episodes of sub-clinical atrial fibrillation in the month preceding their stroke and as many as 16% had episodes of sub-clinical AF detected *after* the event.³⁸ This was further supported by the IMPACT Trial (Combined Use of Biotronik Home Monitoring and Predefined Anticoagulation to Reduce Stroke Risk) where strong association was once again demonstrated but in the absence of a temporal relationship.³⁹ These findings indicate that whilst there is a reproducible association between stroke and AF, a history of AF is not necessarily a prerequisite of a thrombo-embolic stroke.

Restoration of sinus rhythm does not reduce the risk of ischaemic stroke

Finally, a causal relationship between AF and stroke would indicate that in the absence of the arrhythmia, a demonstrable reduction in thromboembolic stroke should be observed. A meta-analysis of rate versus rhythm control therapies in the management of AF showed the number of strokes to be unchanged regardless of strategy (OR 0.99, 95% CI 0.76-1.30), despite

reasonable success at maintaining sinus rhythm (OR 4.39, 95% CI 2.84-6.78).⁴⁰ Whilst these results were largely swayed by a single, large randomised control study, the AFFIRM trial, a further meta-analysis of thirteen studies compared long-term stroke rates following catheter ablation with anti-arrhythmic medication and found no evidence of reduced stroke risk.⁴¹ This was evident despite improved effectiveness in maintaining sinus rhythm following catheter ablation compared with anti-arrhythmic medication with 445 out of 855 patients crossing over from the medical to interventional arms of the studies due to drug failure or intolerance.⁴¹

These findings may be explained by atrial structural remodelling that exists even in the presence of adequate rhythm control, however a previous large retrospective study suggests that risk factors for stroke may be different following catheter ablation and alternative risk scores have been proposed that may improve stroke-risk stratification after catheter ablation treatment.⁴²

Can ischaemic stroke cause AF?

Although less commonly explored, stroke has been hypothesised to cause AF due to autonomic nervous system activation leading to changes in cardiac electrophysiology providing a substrate for atrial arrhythmias.^{13,43} Previous clinical studies have observed a higher incidence of AF in the early post-stroke period that may support this hypothesis.^{8,44} However, patients in these studies had no prior heart rhythm monitoring for AF detection. Furthermore, the ASSERT trial observed a median time of 101 days prior to AF detection, suggesting that a proportion of patients within these studies may have had undiagnosed AF that preceded the stroke event.

The inconsistencies described above add further weight to the emerging concept that the relationship between AF and stroke may represent association rather than causation and

therefore consideration must be given to the implications of this on therapeutic considerations for stroke prevention.

4. Limitations of Current Treatment Strategies

Whilst the AF-stroke relationship remains incompletely understood, meta-analyses have rigorously proven anticoagulation therapy with vitamin-K antagonists or direct oral anticoagulants in AF patients significantly reduces the risk of stroke.^{45,46} However, a delay in the detection of AF leads to further delays in initiation of anticoagulation therapy in a high risk cohort. Previous studies show a reduction in thromboembolic stroke risk by as much as 60% following anticoagulation therapy, compared with 20% whilst on an antiplatelet agent, with the risk of stroke recurrence being highest in the initial 12 months post event.^{46,47} A large proportion of ESUS patients are suspected to have undiagnosed AF, yet do not receive anticoagulation.⁷ With stroke recurrence rates as high as 30% in this cohort, it is conceivable that a proportion of these strokes are preventable with a change in treatment strategy tailored to the individual.

The limitations of the currently accepted AF-stroke model have several implications pertaining to therapy. First, it relies on an episode of AF being detected prior to initiation of anticoagulation therapy. The temporal dissociation described above suggests that the risk of stroke associated with AF may occur distant from the AF episode.⁴⁸ Even with an implantable loop recorder in situ, AF may only be detected in around 12% of patients with cryptogenic stroke over a 1-year period.⁸ With historical studies demonstrating a reduction in secondary stroke risk from 12% to 4% per year (HR 0.34; 95% CI 0.36-0.79) in patients randomised to receive anticoagulation or placebo post-stroke, the risks of delaying anticoagulation in this cohort of patients are likely to be significant.⁴⁹ This observation was recently supported in the Early Rhythm Control Therapy in Patients with Atrial Fibrillation Study (EAST-AFNET 4)

where lower stroke rates were documented in patients receiving early rhythm control compared to usual care (treatment effect HR 0.65; 95% CI 0.44-0.97).⁵⁰

Furthermore, a proportion of patients may never develop AF but remain at high risk of embolic stroke due to underlying atrial disease in keeping with an atrial cardiomyopathy. With current risk stratification models requiring a diagnosis of AF prior to initiation of anticoagulation therapy, this cohort do not currently meet the criteria for treatment.

5. Atrial Biomarkers for Risk Stratification of ESUS Patients

The CHA₂DS₂VASc scoring system is unanimously the risk stratification tool of choice to identify main predictors of stroke in the context of AF.⁵¹ AF patients with no identifiable risk factors are at only a marginally increased risk of stroke compared with patients without AF.⁵² It is interesting to note that anticoagulation guidance has little focus on the clinical characteristics of atrial fibrillation such as arrhythmia duration, sustained rapid ventricular response or arrhythmia persistence, but rather on co-morbidities associated with a cardiomyopathic state.

Recently, alternative biomarkers associated with an atrial cardiomyopathy have been associated with the risk of stroke independent of AF. Identification of such biomarkers, as outlined below, may represent a new approach for risk stratification in prevention of ischaemic stroke, providing anticoagulation guidance in the absence of documented AF.

5.1 Cardiac Imaging Biomarkers:

Atrial Fibrosis:

Atrial fibrosis has been identified as a potential precursor leading to the initiation and perpetuation of AF. Fibrosis occurs as a result of excessive deposition of extracellular matrix predominantly containing collagen I, collagen III and fibronectin.⁵³ Atrial structural

remodelling at the tissue level comprises myocyte apoptosis, alterations in cellular metabolism and accumulation of glycogen.^{54,55} Together, these changes predispose to alterations in the orientation of cardiac myocytes, re-distribution of gap junctions and therefore velocity of electrical conduction, giving rise to electrical conduction delay and atrial substrate for sustaining AF. Furthermore, a thrombogenic substrate is generated due to increased inflammation, endothelial dysfunction and blood stasis.⁵⁴ Together, these changes could lead to the development of a procoagulant state resulting in increased risk of thromboembolism and clinical stroke. Atrial structural remodelling could be caused by AF, but may also be promoted by other factors such as age, hypertension, diabetes, inflammation and heart failure suggesting that stroke may be caused by both AF and/or abnormal atrial substrate.^{54,56} Atrial fibrosis may therefore precede AF, or be present even in the absence of AF.

In order to use the extent of left atrial fibrosis as a means of stroke risk stratification and a guide for anticoagulation therapy, a biological relationship in which severity of fibrosis is directly related to stroke risk needs to be demonstrated in order to prove causation. Indeed, accumulating evidence suggests that such a relationship may exist. An observational, case-control study of 111 patients detected a statistically significant increase in left atrial fibrosis of patients with cryptogenic stroke compared with stroke of other specific causes (excluding AF).^{57,58} The left atrial phenotype of those with cryptogenic stroke appeared to be an intermediate between those with AF and other specific causes for stroke, possibly representing earlier detection of an atrial cardiomyopathy that may ultimately progress to AF with disease progression.^{57,58} The process of disease progression is further supported in histological studies where greater fibrosis has been demonstrated in autopsies of patients with permanent AF compared with paroxysmal AF, although the extent of fibrosis was not found to correlate with the duration of AF history.⁵⁹

Atrial cardiac magnetic resonance imaging is currently the gold standard tool for non-invasive atrial tissue characterisation and has been used in the quantification of atrial fibrosis by identifying areas of late gadolinium enhancement (LGE) in the atrial wall (*Figure 3*).⁶⁰ However, there are a number of limitations of atrial late gadolinium enhancement assessment which could hinder its use as a risk assessment tool. Firstly, cardiac magnetic resonance imaging is expensive and less readily available compared with alternative imaging techniques such as echocardiography. Secondly, accurate quantification of atrial fibrosis is dependent on the image quality acquired. There is currently a large variability in image acquisition parameters, image analysis and post-processing tools across different centres.⁶¹

Atrial Enlargement and Geometry:

A susceptibility to atrial arrhythmias has been demonstrated during abnormal myocardial loading that results in atrial stretch and dilatation, playing an important role in the pathogenesis of AF.³⁷ During AF, uncontrolled ventricular rates can lead to increased haemodynamic overload and consequently cardiac failure, providing an underlying mechanism for atrial dilatation. However, co-morbidities such as hypertension, heart failure or mitral valve disease also contribute to atrial dilatation, increasing the vulnerability of the left atrium to AF. Ex-vivo studies in guinea-pig hearts demonstrated electrophysiological changes in atrial tissue following increased left atrial volume.⁶² A further study in Langendorff-perfused rabbit hearts showed a graded increase in AF vulnerability that correlated with increasing atrial pressures.⁶³ Later, in-vitro studies of atrial stretch have shown downstream effects of hypertrophy and apoptosis that feed into the structural and electrophysiological changes described above. Atrial dilatation is related to both the severity of the underlying disease as well as the AF burden and may therefore precede AF manifestation in the context of other risk factors.³⁷

The relationship between atrial size and stroke risk has been controversial, in large part due to the significant limitations of studies that were conducted in patients with atrial fibrillation or

mitral valve disease. A sub-group analysis of the Framingham Study assessed left atrial size and stroke risk in all patients over the age of 50 years.⁶⁴ After multivariable adjustment, left atrial size was found to be an independent predictor of stroke with every 10mm increase in left atrial size resulting in a relative risk ratio of 2.4 for men (95% CI, 1.6-3.7) and 1.4 for women (95% CI, 0.9-2.1).^{64,65} Further population based studies in the Northern Manhattan Stroke Study assessed the role of atrial enlargement as an independent risk factor for recurrent ischaemic stroke in a multi-ethnic population.^{65,66} Moderate to severe left atrial enlargement was demonstrated in patients with a recurrent cardioembolic or cryptogenic stroke with a hazard ratio of 2.83 after multivariable adjustment (95% CI, 1.03-7.81). These findings were attributed to reduced flow velocity in the left atrial appendage or left atrial enlargement representing a marker of end-organ damage from hypertension, which is a known risk factor for recurrent stroke.

Echocardiography is the most common modality of assessing atrial size and has dominated previous studies, however a significant limitation is that it assumes a spherical left atrial shape, therefore frequently underestimating left atrial volume.⁶⁷⁻⁷⁰ Improved image quality with cardiac magnetic resonance imaging has since increased our understanding of atrial shape.⁷¹ The left atrium can often be asymmetrical and AF disease progression frequently results in geometrical changes such as sphericity, trapezoidal deformation or alterations in roof shape.⁷²⁻

74

Left Atrial Appendage:

The majority of left atrial thrombi have been reported to originate from the left atrial appendage.⁷⁵⁻⁷⁷ Furthermore, trials of left atrial appendage closure were non-inferior to warfarin in reducing the risk of ischaemic stroke, highlighting the significance of left atrial appendage in stroke risk.^{78,79}

Non-chicken wing left atrial appendage morphologies including cactus, windsock and cauliflower have been shown in some trials to increase stroke risk due to more extensive left atrial appendage trabeculations and smaller left atrial appendage orifice diameter.^{80,81} However, a key limitation of these studies has often been inter-observer reproducibility with one study having found left atrial appendage morphology agreement to be 58.82%. To date, classification remains subjective and there are no automated or semi-automated algorithms available between centres to ensure reproducibility.⁸⁰ In addition, left atrial appendage morphology has only been investigated in AF patients and the significance of morphology in the absence of AF remains unknown.

A cross-sectional study also demonstrated reduced left atrial appendage flow velocity on echocardiography between AF and non-AF patients, which may increase blood pooling and stasis leading to clot formation.⁸²⁻⁸⁴ Whilst further studies are required to understand the relevance of left atrial appendage morphology in the absence of AF, there is some evidence that left atrial appendage flow may represent a biomarker of recurrent ischaemic stroke risk.

5.2 Electrocardiographic Biomarkers:

P-wave terminal force in lead V₁

P-wave terminal force in lead V₁ (PTFV₁) is defined as the duration (in seconds) of the negative terminal deflection of the p-wave in lead V₁ multiplied by the absolute value of its amplitude (in mm) and is considered a marker of left atrial abnormality.^{85,86} Several studies have shown elevated P-wave terminal force in lead V₁ to be associated with atrial disease including dilatation, hypertrophy, fibrosis and abnormal inter-atrial conduction, all of which are consistent with structural and electrical remodelling changes that occur in atrial cardiomyopathy.⁸⁷ A 2-year study of 295 consecutive ischaemic stroke patients, found P-wave terminal force in lead V₁ to be a strong predictor of paroxysmal AF in ischaemic stroke patients

and noted a cut-off value of 0.04mm/s for paroxysmal AF prediction (sensitivity of 80% and specificity of 72.2%). P-wave terminal force in lead V₁ was argued to be a cheap and practical method for selecting patients that require more extensive ECG monitoring for the detection of AF in ischaemic stroke patients.⁸⁵ Abnormal P-wave terminal force in lead V₁ has also been shown to independently correlate with diffuse left ventricular fibrosis quantified on CMR and measured as a surrogate for atrial fibrosis.⁸⁸

Evidence of a correlation between P-wave terminal force in lead V₁ with ischaemic stroke independent of AF dates back as far as 2005.⁸⁹ More recently the Atherosclerosis Risk in Communities (ARIC) Study provided further evidence of an association between P-wave terminal force in lead V₁ as both an AF predictor and incident ischaemic stroke.⁹⁰ These findings have been further supported in the Multi-Ethnic Study of Atherosclerosis, Atherosclerosis Risk in Communities study and the Northern Manhattan Study which both showed an association between PTFV₁ and ischaemic stroke independent of AF.^{91,92}

Atrial Ectopy

High atrial ectopic burden has also been associated with an increased risk of developing AF⁹³ and a recent single centre case-control study of 2800 patients found the mean premature atrial complex count on 24-hour monitoring to be 426 in the ischaemic stroke group compared with 105 in the control group (p value <0.001).⁹⁴ This was supported by the Copenhagen Holter Study cohort in which excessive atrial ectopic activity (defined as either more than 30 atrial ectopics per hour daily or runs of over 20 atrial ectopics) was associated with an increased stroke risk even in the absence of electrocardiographic evidence of AF.⁹⁵ A possible mechanism could be atrial ectopy occurring as a precursor to AF. Alternatively, atrial ectopy may be a marker of other risk factors such as hypertension, diabetes or structural remodelling leading to atrial fibrosis and endothelial dysfunction resulting in a hypercoagulable state.⁹⁵

Prolongation of PR interval

Although the PR interval displays circadian variation,⁹⁶ population-based studies have shown prolongation of the electrocardiographic PR interval to be associated with a two-fold increased risk of AF and up to five-fold in those in the upper 95th centile after adjusting for risk factors.^{97,98} PR prolongation could be a manifestation of initial conduction disease occurring as a result of structural cardiac abnormalities and therefore a further biomarker of atrial disease.^{99,100} Alternatively, PR prolongation can be associated with other conduction abnormalities or a sign of slow intra-atrial or inter-atrial conduction providing the atrial mechanism to sustain AF.¹⁰¹ Both a large multi-centre retrospective trial and the CRYSTAL-AF Trial (Cryptogenic Stroke and underlying Atrial Fibrillation) demonstrated a strong association between PR prolongation and increased AF incidence in cryptogenic stroke.^{98,102}

P-wave terminal force in lead V₁, atrial ectopy and PR prolongation have all demonstrated an association with ischaemic stroke in the absence of AF and which may be attributed to an underlying atrial cardiomyopathy. However, despite these associations, predictive models that have implemented these criteria, such as the Framingham AF Score, have displayed moderate or low predictive capacity.^{98,103} These findings highlight the difficulty in application of large population trends on an individual patient level.

5.3 Serum Biomarkers

NT-proBNP (N-terminal pro-brain natriuretic peptide)

NT-proBNP is a neurohormone released by the myocardium in response to increased wall tension. NT-proBNP can be elevated in a number of conditions causing haemodynamic stress including heart failure, AF and structural heart disease.¹⁰⁴ Previous studies have demonstrated elevated NT-proBNP in matched AF patients compared with sinus rhythm as well as a rapid reduction in levels on restoration of sinus rhythm.^{105,106} Furthermore, both the RE-LY sub-

study and later the ARISTOTLE study found the risk of stroke or systemic thromboembolism to be independently related to NT-proBNP in AF patients and more than doubled in the highest compared with the lowest NT-proBNP quartile groups, suggesting there may be a role for inclusion of NT-proBNP in stratifying future stroke.^{107,108} Additional studies advocate use of NT-proBNP levels in the detection of new atrial fibrillation following cryptogenic stroke, however it may not add additional diagnostic information over and above traditional non-invasive methods for AF detection.^{109,110}

Troponin

Troponin is a cardiac enzyme most commonly used as a biomarker in the detection of myocardial infarction or injury but can also be serum detectable at low levels in patients with heart failure, structural heart disease, stable coronary artery disease, renal failure and increasing age. The RE-LY sub-study and ARISTOTLE studies demonstrated rates of stroke were independently related to higher levels of Troponin I. The RE-LY sub-study results further indicated that addition of Troponin I levels to a predictive model for stroke outcomes provided additional prognostic information.

Further studies are required to incorporate serum biomarkers in predictive models of future stroke risk following embolic stroke of unknown source to determine their significance and implications on future anticoagulation.

6. Atrial Cardiomyopathy

The term atrial cardiomyopathy has been defined as ‘any complex of structural, architectural, contractile or electrophysiological changes affecting the atria with the potential to produce clinically relevant manifestations.’ Its clinical value may lie in its role as a precursor to AF, predicting those likely to develop AF as well as allowing substrate classification that may inform individualised patient care strategies. Furthermore, it may define a cohort of patients

with hypercoagulability and at risk of thromboembolic disease (*see Table 2*). Several intra-atrial sampling studies assessing markers of hypercoagulability have shown a statistical difference in marker levels between AF patients and non-AF patients.^{111,112} However, there are no studies as yet that have assessed whether a procoagulant state predates the onset of detectable AF. Furthermore, a limitation of these studies has been the inability to distinguish hypercoagulability secondary to concomitant co-morbidities such as vascular disease or directly attributable to AF.^{113,114}

The EHRA/HRS expert consensus on atrial cardiomyopathy proposes a histological classification of atrial cardiomyopathy (EHRA Class I-IV), emphasising it to be a descriptive classification rather than a pathway of disease progression.¹¹⁵ The classification allows discrete cohorts of atrial cardiomyopathy to be studied further to validate the classification and understand its clinical relevance with regards to individualised therapy for AF patients.

A cross-sectional pilot study of the New York Columbia Collaborative Specialized Program of Translational Research in Acute Stroke registry (NYCC SPOTRIAS) found that 63% of patients had at least one biomarker of left atrial cardiomyopathy (including severe left atrial enlargement, elevated NT-proBNP or elevated PTFV₁). However, the sample size was too small to draw conclusions regarding clinical significance as only 4 patients developed paroxysmal AF after their stroke, of which 2 had evidence of atrial cardiomyopathy as per the classification.

7. Limitations

Several limitations exist in the development of novel risk scores. Firstly, there is significant overlap between several biomarkers which can be difficult to separate.⁵⁷ For example, elevated BNP, a biomarker of atrial wall stress, has been found to be elevated in patients with atrial dilatation.³ Similarly, P-wave terminal force in lead V₁ is a measure of left atrial

activation and has historically been used as a marker of left atrial dysfunction. Other biomarkers such as left atrial size, geometry or fibrosis may also affect the P-wave terminal force in lead V₁ and the risk score therefore compounded. Additionally, further studies are required to assess the comparable significance of each biomarker to understand its individual relevance within the risk score.⁵⁷ These limitations are significant in that the study design of recent and ongoing trials use variable definitions of atrial cardiomyopathy creating difficulty when comparing study outcomes.

A further limitation is the long-term follow-up required for AF detection in the ESUS group. It is well documented that the incidence of AF and atrial disease increases with advanced age presenting further challenges in differentiating ESUS patients at increased risk of developing AF. Furthermore, AF detection post-stroke is affected by a number of external non-cardiac confounders, as yet unaccounted for in many of the risk scores, such as smoking, air-pollution and obesity. These factors likely have increased weighting over time, making it increasingly difficult to identify meaningful risk factors.

With this in mind, it is certainly worth considering recent technological advances and whether there may be a role for the use machine learning in further defining risk factors. However, it is imperative that models are built using high-quality data and appropriate machine learning methodology to ensure reproducibility and clinical utility.

8. Conclusion

The relationship between AF and ischaemic stroke is complex. Atrial cardiomyopathy may be an underlying mechanism for embolic stroke of unknown source but requires further research to create definitions of clinical significance. Imaging and electrocardiographic biomarkers of atrial cardiomyopathy in the absence of AF show promise in observational data (see Table 2), but studies implementing these biomarkers within risk stratification scores have, to date, lacked

sensitivity and specificity highlighting a need for improved risk stratification models. In addition, large randomised studies assessing anticoagulation in the ESUS group have not shown statistically significant benefit as yet, perhaps due to the further need to better identify patients at high risk of stroke recurrence. Ongoing studies, including ARCADIA and ATTICUS, will assess the clinical significance of ESUS in the context of underlying structural or electrical atrial disease. If positive, these studies could lead to the design of randomised trials where anticoagulation therapy is determined not by the presence or absence of AF but by the presence or absence of the underlying atrial cardiomyopathy. This development would result in a step-change in the clinical management of ESUS patients and should be an area of research priority over the coming years.

References

1. Virani SS, Alonso A, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP, Chamberlain AM, Chang AR, Cheng S, Delling FN, et al. Heart Disease and Stroke Statistics—2020 Update. *Circulation*. 2020. 139–596 p.
2. Gladstone DJ, Bui E, Fang J, Laupacis A, Lindsay MP, Tu J V., Silver FL, Kapral MK. Potentially preventable strokes in high-risk patients with atrial fibrillation who are not adequately anticoagulated. *Stroke*. 2009 10.1161/STROKEAHA.108.516344;40(1):235–40.
3. Saposnik G, Gladstone D, Raptis R, Zhou L, Hart RG. Atrial fibrillation in ischemic stroke: Predicting response to thrombolysis and clinical outcomes. *Stroke*. 2013 10.1161/STROKEAHA.112.676551;44(1):99–104.
4. Hindricks G, Potpara T, Dagres N, Arbelo E, Bax JJ, Blomström-Lundqvist C, Boriani G, Castella M, Dan G-A, Dilaveris PE, et al. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association of Cardio-Thoracic Surgery (EACTS). *Eur Heart J* [Internet]. 2020 Aug 29 10.1093/eurheartj/ehaa612 32860505;1–126. Available from: 10.1093/eurheartj/ehaa612
5. Camm AJ, Corbucci G, Padeletti L. Usefulness of continuous electrocardiographic monitoring for atrial fibrillation. *Am J Cardiol* [Internet]. 2012 10.1016/j.amjcard.2012.03.021;110(2):270–6. Available from: 10.1016/j.amjcard.2012.03.021
6. Healey JS, Connolly SJ, Gold MR, Israel CW, Van Gelder IC, Capucci A, Lau CP, Fain E, Yang S, Bailleul C, et al. Subclinical atrial fibrillation and the risk of stroke. *N Engl J Med*. 2012 10.1056/NEJMoa1105575;366(2):120–9.

7. Gladstone DJ, Spring M, Dorian P, Panzov V, Thorpe KE, Hall J, Vaid H, O'Donnell M, Laupacis A, Côté R, et al. Atrial fibrillation in patients with cryptogenic stroke. *N Engl J Med*. 2014 10.1056/NEJMoa1311376;370(26):2467–77.
8. Sanna T, Diener HC, Passman RS, Di Lazzaro V, Bernstein RA, Morillo CA, Rymer MM, Thijs V, Rogers T, Beckers F, et al. Cryptogenic stroke and underlying atrial fibrillation. *N Engl J Med*. 2014 10.1056/NEJMoa1313600;370(26):2478–86.
9. Afzal M, Kanmanthareddy A, Gunda S, Atkins D, Reddy M, Atoui M, Pillarisetti J, Dawn B, Lakkireddy D. Cryptogenic Stroke and Underlying Atrial Fibrillation: a Systematic Review and Meta-Analysis of Randomized Control Trials. *J Am Coll Cardiol* [Internet]. 2015 10.1016/s0735-1097(15)60360-6;65(10):A360. Available from: 10.1016/s0735-1097(15)60360-6
10. Steinberg JS, O'Connell H, Li S, Ziegler PD. Thirty-Second Gold Standard Definition of Atrial Fibrillation and Its Relationship with Subsequent Arrhythmia Patterns: Analysis of a Large Prospective Device Database. *Circ Arrhythmia Electrophysiol*. 2018 10.1161/CIRCEP.118.006274 30002065;11(7):1–9.
11. Hart RG, Diener HC, Coutts SB, Easton JD, Granger CB, O'Donnell MJ, Sacco RL, Connolly SJ. Embolic strokes of undetermined source: The case for a new clinical construct. *Lancet Neurol* [Internet]. 2014 10.1016/S1474-4422(13)70310-7;13(4):429–38. Available from: 10.1016/S1474-4422(13)70310-7
12. Kamel H, Longstreth WT, Tirschwell DL, Kronmal RA, Broderick JP, Palesch YY, Meinzer C, Dillon C, Ewing I, Spilker JA, et al. The AtRial Cardiopathy and Antithrombotic Drugs In prevention After cryptogenic stroke randomized trial: Rationale and methods. *Int J Stroke*. 2019 10.1177/1747493018799981;14(2):207–14.
13. Kamel H, Okin PM, Elkind MSV, Iadecola C. Atrial Fibrillation and Mechanisms of

- Stroke: Time for a New Model. *Stroke*. 2016 10.1161/STROKEAHA.115.012004 26786114;47(3):895–900.
14. Okin PM. P-wave morphology and the risk of incident ischemic stroke in the Multi-Ethnic Study of Atherosclerosis. *Stroke*. 2014 10.1161/STROKEAHA.114.006364;45(9):2786–8.
 15. Mahnkopf C, Badger TJ, Burgon NS, Daccarett M, Haslam TS, Badger CT, McGann CJ, Akoum N, Kholmovski E, MacLeod RS, et al. Evaluation of the left atrial substrate in patients with lone atrial fibrillation using delayed-enhanced MRI: Implications for disease progression and response to catheter ablation. *Heart Rhythm*. 2010 10.1016/j.hrthm.2010.06.030;7(10):1475–81.
 16. Adams H., Bendixen B., Kappelle L., Biller J, Love B., Gordon D., Marsh E. Classification of Subtype of Acute Ischemic Stroke. *Stroke*. 1993 10.1161/01.STR.24.1.35 7678184;23(1):35–41.
 17. Sacco RL, Prabhakaran S, Thompson JLP, Murphy A, Sciacca RR, Levin B, Mohr JP. Comparison of warfarin versus aspirin for the prevention of recurrent stroke or death: Subgroup analyses from the warfarin-aspirin recurrent stroke study. *Cerebrovasc Dis*. 2006 10.1159/000092331;22(1):4–12.
 18. Mohr JP, Thompson JLP, Lazar RM, Levin B, Sacco RL, Furie KL, Kistler JP, Albers GW, Pettigrew LC, Adams HP, et al. A Comparison of Warfarin and Aspirin for the Prevention of Recurrent Ischemic Stroke. *N Engl J Med* [Internet]. 2001 Nov 15 10.1056/NEJMoa011258 11794192;345(20):1444–51. Available from: 10.1056/NEJMoa011258
 19. Santamarina E, Penalba A, Garcia-Berrocso T, Delgado P, Quintana M, González-Alujas T, Ribo M, Maisterra O, Molina CA, Evangelista A, et al. Biomarker level

- improves the diagnosis of embolic source in ischemic stroke of unknown origin. *J Neurol*. 2012 10.1007/s00415-012-6532-4;259(12):2538–45.
20. Boeckh-Behrens T, Kleine JF, Zimmer C, Neff F, Scheipl F, Pelisek J, Schirmer L, Nguyen K, Karatas D, Poppert H. Thrombus histology suggests cardioembolic cause in cryptogenic stroke. *Stroke*. 2016 10.1161/STROKEAHA.116.013105;47(7):1864–71.
 21. Hart RG, Sharma M, Mundl H, Kasner SE, Bangdiwala SI, Berkowitz SD, Swaminathan B, Lavados P, Wang Y, Wang Y, et al. Rivaroxaban for stroke prevention after embolic stroke of undetermined source. *N Engl J Med*. 2018 10.1056/NEJMoa1802686 29766772;378(23):2191–201.
 22. Hart RG, Sharma M, Mundl H, Shoamanesh A, Kasner SE, Berkowitz SD, Pare G, Kirsch B, Pogue J, Pater C, et al. Rivaroxaban for secondary stroke prevention in patients with embolic strokes of undetermined source: Design of the NAVIGATE ESUS randomized trial. *Eur Stroke J*. 2016 10.1177/2396987316663049;1(3):146–54.
 23. Diener HC, Sacco RL, Donald Easton J, Granger CB, Bernstein RA, Uchiyama S, Kreuzer J, Cronin L, Cotton D, Grauer C, et al. Dabigatran for prevention of stroke after embolic stroke of undetermined source. *N Engl J Med*. 2019 10.1056/NEJMoa1813959 31091372;380(20):1906–17.
 24. Paciaroni M, Kamel H. Do the Results of RE-SPECT ESUS Call for a Revision of the Embolic Stroke of Undetermined Source Definition? *Stroke* [Internet]. 2019 Apr 10.1161/STROKEAHA.118.024160;50(4):1032–3. Available from: 10.1161/STROKEAHA.118.024160
 25. Geisler T, Poli S, Meisner C, Schreieck J, Zuern CS, Nägele T, Brachmann J, Jung W, Gahn G, Schmid E, et al. Apixaban for treatment of embolic stroke of undetermined source (ATTICUS randomized trial): Rationale and study design. *Int J Stroke*. 2017

- 10.1177/1747493016681019;12(9):985–90.
26. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: The framingham study. *Stroke*. 1991 10.1161/01.STR.22.8.983 1866765;22(8):983–8.
 27. Wolf PA, Dawber TR, Thomas HE, Kannel WB. Epidemiologic assessment of chronic atrial fibrillation and risk of stroke: The framingham study. *Neurology*. 1978 10.1212/wnl.28.10.973 570666;28(10):973–7.
 28. Freedman B, Potpara TS, Lip GYH. Stroke prevention in atrial fibrillation. *Lancet* [Internet]. 2016 10.1016/S0140-6736(16)31257-0;388(10046):806–17. Available from: 10.1016/S0140-6736(16)31257-0
 29. Zatuchni J, Sanfilippo AJ, Weyman AE. Atrial enlargement as a consequence of atrial fibrillation [1]. *Circulation*. 1991 10.1161/circ.83.4.1826477 1826477;83(4):1458.
 30. Mihm MJ, Yu F, Carnes CA, Reiser PJ, McCarthy PM, Wagoner DR Van, Bauer JA. During Human Atrial Fibrillation. 2015;174–81.
 31. Schotten U, Ausma J, Stellbrink C, Sabatschus I, Vogel M, Frechen D, Schoendube F, Hanrath P, Allessie MA. Cellular mechanisms of depressed atrial contractility in patients with chronic atrial fibrillation. *Circulation*. 2001 10.1161/01.CIR.103.5.691 11156881;103(5):691–8.
 32. Steinberg BA, Hellkamp AS, Lokhnygina Y, Patel MR, Breithardt G, Hankey GJ, Becker RC, Singer DE, Halperin JL, Hacke W, et al. Higher risk of death and stroke in patients with persistent vs. paroxysmal atrial fibrillation: Results from the ROCKET-AF Trial. *Eur Heart J*. 2015 10.1093/eurheartj/ehu359;36(5):288–96.
 33. Disertori M, Franzosi MG, Barlera S, Cosmi F, Quintarelli S, Favero C, Cappellini G, Fabbri G, Maggioni AP, Staszewsky L, et al. Thromboembolic event rate in paroxysmal

- and persistent atrial fibrillation: Data from the GISSI-AF trial. *BMC Cardiovasc Disord.* 2013 10.1186/1471-2261-13-28;13.
34. Hohnloser SH, Pajitnev D, Pogue J, Healey JS, Pfeffer MA, Yusuf S, Connolly SJ. Incidence of Stroke in Paroxysmal Versus Sustained Atrial Fibrillation in Patients Taking Oral Anticoagulation or Combined Antiplatelet Therapy. An ACTIVE W Substudy. *J Am Coll Cardiol.* 2007 10.1016/j.jacc.2007.07.076;50(22):2156–61.
 35. Nieuwlaat R, Prins MH, Le Heuzey JY, Vardas PE, Aliot E, Santini M, Cobbe SM, Widdershoven JWMG, Baur LH, Lévy S, et al. Prognosis, disease progression, and treatment of atrial fibrillation patients during 1 year: Follow-up of the Euro Heart Survey on Atrial Fibrillation. *Eur Heart J.* 2008 10.1093/eurheartj/ehn139;29(9):1181–9.
 36. Flaker G, Ezekowitz M, Yusuf S, Wallentin L, Noack H, Brueckmann M, Reilly P, Hohnloser SH, Connolly S. Efficacy and safety of dabigatran compared to warfarin in patients with paroxysmal, persistent, and permanent atrial fibrillation: Results from the RE-LY (Randomized Evaluation of Long-Term Anticoagulation Therapy) study. *J Am Coll Cardiol [Internet].* 2012 10.1016/j.jacc.2011.10.896;59(9):854–5. Available from: 10.1016/j.jacc.2011.10.896
 37. De Jong AM, Maass AH, Oberdorf-Maass SU, Van Veldhuisen DJ, Van Gilst WH, Van Gelder IC. Mechanisms of atrial structural changes caused by stretch occurring before and during early atrial fibrillation. *Cardiovasc Res.* 2011 10.1093/cvr/cvq357;89(4):754–65.
 38. Brambatti M, Connolly SJ, Gold MR, Morillo CA, Capucci A, Muto C, Lau CP, Van Gelder IC, Hohnloser SH, Carlson M, et al. Temporal relationship between subclinical atrial fibrillation and embolic events. *Circulation.* 2014 10.1161/CIRCULATIONAHA.113.007825;129(21):2094–9.

39. Martin DT, Bersohn MM, L.waldo A, Wathen MS, Choucair WK, Lip GYH, Ip J, Holcomb R, Akar JG, Halperin JL. Randomized trial of atrial arrhythmia monitoring to guide anticoagulation in patients with implanted defibrillator and cardiac resynchronization devices. *Eur Heart J*. 2015 10.1093/eurheartj/ehv115;36(26):1660–8.
40. Al-Khatib SM, Allen LaPointe NM, Chatterjee R, Crowley MJ, Dupre ME, Kong DF, Lopes RD, Povsic TJ, Raju SS, Shah B, et al. Rate- and Rhythm-Control Therapies in Patients With Atrial Fibrillation. *Ann Intern Med* [Internet]. 2014 Jun 3 10.7326/M13-1467;160(11):760. Available from: 10.7326/M13-1467
41. Zheng YR, Chen ZY, Ye LF, Wang LH. Long-term stroke rates after catheter ablation or antiarrhythmic drug therapy for atrial fibrillation: A meta-analysis of randomized trials. *J Geriatr Cardiol*. 2015 10.11909/j.issn.1671-5411.2015.05.012;12(5):507–14.
42. Kim YG, Shim J, Oh SK, Lee KN, Choi J Il, Kim YH. Risk Factors for Ischemic Stroke in Atrial Fibrillation Patients Undergoing Radiofrequency Catheter Ablation. *Sci Rep*. 2019 10.1038/s41598-019-43566-z 31065030;9(1):1–8.
43. Chen PS, Chen LS, Fishbein MC, Lin SF, Nattel S. Role of the autonomic nervous system in atrial fibrillation: Pathophysiology and therapy. *Circ Res*. 2014 10.1161/CIRCRESAHA.114.303772;114(9):1500–15.
44. Cotter PE, Martin MPJ, Ring L, Warburton EA, Belham M, Pugh PJ. Incidence of atrial fibrillation detected by implantable loop recorders in unexplained stroke. *Neurology*. 2013 10.1212/WNL.0b013e31828f1828 23535493;80(17):1546–50.
45. López-López JA, Sterne JAC, Thom HHZ, Higgins JPT, Hingorani AD, Okoli GN, Davies PA, Bodalia PN, Bryden PA, Welton NJ, et al. Oral anticoagulants for prevention of stroke in atrial fibrillation: Systematic review, network meta-Analysis, and cost effectiveness analysis. *BMJ*. 2017 10.1136/bmj.j5058;359.

46. Hart RG, Pearce LA, Aguilar MI. Meta-analysis: Antithrombotic therapy to prevent stroke in patients who have nonvalvular atrial fibrillation. *Ann Intern Med.* 2007 10.7326/0003-4819-146-12-200706190-00007 17577005;146(12):857–67.
47. Amarenco P, Lavallée PC, Labreuche J, Albers GW, Bornstein NM, Canhão P, Caplan LR, Donnan GA, Ferro JM, Hennerici MG, et al. One-year risk of stroke after transient ischemic attack or minor stroke. *N Engl J Med.* 2016 10.1056/NEJMoa1412981;374(16):1533–42.
48. Turakhia MP, Ziegler PD, Schmitt SK, Chang Y, Fan J, Than CT, Keung EK, Singer DE. Atrial Fibrillation Burden and Short-Term Risk of Stroke: Case-Crossover Analysis of Continuously Recorded Heart Rhythm from Cardiac Electronic Implanted Devices. *Circ Arrhythmia Electrophysiol.* 2015 10.1161/CIRCEP.114.003057;8(5):1040–7.
49. European Atrial Fibrillation Trial Study. Secondary prevention in non-rheumatic atrial fibrillation after transient ischaemic attack or minor stroke. *Lancet.* 1993;342:1255–62.
50. Kirchhof P, Camm AJ, Goette A, Brandes A, Eckardt L, Elvan A, Fetsch T, van Gelder IC, Haase D, Haegeli LM, et al. Early Rhythm-Control Therapy in Patients with Atrial Fibrillation. *N Engl J Med.* 2020 10.1056/nejmoa2019422 32865375;383(14):1305–16.
51. Alkhouli M, Friedman PA. Ischemic Stroke Risk in Patients With Nonvalvular Atrial Fibrillation: JACC Review Topic of the Week. *J Am Coll Cardiol.* 2019 10.1016/j.jacc.2019.10.040 31865973;74(24):3050–65.
52. Jahangir A, Lee V, Friedman PA, Trusty JM, Hodge DO, Kopecky SL, Packer DL, Hammill SC, Shen WK, Gersh BJ. Long-term progression and outcomes with aging in patients with lone atrial fibrillation: A 30-year follow-up study. *Circulation.* 2007 10.1161/CIRCULATIONAHA.106.644484 17548732;115(24):3050–6.
53. Swynghedauw B. Molecular mechanisms of myocardial remodeling. *Physiol Rev.* 1999

10.1152/physrev.1999.79.1.215 9922372;79(1):215–62.

54. Hirsh BJ, Copeland-Halperin RS, Halperin JL. Fibrotic atrial cardiomyopathy, atrial fibrillation, and thromboembolism: Mechanistic links and clinical inferences. *J Am Coll Cardiol* [Internet]. 2015 10.1016/j.jacc.2015.03.557;65(20):2239–51. Available from: 10.1016/j.jacc.2015.03.557
55. Boldt A, Wetzel U, Lauschke J, Weigl J, Gummert J, Hindricks G, Kottkamp H, Dhein S. Fibrosis in left atrial tissue of patients with atrial fibrillation with and without underlying mitral valve disease. *Heart*. 2004 10.1136/hrt.2003.015347 15020515;90(4):400–5.
56. Guichard JB, Nattel S. Atrial Cardiomyopathy: A Useful Notion in Cardiac Disease Management or a Passing Fad? *J Am Coll Cardiol* [Internet]. 2017 10.1016/j.jacc.2017.06.033;70(6):756–65. Available from: 10.1016/j.jacc.2017.06.033
57. Yaghi S, Kamel H, Elkind MS V. Atrial cardiopathy: a mechanism of cryptogenic stroke. *Expert Rev Cardiovasc Ther* [Internet]. 2017 Aug 10.1080/14779072.2017.1355238 28718666;15(8):591–9. Available from: 10.1080/14779072.2017.1355238
58. Fonseca AC, Alves P, Inácio N, Marto JP, Viana-Baptista M, Pinho-E-Melo T, Ferro JM, Almeida AG. Patients with undetermined stroke have increased atrial fibrosis: A cardiac magnetic resonance imaging study. *Stroke*. 2018 10.1161/STROKEAHA.117.019641 29371431;49(3):734–7.
59. Platonov PG, Mitrofanova LB, Orshanskaya V, Ho SY. Structural abnormalities in atrial walls are associated with presence and persistency of atrial fibrillation but not with age. *J Am Coll Cardiol*. 2011 10.1016/j.jacc.2011.05.061;58(21):2225–32.
60. Oakes RS, Badger TJ, Kholmovski EG, Segerson NM, Burgon NS, Fish EN, Blauer JJ,

- Rao SN, DiBella E V, Akoum N, et al. Detection and Quantification of Left Atrial Structural Remodeling Using Delayed Enhancement MRI in Patients with Atrial Fibrillation. *Circulation*. 2009 10.1161/CIRCULATIONAHA.108.811877.Detection;119(13):1758–67.
61. Benito EM, Carlosena-Remirez A, Guasch E, Prat-González S, Perea RJ, Figueras R, Borràs R, Andreu D, Arbelo E, Tolosana JM, et al. Left atrial fibrosis quantification by late gadolinium-enhanced magnetic resonance: A new method to standardize the thresholds for reproducibility. *Europace*. 2017 10.1093/europace/euw219;19(8):1272–9.
 62. Nazir SA, Lab MJ. Mechanoelectric feedback in the atrium of the isolated guinea-pig heart. *Cardiovasc Res*. 1996 10.1016/0008-6363(96)00077-6 8776408;32(1):112–9.
 63. Bode F, Katchman A, Woosley RL, Franz MR. Gadolinium decreases stretch-induced vulnerability to atrial fibrillation. *Circulation*. 2000 10.1161/01.CIR.101.18.2200 10801762;101(18):2200–5.
 64. Benjamin EJ, D’Agostino RB, Belanger AJ, Wolf PA, Levy D. Left Atrial Size and the Risk of Stroke and Death. *Circulation* [Internet]. 1995 Aug 15 10.1161/01.CIR.92.4.835;92(4):835–41. Available from: 10.1161/01.CIR.92.4.835
 65. Tullio MR Di, Sacco RL, Sciacca RR, Homma S. Ethnically Mixed Population. *Stroke*. 1999;3–8.
 66. Yaghi S, Moon YP, Mora-Mclaughlin C, Willey JZ, Cheung K, Di Tullio MR, Homma S, Kamel H, Sacco RL, Elkind MSV. Left Atrial Enlargement and Stroke Recurrence: The Northern Manhattan Stroke Study. *Stroke*. 2015 10.1161/STROKEAHA.115.008711;46(6):1488–93.
 67. Lang RM, Badano LP, Victor MA, Afzal J, Armstrong A, Ernande L, Flachskampf

- FA, Foster E, Goldstein SA, Kuznetsova T, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: An update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* [Internet]. 2015 10.1016/j.echo.2014.10.003 25559473;28(1):1-39.e14. Available from: 10.1016/j.echo.2014.10.003
68. Rodevan O, Bjornerheim R, Ljosland M, Maehle J, Smith HJ, Ihlen H. Left atrial volumes assessed by three- and two-dimensional echocardiography compared to MRI estimates. *Int J Card Imaging* [Internet]. 1999 Oct 10.1023/a:1006276513186 10595406;15(5):397–410. Available from: 10.1023/a:1006276513186
 69. Shimada YJ, Shiota T. Underestimation of left atrial volume by three-dimensional echocardiography validated by magnetic resonance imaging: A meta-analysis and investigation of the source of bias. *Echocardiography*. 2012 10.1111/j.1540-8175.2011.01593.x;29(4):385–90.
 70. Krohn Therkelsen S, Aaris Groenning B, Hastrup Svendsen J, Boje Jensen G. Atrial and Ventricular Volume and Function Evaluated by Magnetic Resonance Imaging in Patients With Persistent Atrial Fibrillation Before and After Cardioversion. *Am J Cardiol*. 2006 10.1016/j.amjcard.2005.11.040;97(8):1213–9.
 71. Anselmino M, Blandino A, Beninati S, Rovera C, Boffano C, Belletti M, Caponi D, Scaglione M, Cesarani F, Gaita F. Morphologic analysis of left atrial anatomy by magnetic resonance angiography in patients with atrial fibrillation: A large single center experience. *J Cardiovasc Electrophysiol*. 2011 10.1111/j.1540-8167.2010.01853.x;22(1):1–7.
 72. Cozma D, Popescu BA, Lighezan D, Lucian P, Mornos C, Ginghina C, Dragulescu SI. Left atrial remodeling: Assessment of size and shape to detect vulnerability to atrial

- fibrillation. *PACE - Pacing Clin Electrophysiol.* 2007 10.1111/j.1540-8159.2007.00626.x;30(SUPPL. 1):147–50.
73. Kurotobi T, Iwakura K, Inoue K, Kimura R, Toyoshima Y, Ito N, Mizuno H, Shimada Y, Fujii K, Nanto S, et al. The significance of the shape of the left atrial roof as a novel index for determining the electrophysiological and structural characteristics in patients with atrial fibrillation. *Europace.* 2011 10.1093/europace/eur039;13(6):803–8.
 74. Bisbal F, Guiu E, Calvo N, Marin D, Berruezo A, Arbelo E, Ortiz-Pérez J, De Caralt TM, Tolosana JM, Borràs R, et al. Left atrial sphericity: A new method to assess atrial remodeling. impact on the outcome of atrial fibrillation ablation. *J Cardiovasc Electrophysiol.* 2013 10.1111/jce.12116;24(7):752–9.
 75. Scherr D, Dalal D, Chilukuri K, Dong J, Spragg D, Henrikson CA, Nazarian S, Cheng A, Berger RD, Abraham TP, et al. Incidence and predictors of left atrial thrombus prior to catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol.* 2009 10.1111/j.1540-8167.2008.01336.x;20(4):379–84.
 76. Savelieva I, Bajpai A, Camm AJ. Stroke in atrial fibrillation: Update on pathophysiology, new antithrombotic therapies, and evolution of procedures and devices. *Ann Med.* 2007 10.1080/07853890701320662 17701479;39(5):371–91.
 77. Puwanant S, Varr BC, Shrestha K, Hussain SK, Tang WHW, Gabriel RS, Wazni OM, Bhargava M, Saliba WI, Thomas JD, et al. Role of the CHADS2 Score in the Evaluation of Thromboembolic Risk in Patients With Atrial Fibrillation Undergoing Transesophageal Echocardiography Before Pulmonary Vein Isolation. *J Am Coll Cardiol [Internet].* 2009 10.1016/j.jacc.2009.07.037;54(22):2032–9. Available from: 10.1016/j.jacc.2009.07.037
 78. Belgaid DR, Khan Z, Zaidi M, Hobbs A. Prospective randomized evaluation of the

- watchman left atrial appendage closure device in patients with atrial fibrillation versus long-term warfarin therapy the PREVAIL trial. *Int J Cardiol* [Internet]. 2016 10.1016/j.ijcard.2016.06.041;219:177–9. Available from: 10.1016/j.ijcard.2016.06.041
79. Holmes DR, Kar S, Price MJ, Whisenant B, Sievert H, Doshi SK, Huber K, Reddy VY. Prospective randomized evaluation of the watchman left atrial appendage closure device in patients with atrial fibrillation versus long-term warfarin therapy: The PREVAIL trial. *J Am Coll Cardiol* [Internet]. 2014 10.1016/j.jacc.2014.04.029;64(1):1–12. Available from: 10.1016/j.jacc.2014.04.029
80. Khurram IM, Dewire J, Mager M, Maqbool F, Zimmerman SL, Zipunnikov V, Beinart R, E. Marine J, Spragg DD, Berger RD, et al. Relationship between left atrial appendage morphology and stroke in patients with atrial fibrillation. *Hear Rhythm* [Internet]. 2013 10.1016/j.hrthm.2013.09.065;10(12):1843–9. Available from: 10.1016/j.hrthm.2013.09.065
81. Di Biase L, Santangeli P, Anselmino M, Mohanty P, Salvetti I, Gili S, Horton R, Sanchez JE, Bai R, Mohanty S, et al. Does the left atrial appendage morphology correlate with the risk of stroke in patients with atrial fibrillation? Results from a multicenter study. *J Am Coll Cardiol* [Internet]. 2012 10.1016/j.jacc.2012.04.032;60(6):531–8. Available from: 10.1016/j.jacc.2012.04.032
82. Lee JM, Seo J, Uhm JS, Kim YJ, Lee HJ, Kim JY, Sung JH, Pak HN, Lee MH, Joung B. Why is left atrial appendage morphology related to strokes? an analysis of the flow velocity and orifice size of the left atrial appendage. *J Cardiovasc Electrophysiol*. 2015 10.1111/jce.12710;26(9):922–7.
83. Goldman ME, Pearce LA, Hart RG, Zabalgoitia M, Asinger RW, Safford R, Halperin JL. Pathophysiologic correlates of thromboembolism in nonvalvular atrial fibrillation:

- I. Reduced flow velocity in the left atrial appendage (The stroke prevention in atrial fibrillation [SPAF-III] study). *J Am Soc Echocardiogr.* 1999 10.1016/S0894-7317(99)70105-7 10588784;12(12):1080–7.
84. Lee JM, Shim J, Uhm JS, Kim YJ, Lee HJ, Pak HN, Lee MH, Joung B. Impact of increased orifice size and decreased flow velocity of left atrial appendage on stroke in nonvalvular atrial fibrillation. *Am J Cardiol* [Internet]. 2014 10.1016/j.amjcard.2013.11.058;113(6):963–9. Available from: 10.1016/j.amjcard.2013.11.058
85. Goda T, Sugiyama Y, Ohara N, Ikegami T, Watanabe K, Kobayashi J, Takahashi D. P-Wave Terminal Force in Lead V1 Predicts Paroxysmal Atrial Fibrillation in Acute Ischemic Stroke. *J Stroke Cerebrovasc Dis* [Internet]. 2017 10.1016/j.jstrokecerebrovasdis.2017.06.031;26(9):1912–5. Available from: 10.1016/j.jstrokecerebrovasdis.2017.06.031
86. Morris JJ, ESTES EH, WHALEN RE, THOMPSON HK, MCINTOSH HD. P-Wave Analysis in Valvular Heart Disease. *Circulation* [Internet]. 1964 Feb 10.1161/01.CIR.29.2.242 14119389;29(2):242–52. Available from: 10.1161/01.CIR.29.2.242
87. Hancock EW, Deal BJ, Mirvis DM, Okin P, Kligfield P, Gettes LS. AHA/ACCF/HRS Recommendations for the Standardization and Interpretation of the Electrocardiogram. Part V: Electrocardiogram Changes Associated With Cardiac Chamber Hypertrophy A Scientific Statement From the American Heart Association Electrocardiography . *J Am Coll Cardiol* [Internet]. 2009 10.1016/j.jacc.2008.12.015;53(11):992–1002. Available from: 10.1016/j.jacc.2008.12.015
88. Win TT, Venkatesh BA, Volpe GJ, Mewton N, Rizzi P, Sharma RK, Strauss DG, Lima

- JA, Tereshchenko LG. Associations of electrocardiographic P-wave characteristics with left atrial function, and diffuse left ventricular fibrosis defined by cardiac magnetic resonance: The PRIMERI study. *Hear Rhythm*. 2015 10.1016/j.hrthm.2014.09.044;12(1):155–62.
89. Kohsaka S, Sciacca RR, Sugioka K, Sacco RL, Homma S, Di Tullio MR. Electrocardiographic left atrial abnormalities and risk of ischemic stroke. *Stroke*. 2005 10.1161/01.STR.0000185682.09981.26 16210557;36(11):2481–3.
 90. Soliman EZ, Prineas RJ, Case LD, Zhang Z, Goff DC. Ethnic Distribution of ECG Predictors of Atrial Fibrillation and Its Impact on Understanding the Ethnic Distribution of Ischemic Stroke in the Atherosclerosis Risk in Communities (ARIC) Study. *Stroke* [Internet]. 2009 Apr 10.1161/STROKEAHA.108.534735 1000000221;40(4):1204–11. Available from: 10.1161/STROKEAHA.108.534735
 91. Kamel H, O’Neal WT, Okin PM, Loehr LR, Alonso A, Soliman EZ. Electrocardiographic left atrial abnormality and stroke subtype in the atherosclerosis risk in communities study. *Ann Neurol* [Internet]. 2015 Nov 10.1002/ana.24482;78(5):670–8. Available from: 10.1002/ana.24482
 92. Kamel H, Hunter M, Moon YP, Yaghi S, Cheung K, Di Tullio MR, Okin PM, Sacco RL, Soliman EZ, Elkind MSV. Electrocardiographic left atrial abnormality and risk of stroke: Northern manhattan study. *Stroke*. 2015 10.1161/STROKEAHA.115.009989 26396031;46(11):3208–12.
 93. Thomas A. Dewland, MD; Eric Vittinghoff, PhD, MPH; Mala C. Mandyam, MD; Susan R. Heckbert, MD, PhD; David S. Siscovick, MD M, Phyllis K. Stein, PhD; Bruce M. Psaty, MD, PhD; Nona Sotoodehnia, MD; John S. Gottdiener, MD; and Gregory M. Marcus, MD M. Original Research Atrial Ectopy as a Predictor of Incident Atrial

- Fibrillation. *Ann Intern Med*. 2013;159:721–8.
94. Ninan Koshy A, Sajeev J, Rajakariar K, Briganti E, Andrew P, Zureik M, Quine E, Parfrey S, Venkataraman P, Roberts L, et al. High Atrial Ectopy in Stroke: an Independent Risk Factor? *J Am Coll Cardiol* [Internet]. 2017 10.1016/s0735-1097(17)33727-0;69(11):338. Available from: 10.1016/s0735-1097(17)33727-0
 95. Larsen BS, Kumarathurai P, Falkenberg J, Nielsen OW, Sajadieh A. Excessive Atrial Ectopy and Short Atrial Runs Increase the Risk of Stroke beyond Incident Atrial Fibrillation. *J Am Coll Cardiol*. 2015 10.1016/j.jacc.2015.05.018;66(3):232–41.
 96. Dilaveris PE, Färbon P, Batchvarov V, Ghuran A, Malik M. Circadian behavior of P-wave duration, P-wave area, and PR interval in healthy subjects. *Ann Noninvasive Electrocardiol*. 2001 10.1111/j.1542-474X.2001.tb00092.x;6(2):92–7.
 97. Cheng S. Long-term Outcomes in Individuals With Prolonged PR Interval or First-Degree Atrioventricular Block. *JAMA* [Internet]. 2009 Jun 24 10.1001/jama.2009.88819549974;301(24):2571. Available from: 10.1001/jama.2009.888
 98. Thijs VN, Brachmann J, Morillo CA, Passman RS, Sanna T, Bernstein RA, Diener H-C, Di Lazzaro V, Rymer MM, Hogge L, et al. Predictors for atrial fibrillation detection after cryptogenic stroke. *Neurology* [Internet]. 2016 Jan 19 10.1212/WNL.0000000000002282 26683642;86(3):261–9. Available from: 10.1212/WNL.0000000000002282
 99. Lev M. Anatomic basis for atrioventricular block. *Am J Med*. 1964 10.1016/0002-9343(64)90022-1;37(5):742–8.
 100. Fleg JL, Das DN, Wright J, Lakatta EG. Age-Associated Changes in the Components of Atrioventricular Conduction in Apparently Healthy Volunteers. *J Gerontol* [Internet]. 1990 May 1 10.1093/geronj/45.3.M95 2335725;45(3):M95–100. Available from:

10.1093/geronj/45.3.M95

101. Agarwal YK, Aronow WS, Levy JA, Spodick DH. Association of interatrial block with development of atrial fibrillation. *Am J Cardiol*. 2003 10.1016/S0002-9149(03)00027-4;91(7):882.
102. Montalvo M, Tadi P, Merkler A, Gialdini G, Martin-Schild S, Navalkele D, Samai A, Nouh A, Hussain M, Goldblatt S, et al. PR Interval Prolongation and Cryptogenic Stroke: A Multicenter Retrospective Study. *J Stroke Cerebrovasc Dis* [Internet]. 2017 10.1016/j.jstrokecerebrovasdis.2017.05.036;26(10):2416–20. Available from: 10.1016/j.jstrokecerebrovasdis.2017.05.036
103. Schnabel RB, Sullivan LM, Levy D, Pencina MJ, Massaro JM, D'Agostino RB, Newton-Cheh C, Yamamoto JF, Magnani JW, Tadros TM, et al. Development of a risk score for atrial fibrillation (Framingham Heart Study): a community-based cohort study. *Lancet* [Internet]. 2009 10.1016/S0140-6736(09)60443-8;373(9665):739–45. Available from: 10.1016/S0140-6736(09)60443-8
104. Hijazi Z, Oldgren J, Siegbahn A, Granger CB, Wallentin L. Biomarkers in atrial fibrillation : a clinical review *Cardiac biomarkers*. 2013 10.1093/eurheartj/ehf024;1475–80.
105. Shelton RJ, Clark AL, Goode K, Rigby AS, Cleland JGF. The diagnostic utility of N-terminal pro-B-type natriuretic peptide for the detection of major structural heart disease in patients with atrial fibrillation. 2006 10.1093/eurheartj/ehl233;2353–61.
106. Silvet H, Young-xu Y, Walleigh D, Ravid S. Brain Natriuretic Peptide Is Elevated in Outpatients. 2003 10.1016/S0002-9149(03)01053-1;92(03):2–5.
107. Hijazi Z, Wallentin L, Siegbahn A, Andersson U, Ms C, Christersson C, Ezekowitz J, Mbbc H, Ms C, Gersh BJ, et al. N-Terminal Pro – B-Type Natriuretic Peptide for Risk

- Assessment in Patients With Atrial Fibrillation Insights From the ARISTOTLE Trial (Apixaban for the Prevention of Stroke in Subjects With Atrial Fibrillation). 2013 10.1016/j.jacc.2012.11.082;61(22).
108. Hijazi Z, Oldgren J, Andersson U, Connolly SJ, Ezekowitz MD, Hohnloser SH, Reilly PA. Cardiac Biomarkers Are Associated With an Increased Risk of Stroke and Death in Patients With Atrial Fibrillation A Randomized Evaluation of Long-Term Anticoagulation Therapy. 2012 10.1161/CIRCULATIONAHA.111.038729;
 109. Shibazaki K, Kimura K, Fujii S, Sakai K. Brain Natriuretic Peptide Levels as a Predictor for New Atrial Fibrillation During Hospitalization in Patients With Acute Ischemic Stroke. *AJC* [Internet]. 2012 10.1016/j.amjcard.2011.12.022;109(9):1303–7. Available from: 10.1016/j.amjcard.2011.12.022
 110. Suissa L, Bresch S, Lachaud S, Marie H. Brain Natriuretic Peptide : A Relevant Marker to Rule Out Delayed Atrial Fibrillation in Stroke Patient. 2013 10.1016/j.jstrokecerebrovasdis.2012.08.010;22(7):103–10.
 111. Li-Saw-Hee FL, Blann AD, Goldsmith I, Lip GYH. Indexes of hypercoagulability measured in peripheral blood reflect levels in intracardiac blood in patients with atrial fibrillation secondary to mitral stenosis. *Am J Cardiol*. 1999 10.1016/S0002-9149(99)00060-0;83(8):1206–9.
 112. Akar JG, Jeske W, Wilber DJ. Acute Onset Human Atrial Fibrillation Is Associated With Local Cardiac Platelet Activation and Endothelial Dysfunction. *J Am Coll Cardiol*. 2008 10.1016/j.jacc.2007.11.083;51(18):1790–3.
 113. Choudhury A, Chung I, Blann AD, Lip GYH. Elevated platelet microparticle levels in nonvalvular atrial fibrillation: Relationship to P-selectin and antithrombotic therapy. *Chest* [Internet]. 2007 10.1378/chest.06-2039;131(3):809–15. Available from:

10.1378/chest.06-2039

114. Tan KT, Lip GYH. Atrial fibrillation: should we target platelets or the coagulation pathway? Card Electrophysiol Rev. 2003 10.1023/B:CEPR.0000023141.17553.f1;7(4):370–1.
115. Goette A, Kalman JM, Aguinaga L, Akar J, Cabrera JA, Chen SA, Chugh SS, Corradi D, D'Avila A, Dobrev D, et al. EHRA/HRS/APHRS/SOLAECE expert consensus on atrial cardiomyopathies: Definition, characterization, and clinical implication. Europace. 2016 10.1093/europace/euw161 27402624;18(10):1455–90.




Figure 1:

Schematic representation of the relationship between atrial cardiomyopathy, atrial fibrillation and thromboembolic disease⁵⁴




Figure 2: Graphical representation showing temporal dissociation between Atrial Tachycardia/Atrial Fibrillation (AT/AF) and Thromboembolic stroke (TE) in ASSERT, TRENDS and IMPACT studies. Reproduced with permission from Hirsch et al. (2015).⁵⁴




Figure 3:

Representative left atrial cardiac magnetic resonance imaging (CMR) showing various degrees of late gadolinium enhancement suggestive of a normal left atrium (A), focal area of fibrosis on the posterior wall of the left atrium (B), diffuse posterior wall fibrosis (C) and circumferential fibrosis (D).

Study Name	Study Design	Sample Size	Patient cohort	Treatment group	Primary Outcome Measure	Findings
WARSS (1993-2000) ¹⁷	Multi-center, double-blind, randomised control study	2206	Cryptogenic stroke	Warfarin	Recurrent stroke or death	No difference between aspirin and warfarin in prevention of recurrent stroke/death or rate of major haemorrhage
NAVIGATE ESUS (2014-2018) ²²	Multi-center, double-blind, randomised control study	7213	ESUS	Rivaroxaban	Recurrent ischaemic /haemorrhagic stroke or systemic embolism	Rivaroxaban was not superior to aspirin in prevention of recurrent stroke and was associated with a higher risk of bleeding
RE-SPECT ESUS (2014-2018) ²³	Multi-center, double-blind, randomised control study	5390	ESUS	Dabigatran	Recurrent stroke	Dabigatran was not superior to aspirin, but had a higher risk of non-major haemorrhage
ATTICUS (2015-2021) ²⁵	Multi-center, open label, randomised control trial	Approx. 500	ESUS and non-major risk factors for cardio-embolic stroke	Apixaban	At least one new ischaemic lesion identified on MRI imaging at 12 months	Study in progress
ARCADIA (2018-2022) ¹²	Multi-center, double-blind, randomised control study	Approx. 1100	ESUS and biomarkers of atrial cardiomyopathy	Apixaban	Recurrent ischaemic/haemorrhagic stroke	Study in progress

Table 1: Studies assessing aspirin versus anticoagulation therapy in reducing stroke recurrence in ischaemic stroke patients.

Table 2: Table showing features of atrial cardiomyopathy and evidence of association with AF and stroke.

	Study/Author	Sample size	Association with AF	Association with Stroke	Limitations
Atrial Fibrosis	Platonov et al. ⁵⁹	30 patients	Patients with PsAF had greater fibrosis than those with PAF. 2-3x increase in fibrosis and fatty infiltration in patients with AF compared with sinus rhythm.	N/A	CMR imaging is expensive Accuracy of findings, in particular atrial fibrosis, is dependent on image quality acquired Large interobserver reproducibility of left atrial appendage morphology Large variability in image acquisition parameters, image analysis and post-processing tools across different centres
	Boldt et al. ⁵⁵	118 patients	Increased protein concentrations in patients with AF compared with those in sinus rhythm. No systematic difference in ECM expression between PAF and PsAF.	N/A	
	Fonseca et al. ⁵⁸	111 patients	N/A	Cryptogenic stroke patients had a higher percentage of LA fibrosis compared with other stroke types ($p=0.03$). Cryptogenic stroke patients had an Intermediate LA phenotype compared with those with AF and other stroke causes.	
Atrial enlargement and Geometry	Nazir et al. ⁶²	12 guinea-pigs (ex vivo)	Increased LA volume and load changes resulted increased arrhythmia occurrence or premature atrial beats.	N/A	
	Bode et al. ⁶³	16 rabbits (ex vivo)	Acute atrial stretch significantly enhances vulnerability to AF	N/A	

	<i>Framingham Study Benjamin et al.⁶⁴</i>	<i>3099 patients (sub-group analysis)</i>	<i>N/A</i>	<i>After multivariable adjustment, for every 10mm increase in LA size, the RR of stroke was 2.4 in men (95% CI, 1.0-1.5) and 1.4 in women (95% CI, 1.1-1.7).</i>
	<i>Tullio et al.⁶⁵</i>	<i>721 patients (352 study group, 369 control group)</i>	<i>N/A</i>	<i>LA index was associated with ischaemic stroke in the overall group (adjusted OR 1.47 per 10mm/1.7m² of BSA; 95% CI 1.03-2.11).</i>
	<i>Cozma et al.⁷²</i>	<i>112 patients</i>	<i>LA dilatation is accompanied by changes in LA shape. Proportion of patients with trapezoidal LA shape was increased. LA dilatation and electrical remodelling are interrelated.</i>	<i>N/A</i>
	<i>Bisbal et al.⁷⁴</i>	<i>106 patients</i>	<i>Left atrial sphericity is an independent predictor of recurrence of AF after AF ablation.</i>	<i>N/A</i>
<i>Left atrial appendage (LAA)</i>	<i>Belgaid et al.⁷⁸</i>	<i>407 patients</i>	<i>N/A</i>	<i>LAA occlusion is non-inferior to warfarin in secondary stroke prevention in non-valvular AF.</i>
	<i>Holmes et al.⁷⁹</i>	<i>407 patients</i>	<i>N/A</i>	<i>LAA occlusion is non-inferior to warfarin in secondary stroke prevention in non-valvular AF.</i>
	<i>Khurram et al.⁸⁰</i>	<i>65 patients</i>	<i>N/A</i>	<i>Extent of LAA trabeculations and smaller LAA orifice diameter are associated with stroke prevalence in AF patients.</i>

				LAA morphologies were not associated with stroke risk.	
	Di Biase et al. ⁸¹	932 patients	N/A	Chicken Wing morphology is 79% less likely to have a stroke/TIA history (OR: 0.21, CI 95% 0.05-0.91, p=0.036) in AF patients. Cauliflower morphology is 8 times (p=0.056) more likely to have had a stroke/TIA.	
	Lee et al. ⁸²	218 patients (study group =67, control group = 151)	N/A	In patients with flow velocity <37.0cm/s, larger LAA orifice (>3.5cm ²) was associated with a greater incidence of stroke in AF patients	
	SPAF-III Study Goldman et al. ⁸³ .	721 patients	N/A	Patients in AF displayed lower peak antegrade (emptying) flow velocity than those in SR or with intermittent AF. AVp<20cm/s was associated with appendage thrombus (p<0.01) and cardioembolic events (P<0.01).	
P-wave terminal force in lead V ₁	Goda et al. ⁸⁵	295 patients	PTFV ₁ was significantly higher in patients with PAF than those without PAF (-.051 vs. 0.027 mm.s, p<0.001).	N/A	Despite strong association, predictive models have moderate or low predictive capacity.
	Northern Manhattan Study Kamel et al. ⁹²	1039 patients (study group =241, control group = 798)	N/A	PTFV ₁ was associated with cryptogenic or cardioembolic stroke (HR: 1.14 95%CI 1.08-1.58).	

	<i>Atherosclerosis risk in Communities Study Kamel et al.⁹¹</i>	14,552 patients	N/A	<i>PTFV₁ >40ms.mm was associated with incident ischaemic stroke (HR 1.33, 95% CI: 1.11-1.59) in the absence of AF.</i>	
	<i>ARIC Study Soliman et al.⁹⁰</i>	15,429 patients	<i>HR for incident AF associated with PTFV₁ was 1.22 (CI: 1.14-1.31) when corrected for demographic and clinical variables.</i>	<i>PTFV₁ was the strongest AF predictor for incident stroke. HR: 2.28 (95% CI 1.79-2.90).</i>	
	<i>Kohsaka et al.⁸⁹</i>	341 patients (study group =146, control group = 195 control)	N/A	<i>PTFV₁ >40ms.mm was associated with ischaemic stroke after adjustment for other stroke risk factors (OR: 2.32, 95%CI 1.29-1.48). Effect of PTFV₁ on stroke risk was strongly dependent on LV mass.</i>	
<i>Atrial ectopy</i>	<i>Koshy et al.⁹⁴</i>	2800 patients	N/A	<i>Mean premature atrial complex count on 24-hour monitoring 426 vs 105 (study group vs. control group) with P-value <0.001.</i>	
	<i>Copenhagen Holter Study Cohort Larsen et al.⁹⁵</i>	678 patients	N/A	<i>Excessive atrial ectopic activity (>30 ectopics/hr or runs of over 20 ectopics) associated with an increase risk of stroke in absence of AF.</i>	
	<i>Dewland et al.⁹³</i>	5201 patients	<i>Atrial ectopy showed similar AF risk discrimination compared with AF Framingham Risk score. A 100% increase in hourly atrial ectopy was associated with a significant increase in AF risk (HR: 1.17, 95% CI 1.13 – 1.22, P<0.001).</i>		
<i>Prolongation of PR interval</i>	<i>Framingham Heart Study Cheng et al⁹⁷</i>	7575 patients	<i>PR >200ms associated with a two-fold adjusted risk of AF (HR 2.06, 95% CI: 1.36-3.12, p<0.001).</i>	N/A	

	<i>CRYSTAL AF Thijis et al.⁸</i>	<i>221 patients</i>	<i>PR prolongation was independently associated with an increased AF incidence in cryptogenic stroke patients (HR 1.3 per 10ms, 95% CI 1.2-1.4, P<0.0001).</i>	<i>N/A</i>	
	<i>Montalvo et al.¹⁰²</i>	<i>687 patients</i>		<i>Mean PR interval was longer in cryptogenic stroke patients compared with other known stroke aetiology (175.4 vs. 166.5). 23.2% of patients with cryptogenic stroke had a PR interval >200ms compared with 13.8% of those with alternative stroke aetiology (P = 0.009).</i>	

(AF = atrial fibrillation, PAF = paroxysmal atrial fibrillation, PsAF = persistent atrial fibrillation, ECM = extracellular matrix, LA = left atrium, LAA = left atrial appendage, RR = relative risk, OR = odds ratio, HR = hazard ratio, CI = confidence interval)