

Electrophysiological responses of *Cydia pomonella* to codlemone and pear ester ethyl (*E,Z*)-2,4-decadienoate: peripheral interactions in their perception and evidences for cells responding to both compounds

Antonio DE CRISTOFARO¹, Claudio IORIATTI², Edison PASQUALINI³, Gianfranco ANFORA^{1,2}, Giacinto Salvatore GERMINARA¹, Michela VILLA^{2,4}, Giuseppe ROTUNDO¹

¹Dipartimento di Scienze Animali, Vegetali e dell'Ambiente, Università degli Studi del Molise, Campobasso, Italy

²Istituto Agrario di San Michele all'Adige, San Michele all'Adige, Trento, Italy

³Dipartimento di Scienze e Tecnologie Agroambientali - Entomologia, Università di Bologna, Italy

⁴Istituto di Entomologia e Patologia Vegetale, Università Cattolica del Sacro Cuore, Piacenza, Italy

Abstract

Electroantennography (EAG) recordings were made from both virgin and mated males and females of *Cydia pomonella* (L.) (Lepidoptera Tortricidae) (CM) on stimulation with the main component of its sex pheromone (*E,E*)-8,10-dodecadien-1-ol (*E8E10-12:OH*) and ethyl (*E,Z*)-2,4-decadienoate (Et-2*E,4Z*-DD), a ripe pear-derived volatile attractant. CMs of 7 Italian populations collected on different host-plants (3 apple, 3 walnut, 1 pear) were used. The dose-response curves to the volatile compounds (9 doses) were calculated and differences in olfactory sensitivity of virgin and mated males and females of the different populations are discussed. As expected, females were much less sensitive than males to *E8E10-12:OH* but generally showed a similar response to the higher doses of Et-2*E,4Z*-DD. EAG amplitudes were not significantly different among virgin and mated adults, as well as between two populations collected in a chemical treated apple orchard and an organic farm, respectively. The males of the walnut populations showed a lightly higher sensitivity to *E8E10-12:OH*.

Differential saturation experiments (DS-EAG) showed that the antennae of *C. pomonella* clearly reduce their response to Et-2*E,4Z*-DD when continuously stimulated with *E8E10-12:OH*. Recordings from single antennal olfactory cells SCR showed the presence of cells responding only to *E8E10-12:OH* or Et-2*E,4Z*-DD but also to the 2 compounds.

During field tests, all doses of Et-2*E,4Z*-DD showed an attractant activity on both sexes of CM. A synergistic effect on male attraction was not observed by adding Et-2*E,4Z*-DD to the pheromone; nevertheless, a clear interference on pheromone trap efficiency was detected.

These observations have to be highly considered when setting up mixtures of *E8E10-12:OH* and Et-2*E,4Z*-DD to monitor codling moth populations.

Key words: Sex pheromones, kairomones, EAG, DS-EAG, SCR, field test, codling moth.

Introduction

The codling moth (CM), *Cydia pomonella* (L.) (Lepidoptera Tortricidae), is the key insect pest of the apple orchard. Insecticides-based control strategies are still widely used. The CM insecticide resistance induced to change the traditional approach and, as a consequence, virus (CMGV) and semiochemical-based control methods (mating disruption) were introduced (Ioriatti and Bouvier, 2000).

New control strategies need of a more efficient monitoring lure able to provide information on both males and females field behaviour. Ethyl (*E,Z*)-2,4-decadienoate (Et-2*E,4Z*-DD), a ripe pear derived kairomone, is reported as highly attractive to the CM larvae (Knight and Light, 2001) and both male and female (virgin and mated) adults in walnut and apple orchards (Light *et al.*, 2001; Coracini *et al.*, 2004).

Host-plant volatiles can synergize the synthetic pheromone attractant power in CM and other insect species (Dickens *et al.*, 1990; Light *et al.*, 1993; Yang *et al.*, 2004). In order to improve the efficiency of the traps and the Attract and Kill control method, the hypothesized synergistic effect of Et-2*E,4Z*-DD on (*E,E*)-8,10-

dodecadien-1-ol (*E8E10-12:OH*, codlemone) attractancy needs to be investigated. It is also important to underline that traps baited with Et-2*E,4Z*-DD showed a different attractiveness according to the CM host-plant (Light *et al.*, 2001).

In addition, insecticide resistance may limit the effectiveness and attractiveness of mixed formulations where resistance pleiotropic effects influence pheromone perception. It has already been shown that the dose-response relationships of males of susceptible and resistant strains of the CM to *E8E10-12:OH* does not differ significantly (Beslay *et al.*, 2000). This occurrence could be different for the response to Et-2*E,4Z*-DD, as far as the enzymatic system mainly involved in insecticide detoxification in CM, like monooxygenases (Sauthanor *et al.*, 1997), are also involved in the recognition of the host-plant (Feyresen, 1999).

In the present paper electrophysiological responses (electroantennography, EAG; differential saturation-electroantennography, DS-EAG; single cell recording, SCR) to *E8E10-12:OH* and Et-2*E,4Z*-DD of 7 CM populations living on different host plant (apple, pear, walnut) and under different cropping conditions (organic and chemical-based control methods) were re-

corded, focusing the attention on the possible influence of the kairomone on the perception of the sex pheromone main component. Moreover, field tests were carried out using both compounds and some blends.

Materials and methods

Insects

Overwintering larvae of CM were collected in October 2001 using cardboard bands in 3 walnut (Bleggio, TN; Rovigo, RO; Ceresi, MO), 3 apple (San Michele all'Adige, Gardolo, Borgo Valsugana, TN) and one pear (Maieutica, BO) orchards. S. Michele all'Adige population showed a reduced susceptibility to the insecticides (Ioriatti *et al.*, 2003a), while the Borgo Valsugana one originated from an organic orchard. Larvae were kept in an outdoor insectary till March 2002, than the obtained pupae were isolated in plastic Petri dishes (i.d. 5 cm) and transferred to the laboratory at 16:8 L:D cycle, $23\pm 2^\circ\text{C}$ and $70\pm 5\%$ RH. Moths emerged from the middle of April to the beginning of June. Males and females were kept in different rearing rooms. Adults were fed with a 10% (W/V) sucrose aqueous solution absorbed on a cotton pad. During the experiments, 3-6 day old moths (virgin and mated) were used.

Stimuli

Stimuli were the main component of the CM sex pheromone (*E8E10-12:OH*; >99% pure) and the pear ester kairomone (*Et-2E,4Z-DD*; >97% pure) (figure 1). Gas chromatography (GC) and GC linked to mass spectrometry (GC-MS) analyses of hexane solution (100 ng/ μl) of synthetic compounds did not detect the presence of related isomers or traces of *E8E10-12:OH* in *Et-2E,4Z-DD*. GC coupled to electroantennography detector (GC-EAD) analyses did not reveal any EAG activity at the *E8E10-12:OH* retention time. Attractants were supplied by Sigma-Aldrich (Germany). Aliquots of 10 μl of a mineral oil solution of a chemical were absorbed on a piece of filter paper (1 cm^2). For each compound, nine different stimuli containing from 10^{-6} to 10^2 μg were prepared. During EAG and DS-EAG tests all stimuli were applied; in SCRs, after preliminary assays, the more useful dose (1.0 μg) in discerning the responding cells to one or both compounds was adopted.




Figure 1. Compounds used during the electrophysiological studies: the main component of *C. pomonella* sex pheromone (*E8E10-12:OH*) and the ripe pear ester acting as kairomone (*Et-2E,4Z-DD*).

Electrophysiology

EAG responses (Rotundo and Tremblay, 1993; Den Otter *et al.*, 1996; De Cristofaro *et al.*, 2000) to *E8E10-12:OH* and *Et-2E,4Z-DD* of the 7 populations were recorded. The differential saturation-EAG (DS-EAG) (Baker and Roelofs, 1976; Miller *et al.*, 1977; Nishino and Manabe, 1984) and single cell recordings (SCR, surface contact technique) (Den Otter *et al.*, 1980; 1996) were performed on one population (Ceresi, MO).

The proximal end of an amputated antenna was inserted into a glass pipette indifferent electrode (i.d. tip 1 mm) filled with Beadle-Ephrussi saline containing 5 g/l polyvinylpyrrolidone K90 (Fluka Chemie, Switzerland). For EAG and DS-EAG recordings the different electrode was a similar pipette brought into contact with the distal end of the antenna from which 1-2 segments had been excised. The scale-less area of the antenna lodging, like in other tortricid moth, the highest number of olfactory hairs, was placed perpendicularly to the stimulus direction (De Cristofaro *et al.*, 1997; 2000).

During SCR, spikes from individual cells were recorded by gently pressing the tip (i.d. < 3 μm) of the different glass electrode against the cuticle of the antenna; recordings were made from the rostral-ventral scale-less area of the antenna, at the distal end of a medial segment, where rows of hooked long olfactory sensilla trichoidea and sensilla auriculica are present (figure 2). Silver wires inserted in the pipettes were connected to a portable EAG/SCR equipment (Van der Pers and Minks, 1993). During the EAG and SCR recordings, the antenna was continuously flushed with charcoal-filtered, humidified air at room temperature (23°C). During the DS-EAG measurements the continuous air stream originated from a bottle containing a mineral oil solution (1 $\mu\text{g}/\mu\text{l}$) of a chemical (*E8E10-12:OH* or *Et-2E,4Z-DD*).

The air flowed at 50 cm^3/s through a stainless steel tube (i.d. 8 mm) the outlet of which was about 1 cm from the preparation. Through a hole (i.d. 4 mm), 9 cm from the outlet, stimuli were added to the air stream by injecting vapour from an odour cartridge. During injection, a stimulus controller (CS-01, Syntech, Hilversum, NL) was used to keep constant the flow over the antenna.

In EAG and DS-EAG recordings 2.5 cm^3 were injected during 0.1 s, in SCR experiments 12.5 cm^3 in 500 ms. Intervals between stimuli were 1 min. In EAG and DS-EAG tests, the doses of different chemicals were applied in ascending order; one minute before each series of stimulation with the same dose of chemicals, a reference stimulus (0.01 μg of *E8E10-12:OH*) was applied to correct for changes in EAG responsiveness. In order to verify the effect of saturation on the antennal sensitivity, during DS-EAG tests 2 plant volatile (heptanal, Z3-hexen-1-ol) stimuli (1.0 μg) were applied. EAGs were analysed using a proper programme (EAG 2.0, Syntech, Hilversum, NL). The absolute EAG amplitudes were calculated as described by Den Otter *et al.* (1988; 1991).

Action potentials were stored on tape and analysed using a suitable programme (Autospike 1.1., Syntech, Hilversum, NL); spikes were distinguished on the basis of shape, amplitude and frequency. Only SCRs showing




Figure 2. An antennal segment (33rd) of *C. pomonella* male showing the rostral-ventral scale-less area. The arrows indicate some sensilla trichoidea (long hooked olfactory hairs; i.e. A arrow) and auricillica (rabbit-ear shaped; i.e. B arrow) and where the recording electrode was placed (i.e. C arrow) during SCR recordings.

the activity of one sensitive cell, and during which it was possible to apply the different stimuli at least 3 times (complete recordings), were considered in the data analysis. The response frequency (spikes/s) were referred to the first 100 ms (spikes/100 ms x 10) and subsequently analyzed (LSD test).

Field tests

Trials have been carried out during two years (2003-04) in an organic orchard (Colletorto, CB, South Italy). Pherocon IIB (Trécé) traps were suspended from trees, 2-3 m above ground and spaced 15-20 meters apart.

Based on previous field studies (Ioriatti *et al.*, 2003b), 2 doses of *E8E10-12:OH* (0.1, 1.0 mg), 5 doses of *Et-2E,4Z-DD* (0.1, 1.0; 3.0, 10.0, 40.0 mg), and 2 blends (0.1 + 0.1 mg and 1.0 + 0.1 mg, respectively) were evaluated. Rubber septa dispensers, placed at the centre of the trap bottom, were adopted. Baits were replaced every 3 weeks. Three replicates for each treatment as well as 3 unbaited traps (blank) were used.

Traps were checked weekly and bottoms of those containing insects were replaced; male and female catches were recorded.

Data were analysed by ANOVA and means separated by Duncan test ($P = 0.05$). Prior to the statistical analysis, data were transformed to $\sqrt{x+0.5}$ and submitted to Levene test for homogeneity of variance. For the same bait, differences among male and female catches were evaluated by Student's *t*-test.

Results

Laboratory

For each population, EAG responses to 1.0 μg of *E8E10-12:OH* or *Et-2E,4Z-DD* were not significantly different (*t*-test; $P=0.05$) between virgin and mated males or females (table 1). Comparing the populations on the basis of their host plant (apple, walnut, pear) no statistical differences (Duncan test; $P=0.05$) emerged (table 1).

EAG responses from virgin and mated males and females collected in a chemical treated orchard with a reduced sensitivity to insecticides (San Michele all'Adige, TN) were similar to those of individuals coming from an organic farm (Borgo Valsugana, TN) (table 2).

As expected, females were less sensitive than males to *E8E10-12:OH* (figure 3). This difference was not observed in respect to *Et-2E,4Z-DD*, particularly at higher doses (figure 4).

DS-EAG revealed that the antennae of the CM clearly reduce their response to *E8E10-12:OH* when continuously flushed with *Et-2E,4Z-DD* (figure 3) as well as their response to *Et-2E,4Z-DD* when saturated with *E8E10-12:OH* (figure 4). Responses of virgin adults ($n = 10$) to heptanal (male: 1.43 ± 0.28 mV; female: 1.27 ± 0.32 mV) and *Z3*-hexen-1-ol (male: 1.34 ± 0.36 mV; female: 1.16 ± 0.20 mV) were not significantly reduced by saturation with *E8E10-12:OH* (heptanal: male = 1.26 ± 0.31 mV; female = 1.18 ± 0.26 mV; *Z3*-hexen-1-ol: male = 1.21 ± 0.34 mV; female = 0.96 ± 0.25 mV) or

Table 1. Mean EAG responses of virgin and mated *C. pomonella* adults (n=10/category/population) from 7 Italian populations living on different host plant (3 apple; 3 walnut; 1 pear) to 1.0 µg dose of *E8E10-12:OH* or *Et-2E,4Z-DD*.

		EAG (mV) ± S.D.					
		<i>E8E10-12:OH</i>			<i>Et-2E,4Z-DD</i>		
		Apple ¹	Walnut ¹	Pear ¹	Apple ¹	Walnut ¹	Pear ¹
Females ¹	virgin	0.32±0.09 a	0.38±0.09 a	0.36±0.07 a	1.47±0.23 a	1.56±0.28 a	1.44±0.26 a
	mated	0.36±0.07 a	0.42±0.10 a	0.38±0.08 a	1.56±0.16 a	1.62±0.25 a	1.52±0.28 a
Males ¹	virgin	3.27±0.25 b	3.52±0.34 b	3.32±0.29 b	1.66±0.25 a	1.82±0.28 a	1.62±0.20 a
	mated	3.12±0.29 b	3.32±0.30 b	3.18±0.23 b	1.49±0.28 a	1.69±0.21 a	1.52±0.22 a

Different letters on the same column indicate significant differences between sexes (Duncan test; P=0.05).

¹ Between physiological status (virgin and mated) of both sexes and among populations from different host plant no significant differences were observed (Duncan test; P=0.05).

Table 2. Mean EAG responses of virgin and mated *C. pomonella* adults (n=10/category/population) from a population (I) with a reduced susceptibility to organophosphates (San Michele all'Adige, TN) and a population (II) from an organic orchard (Borgo Valsugana, TN) to 1.0 µg dose of *E8E10-12:OH* or *Et-2E,4Z-DD*.

		EAG (mV) ± S.D.			
		<i>E8E10-12:OH</i>		<i>Et-2E,4Z-DD</i>	
		I	II	I	II
Females	virgin	0.28±0.10 a	0.36±0.10 a	1.56±0.34 a	1.46±0.20 a
	mated	0.36±0.12 a	0.28±0.10 a	1.68±0.26 a	1.58±0.16 a
Males	virgin	3.12±0.24 b	3.22±0.38 a	1.74±0.18 a	1.72±0.30 a
	mated	2.96±0.30 b	3.02±0.34 a	1.72±0.20 a	1.60±0.24 a

Different letters on the same column indicate significant differences (Duncan test; P=0.05).




Figure 3. Mean EAG (mV) dose-response curves of *C. pomonella* (n=10) virgin adults (walnut population) to *E8E10-12:OH* alone and during saturation (DS-EAG) with *Et-2E,4Z-DD*. Vertical bars represent S.D. (two-sided).




Figure 4. Mean EAG (mV) dose-response curves of *C. pomonella* (n=10) virgin adults (walnut population) to *Et-2E,4Z-DD* alone and during saturation (DS-EAG) with *E8E10-12:OH*. Vertical bars represent S.D. (two-sided).

when continuously exposed to *Et-2E,4Z-DD* (heptanal: male = 1.28±0.33 mV; female = 0.98±0.28 mV; Z3-hexen-1-ol: male = 1.11±0.38 mV; female = 1.06±0.22 mV).

Out of 123 attempts, complete recordings (n=20) from single antennal olfactory cells of males showed the presence of different cells (A, B, C) responding only to *E8E10-12:OH* (n=9) or *Et-2E,4Z-DD* (n=3) but also to the two compounds (n=8) (figure 5; table 3). In females,

out of 115 attempts, complete recordings (n=10) revealed the presence of cells (D, E, F) responding to *E8E10-12:OH* (n=2), *Et-2E,4Z-DD* (n=4) and to both compounds (n=4) (table 3).

In all recordings, in response to the adequate stimulus (*E8E10-12:OH*, *Et-2E,4Z-DD* or both compounds separately applied) according to the cell type, spike frequencies significantly increased with respect to control stimulus (mineral oil) (table 3). Significant differences

between the response to the mineral oil and the cell resting activity were not detected.

Field

All doses of Et-2E,4Z-DD showed an attractant activity on both sexes of CM. During 2003, at the highest dose (40 mg) of Et-2E,4Z-DD male and female catches were significantly lower. In 2004, male and female catches did not vary according to the dose (table 4).

Adults of CM were attracted by Et-2E,4Z-DD (0.1, 1.0 mg) in a significant lower number than the corresponding doses of pheromone. A similar result was observed when only male catches were considered. During 2003, male attraction was significantly reduced by addition of Et-2E,4Z-DD (0.1 mg) to the highest dose (1.0 mg) of pheromone and slightly increased when added to the lowest one (0.1 mg).

On the contrary, females were significantly attracted by 0.1 and 1.0 mg doses of pear ester; catches decreased in the presence of pheromone. Some females were caught by pheromone traps in a not significant number

than control (table 4).

During 2004, when the CM population density was low, significant differences among pheromone and binary blends did not emerge (table 4).




Figure 5. Spikes recorded on stimulation of male antennal olfactory cells of *C. pomonella* with solvent (a: mineral oil, 10 µl), Et-2E,4Z-DD (b: 1 µg) and E8E10-12:OH (c: 1 µg). Horizontal line above the recordings indicates the duration of stimulation (500 ms).

Table 3. Mean spike frequency (spikes/s±S.D.) recorded (surface contact technique) from *C. pomonella* male (A: n=9; B: n=3; C: n=8) and female (D: n=3; E: n=5; F: n=5) olfactory cells on stimulation with E8E10-12:OH and Et-2E,4Z-DD (1 µg) in mineral oil solution (10 µl). Mineral oil (10 µl) was used as control stimulus.

Cell	Spikes/s ± S. D.			
	Resting activity	Mineral oil	E8E10-12:OH	Et-2E,4Z-DD
Males				
A	7.89±1.45 a	8.56±1.33 a	47.78±5.07 b	9.22±1.64 a
B	9.33±1.53 a	8.67±1.15 a	9.67±1.53 a	28.67±2.52 b
C	8.13±1.25 a	8.75±1.04 a	43.88±3.72 c	31.25±3.06 b
Females				
D	6.67±1.53 a	7.67±1.53 a	23.33±4.04 b	8.67±2.08 a
E	9.20±1.43 a	8.60±1.14 a	10.20±1.64 a	32.80±4.38 b
F	7.40±1.14 a	9.60±2.30 a	26.60±4.34 b	28.60±4.22 b

Different letters on the same line indicate significant differences (LSD test; P=0.05).

Table 4. Field trapping of *C. pomonella* (no. adults/trap) in Colletorto (CB), Molise region, Italy (3 replicates; 21.V-20.IX.03; 20.V-25.VIII.04).

Compounds and blends	Males		Females		Total	
	2003	2004	2003	2004	2003	2004
E8E10-12:OH (0.1 mg)	67.0±5.0 c	40.0±2.0 a	1.3±1.2 d	0.3±0.6 c	68.3±6.0 c	40.3±1.5 a
E8E10-12:OH (1.0 mg)	110.7±6.4 a	27.7±7.4 b	2.3±0.6 cd	0.3±0.6 bc	113.0±6.1 a	28.0±7.9 b
E8E10-12:OH (0.1 mg) Et-2E,4Z-DD (0.1 mg)	82.0±9.4 b	29.3±4.7 ab	7.3±1.2 b	2.3±2.5 abc	89.3±9.8 b	32.0±4.6 ab
E8E10-12:OH (1.0 mg) Et-2E,4Z-DD (0.1 mg)	56.7±6.1 c	36.3±6.1 ab	4.7±2.5 bc	1.0±1.0 abc	61.3±8.6 c	37.3±7.1 ab
Et-2E,4Z-DD (0.1 mg)	21.3±5.1 e	2.7±1.2 d	15.7±3.1 a	1.3±2.3 abc	37.0±4.4 d	2.7±1.2 de
Et-2E,4Z-DD (1.0 mg)	25.3±5.1 de	1.3±1.5 de	13.7±5.0 a	1.3±1.5 abc	39.0±7.8 d	2.7±1.5 e
Et-2E,4Z-DD (3.0 mg)	27.7±0.6 de	10.3±8.1 c	13.3±6.1 a	2.7±1.2 ab	41.0±6.2 d	13.0±7.2 c
Et-2E,4Z-DD (10 mg)	30.0±5.2 d	5.0±2.6 cd	5.0±1.0 bc	3.3±2.5 a	35.0±6.1 d	8.3±3.5 cd
Et-2E,4Z-DD (40 mg)	11.0±3.0 f	3.0±1.0 d	4.7±0.6 bc	0.3±0.6 bc	15.7±2.5 e	3.3±1.5 e
Control	0.3±0.6 g	0.0±0.0 e	1.0±1.0 d	0.0±0.0 c	1.3±1.5 f	0.0±0.0 f

Means followed by the same letter are not significantly different at P=0.05 by Duncan test following ANOVA of $\sqrt{x+0.5}$ transformed data.

Discussion and conclusions

Sex, mating status and host-plant did not significantly influence the electrophysiological responses to pear ester, although mated females showed a higher response to this compound. Obviously, females were highly less sensitive to codlemone, confirming the results of previous studies and their ability in the auto-detection of sex pheromone (De Cristofaro *et al.*, 2002).

As a general tendency, adults of the walnut populations showed, on the average, a higher sensitivity to both compounds. The larval development on different host plants, affecting the female oviposition behaviour (Ansebo and Witzgall, 2002), could also be related to the perception of substances other than plant volatiles (Lombarkia and Derridj, 2002).

The reduced insecticides susceptibility of the tested population did not affect the electrophysiological responses both to the pear ester and codlemone, confirming the results related to the pheromone perception reported by Beslay *et al.* (2000).

SCR (directly) and DS-EAG (indirectly) results clearly showed the presence of cells responding both to codlemone and pear ester, at comparable doses, in CM male and female.

Olfactory cells sensitive only to a single pheromone component or to a plant volatile have been reported by several authors (i.e. Anderson *et al.*, 1995). Receptor neurons responding to more than one pheromone compound, or to both synergists and antagonists, have been also found in different Lepidoptera species (i.e. Löfqvist *et al.*, 1990; Renou and Lucas, 1994; Hansson *et al.*, 1995; Monti *et al.*, 1995; Wu *et al.*, 1995; Cossè *et al.*, 1998; Bäckman *et al.*, 2000). Cells responding to more than one plant volatile are well known (i.e. Den Otter *et al.*, 1996); most of them respond strongly to one compound and only weakly to a few other related constituents (i.e. Ulland *et al.*, 2003).

Cells responding to pheromone components and to high doses of host-plant volatiles have been already found on male antennae of *Cydia fagiglandana* (Zeller) (Lepidoptera Tortricidae) (De Cristofaro, 1995; Den Otter *et al.*, 1996; De Cristofaro *et al.*, 1997), *Utetheisa ornatrix* (L.) (Lepidoptera Arctiidae) (Bogner *et al.*, 1992), *Agrotis segetum* (Denis & Schiffmüller) (Hansson, 1995) and other insects (De Cristofaro *et al.*, 2004). A male receptor neuron responding to a blend of the main pheromone component (Z11-hexadecenal) with either linalool or Z3-6:OH, but not to the plant compounds alone, has been reported in *Helicoverpa zea* (Boddie) (Ochieng *et al.*, 2002). The biological meaning of these electrophysiological evidences is still not completely understood, since a separate peripheral perception mechanism and response processing of the two categories of volatile substances have been clearly shown (i.e. Mustaparta, 2002). Interestingly, our recordings allow to hypothesize that in CM the behavioural response to the plant volatile Et-2E,4Z-DD and to the main component of the sex pheromone E8E10-

12:OH is partly (at least) mediated through a common sensory channel.

The occurrence of cells sensitive to the sex pheromone main component and pear ester in male and female CM (De Cristofaro *et al.*, 2002; present paper), probably housed in sensilla trichoidea or auricillica, is particularly relevant. The absence of specificity of this receptor may be related to a similar olfactory affinity of the two molecules. In this sense pear ester acts as a codlemone mimic attracting CM and other tortricid moths sharing the same compound in the pheromone blend, like *C. fagiglandana*, *Cydia splendana* (Hübner), *H. nubiferana* (Haworth) (Schmidt *et al.*, 2004). This finding could explain the pheromonal potency of Et-2E,4Z-DD (Light *et al.*, 2001; Ioriatti *et al.*, 2003b) and the peripheral interactions in the perception of the two compounds. A synergistic effect on male attraction was not observed by adding Et-2E,4Z-DD to the pheromone. On the other hand, the frequently observed codlemone male capture efficiency reduction when mixed to the pear ester, like in the first year of our field studies, could be justified by a possible competition of the two substances for the same receptor. When CM population density is low the combined effect of the two substances on field catches is hardly distinguishable.

CM males were more sensitive than females to Et-2E,4Z-DD. An EAG study on *Manduca sexta* (L.) suggested that female antennae have a higher number of olfactory receptor neurons sensitive to non-pheromone stimuli than males (Kalinová and Hansson, 2001). However, the higher number of pheromone olfactory sensilla together with the presence of cells responding to Et-2E,4Z-DD and pheromone components could explain the higher EAG response of CM males to this compound. A higher EAG response of males to plant volatiles has already been reported in other Lepidoptera (i.e. De Cristofaro *et al.*, 1997; 1998; 1999; 2000; Vitagliano *et al.*, 2004).

All these observations have to be highly considered when setting up mixtures of E8E10-12:OH and Et-2E,4Z-DD to monitor different CM populations. As a general consequence, it is necessary to pay more attention when setting up mixed blends of pheromone components and plant volatiles to monitor or control insect pest populations.

Acknowledgements

Research supported by a grant of the Trento Province (Italy), BIOINNOVA project. The authors are indebted to Dr Bill Lingren of Trécé for the cooperation in supplying traps and dispensers, and to Drs Lucia Maiuro, Centro Servizi Interdipartimentali di Microscopia (CSIM), University of Molise, for the SEM observations. The first author thanks Dr César Gemenó, Centre UdL-IRTA de R+D, Àrea de Protecció de Conreus, Lleida, Spain, for useful advice and discussion about DS-EAG experiments.

References

- ANDERSON P., HANSSON B. S., LÖFQVIST J., 1995.- Plant-odor-specific receptor neurons on the antennae of female and male *Spodoptera littoralis*.- *Physiological Entomology*, 20: 189-198.
- ANSEBO L., WITZGALL P., 2002.- Codling moth oviposition response to plant odors, p. 101. In: *Book of Abstracts of the IOBC wprs Working Group Meeting "Pheromones and Other Semiochemicals in Integrated Production"*, Erice (TP), Italy, 22-27 September 2002.
- BÄCKMAN A. C., ANDERSON P., BENGSSON M., LÖFQVIST J., UNELIUS C. R., WITZGALL P., 2000.- Antennal response of codling moth males, *Cydia pomonella* L. (Lepidoptera: Tortricidae), to the geometric isomers of codlemone and codlemone acetate.- *Journal of Comparative Physiology A*, 186: 513-519.
- BAKER T. C., ROELOFS W. L., 1976.- Electroantennogram response of the male moth, *Argyrotaenia velutinana* to mixtures of sex pheromone components of the female.- *Journal of Insect Physiology*, 22: 1357-1364.
- BESLAY D., FRÉROT B., MALOSSE C., RENOU M., BOUVIER J. C., SAUPHANOR B., 2000.- No pleiotropic effect of insecticide resistance on sexual communication in *Cydia pomonella*, p. 0640. In: *Proceedings of XXI International Congress of Entomology*, Foz do Iguassu, Embrapa Soja, Londrina, Brazil, 20-26 August 2000.
- BOGNER F., GRANT A. J., O'CONNELL R. J., 1992.- A potential kairomone stimulates pheromone-responsive receptor neurons in *Utetheisa ornatrix* (Lepidoptera: Arctiidae).- *Journal of Chemical Ecology*, 18 (3): 427-439.
- CORACINI M., BENGSSON M., LIBLIKAS I., WITZGALL P., 2004.- Attraction of codling moth males to apple volatiles.- *Entomologia Experimentalis et Applicata*, 110: 1-10.
- COSSÉ A. A., TODD J. L., BAKER T. C., 1998.- Neurons discovered in male *Helicoverpa zea* antennae that correlate with pheromone-mediated attraction and interspecific antagonism.- *Journal of Comparative Physiology A*, 182: 585-594.
- DE CRISTOFARO A., 1995.- Ruolo dei semiochimici nelle interrelazioni castagno-tortrici: studi elettrofisiologici e comportamentali.- *Ph. D. Thesis*, Italian National Libraries of Rome and Florence: 1-80.
- DE CRISTOFARO A., ROTUNDO G., GERMINARA G. S., 1997.- Risposte elettrofisiologiche e comportamentali delle tortrici delle castagne (Lep.: Tortricidae) agli attrattivi sessuali ed alla pianta ospite, pp. 493-513. In: *Atti Convegno Nazionale sul Castagno*, Cison di Valmarino (TV), Italy, 23-25 October 1997.
- DE CRISTOFARO A., ROTUNDO G., GERMINARA G. S., 1998.- I semiochimici di *Ephestia kuehniella* Zeller (Lepidoptera: Phycitidae): analisi elettroantennografica e risultati preliminari di osservazioni comportamentali, pp. 143-156. In: *Atti VI Simposio Difesa Antiparassitaria nelle Industrie Alimentari e sulla Protezione degli Alimenti*, Piacenza, Italy, 24-26 September 1997.
- DE CRISTOFARO A., FONZO V., ANFORA G., ARNONE S., CRISTOFARO M., 1999.- Electrophysiological and olfactory responses of *Phthorimaea operculella* (Zeller) adults to volatile potato semiochemicals, pp. 217-218. In: *Proceedings XIV Triennial Conference of the European Association for Potato Research, EAPR '99*, Sorrento (NA), Italy, 2-7 May 1999.
- DE CRISTOFARO A., ROTUNDO G., GERMINARA G. S., 2000.- Electrophysiological and olfactory responses of *Ephestia kuehniella* Zeller adults to cereal semiochemicals.- *IOBC/WPRS Bulletin*, 23 (10): 189-194.
- DE CRISTOFARO A., IORIATTI C., MOLINARI F., PASQUALINI E., ROTUNDO G., 2002.- Electrophysiological responses of codling moth populations from different host plants to (E,E)-8,10-dodecadien-1-ol and ethyl (2E,4Z)-2,4-decadienoate and interactions in perception of the two attractants, pp. 108-109. In: *Book of Abstracts of the OILB wprs Working Group Meeting "Pheromones and Other Semiochemicals in Integrated Production"*, Erice (TP), Italy, 22-27 September 2002.
- DE CRISTOFARO A., ANFORA G., IORIATTI C., GERMINARA G. S., ROTUNDO G., 2004.- Occurrence of olfactory cells responding to pheromone components and plant volatile compounds in different species of Lepidoptera and Diptera: possible implications on semiochemical applications, pp. 83-84. In: *Book of Abstracts of the OILB VI International Conference on Integrated Fruit Production*, Baselga di Pinè (TN), Italy, 26-30 September 2004.
- DEN OTTER C. J., BEHAN M., MAES F. W., 1980.- Single cell responses in female *Pieris brassicae* (Lepidoptera: Pieridae) to plant volatiles and conspecific egg odours.- *Journal of Insect Physiology*, 26: 465-472.
- DEN OTTER C. J., TCHICAYA T., VAN DEN BERG, M. J., 1988.- Olfactory sensitivity of five species of tsetse flies (*Glossina* spp.) to 1-octen-3-ol, 4-heptanone, 3-nonanone and acetone.- *Insect Science and Applications*, 9: 213-218.
- DEN OTTER C. J., TCHICAYA T., SCHUTTE A. M., 1991.- Effects of age, sex and hunger on the antennal olfactory sensitivity of tsetse flies.- *Physiological Entomology*, 16: 173-182.
- DEN OTTER C. J., DE CRISTOFARO A., VOSKAMP K. E., ROTUNDO G., 1996.- Electrophysiological and behavioural responses of chestnut moths, *Cydia fagiglandana* and *C. splendana* (Lep., Tortricidae), to sex attractants and odours of host plant.- *Journal of Applied Entomology*, 120: 413-421.
- DICKENS J. C., JANG E. B., LIGHT D. M., ALFORD A. R., 1990.- Enhancement of insect pheromone response by green leaf volatiles.- *Naturwissenschaften*, 77: 29-31.
- FEYEREISEN R., 1999.- Insect P450 enzymes.- *Annual Review of Entomology*, 44: 507-533.
- HANSSON B. S., 1995.- Olfaction in Lepidoptera.- *Experientia*, 51: 1003-1027.
- HANSSON B. S., BLACKWELL A., HALLBERG E., LÖFQVIST J., 1995.- Physiological and morphological characteristics of the sex pheromone detecting system in male corn stem borers, *Chilo partellus* (Lepidoptera: Pyralidae).- *Journal of Insect Physiology*, 41: 171-178.
- IORIATTI C., BOUVIER J. C., 2000.- La resistenza agli insetticidi: il caso della Carpocapsa (*Cydia pomonella* L.).- *Informatore fitopatologico*, 50 (9): 5-10.
- IORIATTI C., BOUVIER J. C., BUTTURINI A., CORNALE R., TISO R., 2003a.- Carpocapsa: la situazione della resistenza ad azinphos methyl e diflubenzuron in Trentino ed Emilia-Romagna.- *Informatore fitopatologico*, 53 (1): 53-59.
- IORIATTI C., DE CRISTOFARO A., MOLINARI F., PASQUALINI E., SCHMIDT S., ESPINHA I., 2003b.- The plant volatile attractant (2E, 4Z)-2,4-ethyl-decadienoate for Codling Moth monitoring.- *Bollettino di Zoologia agraria e di Bachicoltura*, Ser. II, 35 (2): 127-135.
- KALINOVÁ B., HANSSON B. S., 2001.- Characterisation of antennal olfactory receptor cells in the male moth *Manduca sexta* sensitive to non-pheromone odours, p. 16. In: *Book of Abstracts of VII European Symposium for Insect Taste and Olfaction*. Villasimius (CA), Sardinia, Italy, 22-28 September 2001.
- KNIGHT A. L., LIGHT D. M., 2001.- Attractants from 'Bartlett' pear for codling moth, *Cydia pomonella* (L.), larvae.- *Naturwissenschaften*, 88 (8): 339-342.
- LIGHT D. M., FLATH R. A., BUTTERY R. G., ZALOM F. G., RICE R. E., DICKENS J. C., JANG E. B., 1993.- Host-plant green-leaf volatiles synergize the synthetic sex pheromones of the

- corn earworm and codling moth (Lepidoptera).- *Chemoecology*, 4: 145-152.
- LIGHT D. M., KNIGHT A. L., HENRICK C. A., RAJAPASKA D., LINGREN B., DICKENS J. C., REYNOLDS K. M., BUTTERY R. G., MERRIL G., ROITMAN J., CAMPBELL B. C., 2001.- A pear-derived kairomone with pheromonal potency that attracts male and female codling moth, *Cydia pomonella* (L.).- *Naturwissenschaften*, 88 (8): 333-338.
- LÖFQVIST C., HANSSON B. S., DIJKERMAN H. J., HERREBOUT W. M., 1990.- Behavioural and electrophysiological activity of unsaturated analogues of the pheromone tetradecyl acetate of the small ermine moth *Yponomeuta rorellus*.- *Physiological Entomology*, 15: 47-54.
- LOMBARKIA N., DERRIDJ S., 2002.- Incidence of apple fruit and leaf surface metabolites on *Cydia pomonella* oviposition.- *Entomologia Experimentalis et Applicata*, 104: 79-87.
- MILLER J. R., MORI K., ROELOFS W. L., 1977.- Gipsy moth field trapping and electroantennogram studies with pheromone enantiomers.- *Journal of Insect Physiology*, 1447-1453.
- MONTI L., LALANNE-CASSOU B., LUCAS P., MALOSSE C., SILVAIN J. F., 1995.- Differences in sex pheromone communication systems of closely related species *Spodoptera latifascia* (Walker) and *S. descoinsi* Lalanne-Cassou & Silvain (Lepidoptera: Noctuidae).- *Journal of Chemical Ecology*, 21: 641-660.
- MUSTAPARTA H., 2002.- Encoding of plant odour information in insects: peripheral and central mechanisms.- *Entomologia Experimentalis et Applicata*, 104: 1-13.
- NISHINO C., MANABE S., 1984.- Differential saturation electroantennogram of germacrene-D, a sex pheromone mimic of the american cockroach.- *Journal of Pest Science*, 9: 125-130.
- OCHIENG S. A., PARK K. C., BAKER T. C., 2002.- Host plant volatiles synergize responses of sex pheromone-specific olfactory receptor neurons in male *Helicoverpa zea*.- *Journal of Comparative Physiology A*, 188: 325-333.
- RENOU M., LUCAS P., 1994.- Sex pheromone reception in *Mamestra brassicae* L. (Lepidoptera): response of olfactory receptor neurones to minor components of the pheromone blend.- *Journal of Insect Physiology*, 40: 75-85.
- ROTUNDO G., TREMBLAY E., 1993.- Electroantennographic responses of chestnut moths (Lepidoptera: Tortricidae) and their parasitoid *Ascogaster quadridentatus* Wesmael (Hymenoptera: Braconidae) to volatiles from chestnut (*Castanea sativa* Miller) leaves.- *Redia*, 76 (2): 361-373.
- SAUPHANOR B., CUANY A., BOUVIER J. C., BROSSE V., AMICHOT M., BERGÉ J. B., 1997.- Mechanism of resistance to deltamethrin in *Cydia pomonella* L. (Lepidoptera: Tortricidae).- *Pesticide Biochemistry and Physiology*, 58:109-117.
- SCHMIDT S., ANFORA G., DE CRISTOFARO A., IORIATTI C., 2004.- Tortricid species caught by codling moth kairomone ethyl (2E, 4Z)-2,4-decadienoate: trapping experiments and electrophysiological responses, p.153. In: *Book of Abstracts of VI International Conference on Integrated Fruit Production, OILB wprs Working Groups Meeting*, Baselga di Piné (TN), Italy, 26-30 September 2004.
- ULLAND S., MEADOW R., BORG-KARLSON A. K., BONES A., MUSTAPARTA H., 2003.- Olfactory receptor neurones in *Mamestra brassicae* specified for volatiles released by host plants of related species, p. 81. In: *Book of Abstracts of VIII ESITO Meeting*, Harstad, Norway, 2-7 July 2004.
- VAN DER PERS J. N. C., MINKS A. K., 1993.- Pheromone monitoring in the field using single sensillum recording.- *Entomologia Experimentalis et Applicata*, 68: 237-245.
- VITAGLIANO S., ANFORA G., TASIN M., GERMINARA G. S., IORIATTI C., ROTUNDO G., DE CRISTOFARO A., 2004.- Electrophysiological and olfactory responses of *Lobesia botrana* (Den. et Schiff.) (Lepidoptera Tortricidae) to odours of host plant. p. 144. In: *Book of Abstracts of VI International Conference on Integrated Fruit Production, OILB wprs Working Groups Meeting*, Baselga di Piné (TN), Italy, 26-30 September 2004.
- WU W.-Q., HANSSON B. S., LÖFSTEDT C., 1995.- Electrophysiological and behavioral evidence for a fourth sex pheromone component in the turnip moth, *Agrotis segetum*.- *Physiological Entomology*, 20: 81-92.
- YANG Z., BENGTSSON M., WITZGALL P., 2004.- Host plant volatiles synergize response to sex pheromone in codling moth, *Cydia pomonella*.- *Journal of Chemical Ecology*, 30 (3): 619-629.

Corresponding author: Antonio DE CRISTOFARO, (e-mail: decrist@unimol.it) Dipartimento di Scienze Animali, Vegetali e dell'Ambiente, Università degli Studi del Molise, via De Sanctis s.n.c., 86100 Campobasso, Italy.

Received October 5, 2004. Accepted November 4, 2004.